

- 32** Diabase, quartz diabase, gabbro, diorite
- 28** ET-THEN GROUP: conglomerate, sandstone, quartzite
- 27** Granite, granodiorite, diorite, syenite, porphyritic (quartz and feldspar) felsites
- 19** GREAT SLAVE GROUP (Upper Part)-STARK, TOCHATWI, and PEARSON FORMATIONS: dolomite, limestone, shale, argillite, sandstone, andesite, basalt, trachyte
- 17** NONACHO GROUP: arkose, quartzite, slate, greywacke, conglomerate; minor volcanic rocks, phyllite and schist
- 14** GREAT SLAVE GROUP (Lower Part)-SOSAN, KAHOCHELLA, and PETHI FORMATIONS: conglomerate, sandstone, quartzite, shale, slate, argillite, arkose, limestone, dolomite, tuff, breccia, agglomerate, andesite, iron formation
- 13** Granodiorite, granite, quartz diorite, syenite; includes some gneissic rocks; 13c, may include Proterozoic granitic rocks
- 12** Impure and gneissic granitic rocks, sedimentary and volcanic schist and gneiss, mixed sedimentary and granitic schist and gneiss, migmatite, mylonite. May include minor Proterozoic rocks
- 7** TAZIN GROUP: andesite, dacite, rhyolite, tuff, agglomerate, greywacke, arkose, quartzite, argillite, slate, limestone, quartz-mica schist
- 4** Hornfels, nodular quartz-biotite schist, quartzite, siliceous argillite, minor paragneiss and migmatite
- 1** Basalt, andesite, tuff, agglomerate, chert, breccia; derived schist and gneiss; minor dacite and rhyolite

Geological contact (defined or approximate, assumed) .....  
 Fault .....  
 Limit of geological mapping .....  
 Sand and silt, or drift covered areas .....

Geology derived from the 1:1,267,200  
 Geological Map of the District  
 of Mackenzie, Northwest Territories  
 Map 1055A

Geological cartography by the Geological Survey of Canada


Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Geochemical Symbols and Data Presentation


The concentration of an element at a sample site is graphically presented by using one of 15 symbols, if a sample was collected but no relevant data is available a dot is plotted. The symbols are divided into 2 groups of 7 respectively, and one additional symbol. The first 7 increase in size from a small cross to a large open circle. The eighth symbol, the additional one, consists of the large open circle with a dot at its centre. In the last 7 symbols the dot becomes progressively larger until a solid black circle is reached, a circumscribed square in solid black is used for the last symbols. Thus the overall impression is of increasing size and blackness as samples display from low to high elemental content.

The data distributions are depicted using both a cumulative probability plot and a histogram. It should be noted that if the survey covers several map sheets the distribution displays are for the total survey data, not the single sheet being studied. The data intervals used for the histogram are semi-logarithmic, e.g. 1, 2, 5, 10, 20, 50, 100 etc., due to the apparent log-normal characteristic of most trace element distributions. The symbols cover 5 decades, e.g. 1 ppm to 10%, using the above data intervals. As this map is part of a Canada wide series these arbitrary intervals have been chosen to preserve some long-term continuity, however, based on experience at the Geological Survey, we believe them to be an appropriate compromise.

For any particular element the eighth, central, symbol is ascribed to the histogram mode. The median of the data distribution, as defined by the 50% (0.5) point on the cumulative probability plot, usually falls within the mode interval of the histogram. The symbol (+) and (x) are respectively used to display concentrations below and at the analytical detection limit. To illustrate the use of the symbols an example is given below. If the actual data distribution does not require all 15 symbols some are selectively dropped so that maximum overall graphic impact is maintained.



This document was produced by scanning the original publication. Ce document est le produit d'une numérisation par balayage de la publication originale.



MANGANESE IN LAKE SEDIMENTS

URANIUM RECONNAISSANCE PROGRAM  
 NATIONAL GEOCHEMICAL RECONNAISSANCE

Scale 1:250,000



Kilometres 0 6 12 18 Kilometres

Miles 0 4 8 Miles

Universal Transverse Mercator Projection

© Crown Copyrights reserved

75L	75K	75J
O.F. 326		
75E	75F	75G
O.F. 325		
75D	75C	75B
O.F. 324		



OPEN FILE 324

by E.H.W. Hornbrook, R.G. Garrett and J.J. Lynch

Geological Survey of Canada

Geochemistry and Federal-Provincial coordination by E.H.W. Hornbrook  
 Analytical chemistry by J.J. Lynch  
 Data monitoring and compilation by R.G. Garrett and N.G. Lund  
 Cartography and base compilation by Geological Cartography Section

Base-map assembled by the Geological Cartography Unit from maps published at the same scale by the Army Survey Establishment, R.C.E. in 1963

Copies of the topographical maps covering this map-area may be obtained from the Canada Map Office

Mean magnetic declination 1976, 26°04.5' East, decreasing 7.6' annually. Readings vary from 24°40.0' in the SE corner to 27°30.6' in the NW corner of the map-area

Elevations in feet above mean sea-level

Contractors

Sample collection by Trigg, Woollett & Associates Ltd.  
 Chemical analyses by Chemex Labs. Ltd.

Chemical analyses by Chemex Labs Ltd., and Atomic Energy of Canada Ltd., Commercial Products Division

This map forms one a series of 42 sheets released under Geological Survey of Canada Open File 324. The open file consists of data for 12 elements, percent loss on ignition and sample site location, each variable requiring 3 sheets for the total survey area.

The data is also available in digital form from the Computer Science Centre of the Department of Energy, Mines and Resources: For further information please contact:

The Director,  
 Computer Science Centre,  
 Department of Energy, Mines and Resources,  
 Ottawa, Ontario K1A 0E4.

OPEN FILE 324  
 MANGANESE  
 NATIONAL GEOCHEMICAL RECONNAISSANCE  
 NORTHWEST TERRITORIES, 1975