



PRELIMINARY SERIES



- LEGEND
- Note: Weighted legend blocks indicate map-units that appear on this map
- QUATERNARY
- 18 Drift
- ORDOVICIAN
- 17 Limestone; minor shale
- APHEBIAN (?)
- 16 Diabase
- APHEBIAN
- 15 Ultrabasic rocks
 - 14 Hypersthene granite
 - 13 Biotite granite
 - 12 Gneissic granite
 - 11 Migmatite
 - 10 Quartzite
- PROTEROZOIC
- 9 Crystalline limestone
 - 8 Hornblende-pyroxene gneiss
 - 7 Garnet-biotite-quartz-feldspar gneiss
 - 6 Rusty paragneiss; commonly graphite-bearing
 - 5 Sillimanite gneiss and schist
 - 4 Biotite-quartz-feldspar gneiss
 - 3 Quartz-feldspar gneiss; 3a, biotite-quartz-feldspar gneiss
 - 2 Biotite gneiss
 - 1 Volcanic rocks
- Geological boundary (assumed, gradational)
- Bouding (horizontal, inclined, vertical, overturned, dip unknown) + / / /
- Foliation (horizontal, inclined, vertical, dip unknown) + / / /
- Lineation, axes of minor folds (horizontal, inclined, vertical) / / /
- Geology by R. G. Blackadar. Field observations by R. G. Blackadar, 1949; W. L. Davison, 1949-1951; T. O. Fortier, 1949-1952; R. N. McNeely, 1965; P. H. Smith, 1965; F. C. Taylor, 1965
- Base-map cartography by the Geological Survey of Canada, 1967 from 1/250,000 scale map published by Surveys and Mapping Branch in 1960, 1961, 1963, 1964
- All elevations in feet above mean sea-level

MAP 18-1966
PAPER 66-47
GEOLOGY
FROBISHER BAY
DISTRICT OF FRANKLIN



Mean magnetic declination, 43°49' West, decreasing 12.6' annually. Readings vary from 41°43' in the SW corner to 46°34' in the NE corner of the map-area

