

LEGEND

Note: this legend is common to maps 1282A, 1283A and 1284A

NORTHERN PART

SOUTHERN PART

QUATERNARY

26 Unconsolidated glacial and alluvial deposits

CRETACEOUS AND TERTIARY (?)  
UPPER CRETACEOUS AND LATER (?)

22 MONSTER FORMATION: 22a, brown-weathering, thin-bedded, brown chert-grain sandstone, siltstone, shale, and fine chert-pebble conglomerate

20a Orange-to brown-weathering diorite and gabbro; altered equivalents; may be older than 20

TRIASSIC

16 Black-weathering, platy, black limy shale and limestone; thin bands of grey-to buff-weathering limestone

PERMIAN

15 TAHKANDIT FORMATION: white, light grey, and dark grey chert, cherty limestone, and limestone

CARBONIFEROUS TO PERMIAN

14 Buff-weathering, dark grey, thin- to medium-bedded limestone; minor black shale, chert, and chert-pebble conglomerate; 14a, dark shale, argillaceous limestone, and thin-bedded brown sandstone; minor chert-pebble conglomerate; 14b, black- and silvery-weathering shale and slate; minor platy, buff-weathering grey limestone, impure sandstone

DEVONIAN TO CARBONIFEROUS

13 Middle Devonian to Carboniferous: Black shale, argillite, and slate, black platy limestone, chert; minor chert-pebble conglomerate and quartzite; 13a, Nation River Formation: brown-weathering fine chert-pebble conglomerate and chert-grain sandstone may, in part, be younger Monster Formation (22)

DEVONIAN

LOWER MIDDLE DEVONIAN

11 Limestone, dark grey, brown and black, massive to thin-bedded, very fine grained, buff-grey-weathering

10 Limestone and dolomite, light grey and dark brownish grey, fine to medium grained, mostly alternating dark and light beds 2 to 5 feet thick

ORDOVICIAN AND SILURIAN

9 ROAD RIVER FORMATION: mainly interbedded black chert and black argillite, also grey-green, olive-green, and grey chert and grey-green argillite; minor quartzite, and chert-pebble conglomerate

8 Grey- and buff-weathering dolomite and limestone, mostly medium to thick bedded; minor platy black argillaceous limestone and dolomite (may include some 9, 10, and 11); 8a, grey- to dark grey-weathering, dark volcanic rocks many partly serpentinized, brown-weathering grey-green limy tuff and argillite, and thin-bedded brown limestone

CAMBRIAN

MIDDLE (?) AND UPPER CAMBRIAN

6 Buff, brown, and grey-weathering, thin- to medium-bedded limestone, and grey-weathering thin- to thick-bedded dolomite; minor brown and green shale and orange-weathering dolomite

CAMBRIAN (?)

5 Mainly brick-red, thick-bedded to massive sandstone and red to buff massive conglomerate; minor red shale; local andesitic or basaltic flows and sills

PROTEROZOIC

2 Orange-weathering, platy, grey-green dolomite, dark slate; minor phyllite and quartzite; 2a, pink-, orange- and grey-weathering dolomite, grey and maroon shale, white, green and mauve quartzite, minor conglomerate, mottled green and maroon shale and black limestone; 2b, buff and orange dolomite, dark shale; minor quartzite limestone and conglomerate; 2c, massive cherty and quartzose, grey dolomite; thin-bedded, buff-weathering, grey dolomite; minor black shale and white quartzite; 2d, buff-weathering dolomite-boulder conglomerate; 2e, dark shale and argillite, buff-weathering, grey siltstone; minor buff- to orange-weathering dolomite

1 Mainly dark grey, grey-green, and black, thin-bedded argillite, slate and phyllite; minor grey quartzite, orange-weathering dolomite, and conglomerate; 1a, grey-weathering, thinly laminated, silicified limestone

METAMORPHIC ROCKS SOUTHWEST OF TINTINA TRENCH

(occurs only on Map 1284A, Dawson)

E Reddish brown-weathering, dark green serpentinized ultrabasic rocks

D Fine- to medium-grained, granitic textured, quartz-biotite gneiss; minor quartzite, quartz-mica and biotite-chlorite schist, and quartz-feldspar pegmatite

C Dark weathering greenstone and banded amphibolite gneiss; minor chloritic quartz-mica schist, graphitic quartz-mica schist, quartzite, and limestone

B KLONDIKE "SCHIST": mainly buff weathering, light pale green quartz-muscovite-chlorite schist, and schistose, chloritic quartzite, with all intermediate rock types also present; minor silvery muscovite schist, fine-grained quartz-biotite gneiss, thinly laminated quartz-graphite-sericite schist and quartzite

A NASINA "SERIES": grey and grey-green/micaeous quartzite; dark grey, light grey and silvery quartz-mica schist; minor fine-grained quartz biotite gneiss, graphitic schist and quartz-muscovite-chlorite schist; Aa, higher rank metamorphic rocks with biotite and garnet; Ab, coarsely crystalline, whitish limestone

QUATERNARY

26 Unconsolidated glacial and alluvial deposits

TERTIARY

25 Quartz porphyry

24 Dark grey and brown andesite and basalt, commonly porphyritic; minor shale, sandstone, and conglomerate

23 Poorly consolidated, brown, buff, and grey, arkosic and micaceous sandstone, light and dark shale, poorly sorted conglomerate; minor lignite

CRETACEOUS

21 21a, fine- to coarse-grained, uneven textured, biotite granodiorite and biotite quartz monzonite; 21b, mainly hornblende and hornblende/biotite syenite, commonly porphyritic (potassium feldspar phenocrysts), uneven textured, mostly medium grained, locally fine or coarse grained; minor diorite

20 Orange- to brown-weathering diorite and gabbro; altered equivalents; 20a, may be older

19 Mottled green and maroon shale and brown-weathering, thin-bedded, brown siltstone, commonly limy

18 KENO HILL QUARTZITE: grey and blue-grey, massive quartzite; minor slate and phyllite, commonly graphitic, argillaceous quartzite; 18a, thin-bedded and phyllitic quartzite, graphitic and chloritic slate and phyllite, minor limestone and massive quartzite; 18b, as 18 but may be older

JURASSIC

17 LOWER SCHIST division: dark grey argillite, slate, and phyllite, commonly graphitic, thin-bedded dark grey quartzite, platy to phyllitic quartzite; minor phyllite and limy quartzite

TRIASSIC

16 Black-weathering, platy, black limy shale and limestone; thin bands of grey- to buff-weathering limestone

PERMIAN

15a Limestone with some chert

ORDOVICIAN AND SILURIAN

9 ROAD RIVER FORMATION: mainly interbedded black chert and black argillite, also grey-green, olive-green, and grey chert and grey-green argillite, minor quartzite, and chert-pebble conglomerate

PRECAMBRIAN AND/OR LATER

4 Dark brown- and green- to light grey-weathering dark green volcanic rocks, commonly with calcite filled vesicles, breccia, tuff, and agglomerate; minor interbedded shale, chert, siltstone, and limestone; 4a, dark brown to dark green-weathering dark green volcanic rocks, commonly with calcite-filled vesicles, breccia, tuff, and agglomerate. Interbedded with 2d and may be older; 4b, dark green, fine-grained andesite


PRECAMBRIAN AND/OR CAMBRIAN

3 Mainly buff-, brown-, and rusty-weathering, gritty quartzite, sandstone and quartz-pebble conglomerate; black, maroon and green shales, and slates; schistose quartzite, quartz chlorite schist, quartz-mica schist and phyllite; minor limestone and black chert; 3a, thin- to medium-bedded, dark grey limestone

MINERALS

Antimony ..... Sb Lead ..... Pb  
Asbestos ..... asb Silver ..... Ag  
Coal ..... C Tin ..... Sn  
Copper ..... Cu Tungsten ..... W  
Gold placer ..... Au Zinc ..... Zn  
Iron ..... Fe

Geological boundary (defined, approximate, assumed) .....  
Bedding, tops known (horizontal, inclined, vertical) .....  
Bedding, tops unknown (dip known) .....  
Bedding, estimated attitudes, may in part be of foliation:  
horizontal, inclined, vertical (dip: g, gentle; m, medium; s, steep) .....  
Foliation (horizontal, inclined, vertical) .....  
Fault (defined, approximate, assumed) .....  
Thrust fault (teeth in direction of dip: defined, approximate, assumed) .....  
Anticline (defined, approximate; arrow indicates plunge) .....  
Syncline (defined, approximate; arrow indicates plunge) .....  
Anticline, syncline (overturned) .....  
Fossil locality .....  
Mineral occurrence .....  
Goldfield .....  
Geology by L.H. Green and J.A. Roddick, 1961  
To accompany GSC Memoir 364 by L.H. Green  
Geological cartography by the Geological Survey of Canada  
Base-map at the same scale published by the Surveys and Mapping Branch in 1954, 1957 and 1958. Roads were revised by the Geological Survey of Canada for this edition  
Copies of the topographical edition of this map may be obtained from the Map Distribution Office, Department of Energy, Mines and Resources, Ottawa  
Any revisions or additional information known to the user would be welcomed by the Geological Survey of Canada  
The following names have not been approved by the Canadian Permanent Committee on Geographical Names: Mike Lake, Frank Lake, Peanut Lake, Marc Lake, Callison Lake, Joan Lake.




INDEX MAP


NOT TO BE TAKEN FROM LIBRARY  
NE PAS SORTIR DE LA BIBLIOTHÈQUE



GEOLOGICAL SURVEY OF CANADA  
DEPARTMENT OF ENERGY, MINES AND RESOURCES



Diagrammatic cross-sections along lines E-F-G and H




Printed by the Surveys and Mapping Branch

Published 1972  
Copies of this map may be obtained from the Geological Survey of Canada, Ottawa

MAP 1283A  
GEOLOGY  
LARSEN CREEK  
YUKON TERRITORY

Scale 1:250,000



Magnetic declination 1970 varies from 33°17' easterly at centre of west edge to 34°14' easterly at centre of east edge. Mean annual change decreasing 4.2'

Elevations in feet above mean sea-level

116 G-116 F (E 1/2)	116 H	106 E
116 B-116 C (E 1/2)	116 A	106 D
1284A	1283A	1282A
115 O-115 N (E 1/2)	115 P	105 M
711 A	1143 A	890 A

NATIONAL TOPOGRAPHIC SYSTEM REFERENCE AND INDEX TO GEOLOGICAL SURVEY OF CANADA MAPS

LARSEN CREEK  
YUKON TERRITORY



1283A