

BIBLIOTHÈQUE  
SERVICE DES MINES

CANADA  
DEPARTMENT OF MINES AND RESOURCES  
MINES AND GEOLOGY BRANCH

---

GEOLOGICAL SURVEY  
PAPER 45-15

**SHERRIDON**  
**MANITOBA**  
(Map and Descriptive Notes)

BY  
J. D. Bateman and J. M. Harrison



---

OTTAWA

1945

This document was produced  
by scanning the original publication.

Ce document est le produit d'une  
numérisation par balayage  
de la publication originale.

CANADA  
DEPARTMENT OF MINES AND RESOURCES  
MINES AND GEOLOGY BRANCH

---

GEOLOGICAL SURVEY

Paper 45-15

SHERRIDON MAP-AREA;

MANITOBA

(Descriptive Notes)

By

J.D. Bateman

---

OTTAWA, 1945

## CONTENTS

	Page
General geology .....	1
Pre-Sherridon formations .....	1
Sherridon group .....	2
Post-Sherridon formations .....	2
Cree Lake intrusive rocks .....	3
Granite .....	3
Structure .....	3
Mineral deposits .....	4
Types of sulphide deposits .....	5
Localization of sulphide deposits .....	5
Sherritt Gordon mine .....	6
Prospecting in the Kisseynew gneisses .....	6
Acknowledgments .....	7

### Illustration

Preliminary map --- Sherridon, Manitoba.

## SHERRIDON MAP-AREA, MANITOBA

### GENERAL GEOLOGY

Most of the rocks in the map-area are crystalline schists and gneisses, and form a part of the great area of Kisseynew gneisses north and east of Flin Flon in northern Manitoba and Saskatchewan. The gneisses are predominantly of sedimentary origin, having been derived from such rocks as quartzite, arkose, and greywacke, through the intimate intrusion of granitic and pegmatitic material and attendant solutions. The process of metamorphism is, in places, so far advanced that the resulting Kisseynew gneisses have lost their sedimentary structures and have become granitic in appearance. The different beds consist of various proportions of quartz, oligoclase, feldspar, biotite, and garnet, with accessory magnetite and hornblende. These minerals represent a common denominator resulting from the granitic metamorphism of various types of sediments.

Several narrow belts of hornblende-rich gneiss occur, mostly in the lower part of the Kisseynew group. These rocks consist of hornblende and andesine feldspar with more or less garnet and biotite. The hornblende gneisses represent metamorphosed basic igneous rocks and are probably mainly of volcanic origin, as relics of pillows were observed in several localities. In places they are massive and coarse-grained, resembling diorite, but more generally they possess a parallel foliation or stratiform structure that has been brought about by deformation during folding.

Throughout the Kisseynew gneisses it is commonly difficult to distinguish between true bedding and bedded-like or stratiform structures that have resulted from later foliation. In the Sherridon map-area the two structures are essentially parallel, the secondary foliation corresponding to the original bedding, even around the noses of plunging folds. It seems probable that recrystallization was controlled by primary bedding structures, for in places it has resulted in an intensification of stratification. On the accompanying map a symbol for foliation is used to indicate rock structures that are not certainly due to bedding, but that resemble bedding, and are described as stratiform. Such stratiform foliation is developed in some igneous as well as the sedimentary rocks and must, therefore, be a result of deformation.

### Pre-Sherridon Formations

The oldest rocks in the map-area lie in the centre of the Sherritt Gordon structure and pre-date the Sherridon group in which the orebodies occur. They consist of stratiform quartz-oligoclase-biotite gneiss (1), which weathers buff and contains many garnetiferous beds. The gneiss is a metamorphosed sedimentary formation, and is overlain by dark green hornblende-plagioclase gneiss (2) that is locally garnetiferous. The hornblende gneiss may be thinly foliated, but is generally massive and without visible structure. Part of it contains light-coloured lenticular fragments and was probably a volcanic breccia. There is some doubt as to whether the pre-Sherridon rocks belong to the Kisseynew group or whether they represent metamorphosed equivalents of older, Archaean formations, such as occur northeast of Flin Flon.



### Sherridon Group

The Sherridon group (3,4) lies in places upon Sherridon basic gneiss (2), but elsewhere upon still older sedimentary gneiss (1). This may be due to thinning out of the basic hornblende gneiss along its strike, or it may represent an erosional unconformity at the base of the Sherridon. The Sherridon consists of a group of distinctive white to grey quartzites interbedded with dark green to black hornblende-plagioclase gneisses that are metamorphosed volcanic flows. The different beds contain abundant quartz and various minor amounts of feldspar, biotite, hornblende, and garnet. They have a distinctive gneissic texture that is emphasized by the quartz, which stands out in relief on weathered surfaces. This feature alone serves to distinguish the Sherridon quartzites from any other rock in the area. Northeast of Cree Lake the Sherridon group contains many narrow lenticular bodies of anorthosite and gabbro that are not indicated on the map. The hornblende gneisses (4) are most abundant east of the Sherritt Gordon west orebody. Remnants of pillow structures were observed in them, but the rock is generally so thoroughly recrystallized that it resembles diorite, except that it may contain abundant garnet.

The Sherridon group probably represents the lower part of the Kisseynew gneisses. This is partly indicated by evidence outside of the map-area at Weldon Bay near the east end of Kisseynew Lake. Here a section overlying Archaean greenstones resembles in many respects the Sherridon group. The rocks at Weldon Bay, however, are much less metamorphosed, and are also strikingly similar to the Missi sediments near Flin Flon, even to containing scattered pebbles throughout the beds.

### Post-Sherridon Formations

The Sherridon group is overlain by dark green hornblende-rich gneiss (5) that is in sharp contact with the distinctive Sherridon quartzites. The hornblende gneiss is succeeded by widespread metamorphic types characteristic of the prevailing Kisseynew gneisses of the district. They vary from recognizable stratified rocks (6) to others of similar origin that have been so injected by granite and pegmatite that the bulk of the rock is intrusive and, for mapping purposes, is classified as "granitized" gneisses (7). In addition there are granitoid gneisses (8) that resemble granite or granodiorite in the hand specimen, but that have a stratiform structure similar to that of the bedded gneisses. The post-Sherridon gneisses include narrow belts of hornblende-rich gneiss (5), and these are most abundant near the lower part of the Kisseynew group. The prevailing rock of the post-Sherridon formations is a quartz-feldspar-biotite gneiss with accessory garnet and magnetite. Individual beds may be rich in one or other of these minerals. Magnetite and garnet are common in the granitic material associated with the "granitized" gneisses (7). The granitoid gneisses (8) differ from true granites in that they contain more albite and little potash feldspars (microcline and microperthite). They probably represent sedimentary gneisses in an advanced state of granitization.

Conglomerate (6a) was found in several localities in the Kisseynew gneisses. In the south part of the map-area are thick beds of conglomerate, the pebbles and boulders of which are almost all of quartz with disseminated magnetite. Near the outlet of Bartlett Lake and again on Big Island are relic conglomerates. At the latter locality the pebbles are mostly of quartz and granitic rocks. This conglomerate is not near the base of the Kisseynew complex and, so far as is known, is stratigraphically above the Sherridon formation. In this respect it would correspond with the conglomerate at Weldon Bay, which lies above what is believed to be the equivalent of the Sherridon.

### Cree Lake Intrusive Rocks

In the vicinity of Cree Lake and to the northeast the Sherridon and older formations are cut by intrusive rocks that form a related sequence from ultrabasic to granitic types. These include coarse-grained, black to rusty weathering pyroxenite (9) along the northeast shore of Cree Lake, and black to dark green massive peridotite and hornblende metagabbro (10) northeast of the lake. The latter rock locally carries abundant garnet. Between Cree and Found Lakes are bodies of anorthositic<sup>1</sup> and anorthositic gabbro (11) that consist

---

<sup>1</sup> C. H. Stockwell of the Geological Survey has examined this section and regards the anorthositic rocks as altered, limy sediments. Actually the present mineral assemblage of these rocks might be derived from the metamorphism of either impure limy quartzites or anorthositic intrusive rocks; and it is possible that either one or both types are present.

---

principally of plagioclase feldspar with more or less pyroxene, hornblende, carbonate, titanite, and scapolite. Narrow lenses of calcite and rusty weathering carbonate may be found in these rocks, and are believed to be the result of widespread carbonatization. The youngest member of the sequence is a buff-coloured oligoclase granite (12) that is locally sheared and rusty weathering. The Cree Lake intrusive rocks are considerably altered and the pyroxenite and anorthosite have a decayed appearance. The anorthosites, in particular, are foliated and it is probable that all were involved in at least part of the folding of the enclosing rocks. The relation of the Cree Lake intrusive rocks to the post-Sherridon Kisseynew gneisses is not known, and they may be older or younger.

### Granite

The younger granite and granite-gneiss is characteristically pink, and commonly has some faint stratiform structures that correspond in attitude to the bedding in the sedimentary gneisses. The granite is relatively undeformed, and was intruded after the main folding had occurred. Most of it forms small sills or larger bodies that conform to the trend of the structure in the sedimentary gneisses. Crosscutting contacts of the granites with the older rocks are rare and most of the granite may be said to form concordant intrusions. Pegmatite, presumably related to the granite, is widespread throughout the area. It is the ordinary variety of pink pegmatite, but may contain small masses of magnetite as much as a pound or more in weight.

### STRUCTURE

The rocks of the area are folded complexly as a result of several periods or stages of deformation. Sufficient detail work was done in the vicinity of Sherritt Gordon mine<sup>1</sup> to determine the essential

---

<sup>1</sup> Preliminary Map, Sherritt Gordon Mine Area; Geol. Surv., Canada, Paper 44-4 (1944).

---

structural features of that part of the area. Altogether some 3,000 determinations of strike and dip were made.

Throughout the map-area the prevailing dips of the formations are north and east, the fold axes being overturned respectively to the south and west. Bedding and foliation normally dip at angles of between 10 and 50 degrees. Several of the folds are isoclinal, but many are open and, as a result of overturning, steep dips are found on one limb with abnormally low dips on the other. Dips other than in the prevailing directions generally indicate the steeper limb of an open fold, which, as a result of overturning of the fold axis, becomes vertical and then reverses in dip as the nose of a plunging fold is approached.

Folded folds, such as the anticline encircling Found Lake, are probably common, but are difficult to recognize. Another stage of folding is represented in a great fan fold that comprises the Sherritt Gordon structure. The most common folding, however, is along north-south axes that swing easterly in the north part of the map-area. Few faults were recognized; the most important is one that extends north through Molly Lake and shows a relative displacement to the south on the west side.

The structural picture at Sherritt Gordon resembles a large drag-fold, but is rather the result of an uplift across a series of northerly trending folds overturned to the west. The uplift has effected a reversal of the prevailing northward plunge of the folds. It commences as a gentle flexure near Cold Lake and extends eastward between the Sherritt Gordon East and West orebodies, swinging north-eastward across Found Lake and northward across Cree Lake. North of Found Lake it steepens and is a definite anticline. It becomes the principal fold northeast of Cree Lake, where it supersedes the main syncline on the east, the latter dying out to the north. As a result the syncline plunges southward to near the south end of Molly Lake, where the prevailing northward plunge is resumed. The combined effect of these folds has resulted in an anticlinal dome or structural "high" at Sherritt Gordon, in which the older formations of the Kiseynew gneisses, and possibly pre-Kiseynew, Archaean rocks, have been exposed.

#### MINERAL DEPOSITS

Copper and zinc are the only metals that have been found in commercial quantities in the map-area. Gold occurrences, which are common in the Archaean greenstone areas to the south, are rare throughout the area of Kiseynew metamorphism. Following the initial development of the Sherritt Gordon deposit early in 1926 there was considerable prospecting activity in the district until 1930. Several additional copper-zinc occurrences were found during these years, but none except the Sherritt Gordon deposit proved to be of commercial size. Prospecting activities were limited after 1930, and there is little to add to the description of properties recorded in Part C of the Summary Report of the Geological Survey for that year. Subsequent exploration in the district has been carried out almost solely by Sherritt Gordon Mines, Limited. The Company increased the extent of its holdings, engaged in geological mapping, carried out an extensive geophysical survey of the northern claims, and undertook considerable exploratory diamond drilling. These efforts disclosed several barren pyrrhotite deposits. However, in December 1940, another copper-zinc orebody was found by drilling beneath Bob Lake, 3 miles northeast of the main orebodies.

As a result of geological investigations the Company, in 1943, acquired an additional 4,000 acres adjoining the main property on the east. Thirty-two claims northeast of Cree Lake are held by Mr. D. C. Barr of Sherridon, and a few claims lying near the West orebody are owned by the Hudson Bay Mining and Smelting Company; otherwise all

claims covering the Sherritt Gordon structure are owned or controlled by Sherritt Gordon Mines, Limited. The Company commenced production of copper in 1931, the concentrator operating until June 1932, when the plant was closed until August 1937. The copper concentrator has operated continuously since 1937 and, in 1942, a zinc concentrate was also produced. In 1942 the plant capacity was increased by 25 per cent, to 3,000 tons a day. Present production is well below plant capacity, and ore reserves although limited are sufficient for several years. The Bob Lake deposit is low-grade, and is not likely to be developed at current prices for copper.

#### Types of Sulphide Deposits

In contrast with the Flin Flon district, where the copper-zinc ores are associated with pyrite, copper and zinc mineralization in the area of Kiseynew metamorphism is accompanied by pyrrhotite. There are two types of mineral deposits in the area: (1) barren sulphide deposits consisting of pyrrhotite with a little pyrite and insignificant amounts of chalcopyrite and sphalerite; (2) copper-zinc orebodies consisting largely of pyrrhotite, but with appreciable amounts of chalcopyrite and sphalerite and a little pyrite. These two types have close structural and genetic relationships, and as pyrrhotite and pyrite are the earlier formed minerals, it is likely that both types were identical at one stage in their evolution; whereas at a later stage some of the pyrrhotite deposits were sufficiently mineralized with chalcopyrite and sphalerite to form orebodies.

#### Localization of Sulphide Deposits

Most of the sulphide deposits lie beneath narrow depressions or under lakes that occupy such depressions at intervals. As the depressions are partly filled with glacial material there is generally no surface indication of the presence of sulphides in the rocks beneath. Exploration must be conducted by systematic diamond drilling in places where geological information warrants the expense.

The greatest number of sulphide deposits has been found in the Sherridon group, although elsewhere in the area pyrrhotite and copper-zinc sulphides do occur in younger Kiseynew formations. Deposits are generally found in the impure quartzite members of the Sherridon group, especially near the contact of such rocks with dark hornblende gneisses. Shearing and fracturing has been localized near such contacts because the rocks on either side have offered different degrees of resistance under stress, and in this way fractures and channelways were provided that guided and confined the sulphide deposition derived from some igneous source beneath.

Most of the structures suitable for mineralization are localized in an area of intricate folding. The Sherridon group, being at or near the base of the Kiseynew gneisses, and thus in the vicinity of rocks having different physical characteristics, responded more violently to regional folding than did the overlying gneisses. Furthermore, the formations immediately above and below the Sherridon group as well as those comprising it form a greater variety of rock types than is found elsewhere in the area, and, as a result of differences in relative competency, have tended to localize folds and fractures in them during deforming movements. For these reasons the Sherridon provides the most favourable host rocks for mineral deposits.

There is a close association of pegmatite with many of the sulphide deposits. Pegmatite intrudes all members of the Kiseynew gneisses throughout the district, but is more prevalent along zones of weakness, such as contacts between hornblende gneiss and quartzite of the Sherridon group. The Bob Lake deposit is almost completely enclosed



in pegmatite, which is also present, although less obvious, in association with some of the barren pyrrhotite deposits. The time of intrusion of the pegmatite is earlier than the pyrrhotite mineralization, and it may have ascended the same channelways along which the sulphides later came. In some instances the pegmatitic intrusions accommodated themselves by bending the hanging-wall outward; in other places they burst through into the hanging-wall, tapering off into the country rock. Pegmatite is shattered where it is associated with sulphide deposits.

#### Sherritt Gordon Mine

The East and West orebodies of the Sherritt Gordon mine are together one of the longest sulphide deposits in the world, having a combined total length of almost 16,000 feet, of which 3,600 feet, between the two orebodies, carries no ore. As a cross-axial uplift passes through this barren interval it is probable that the two orebodies formed a single one before being reduced to the present erosion surface. The main orebodies have an average width of about 15 feet. The East orebody is approximately 250 feet deep, whereas the West orebody is 500 to 800 feet deep and rakes northward to a maximum depth below surface of about 1,500 feet. The West orebody tends to flatten out down the northward rake, due possibly to an internal fold in the Sherridon group. A relatively pure quartzite member forms the foot-wall (overturned hanging-wall) of the ore, whereas hornblende gneiss is generally on or near the hanging-wall. In the West orebody, particularly, there were subsidiary orebodies containing up to one-half million tons, which occurred as bulges or offsets into the hanging-wall. These offset deposits are almost entirely in pegmatite. The boundaries of the ore are relatively sharp and are somewhat better defined on the hanging- than on the foot-wall. The ore consists chiefly of pyrrhotite containing quartz nodules and more or less chalcopyrite and sphalerite, the East orebody being higher in zinc.

#### PROSPECTING IN THE KISSEYNEW GNEISSES

Certain conclusions may be drawn from observations at Sherritt Gordon mine and applied to the prospecting possibilities of the district generally. An obvious one is that the Sherridon group, on its eastward extension beyond the map-area, is the most promising formation for exploration. The Kisseynew gneisses cover a large area, extending northward beyond Granville Lake and westward into Saskatchewan, in any part of which structural conditions may be favourable for mineral deposition. On the basis of the study at Sherritt Gordon mine it would appear that formations in the more sharply folded anticlines will provide structures most favourable for mineralization. The reason for this is that the folding of the Kisseynew gneisses is characteristically open, and the folds plunge at fairly low angles. Thus large areas present a monotonous succession of the crystalline gneisses; but, wherever sharp anticlinal folds occur, older Kisseynew formations are exposed on the present erosion surface, and in some instances pre-Kisseynew, Archaean rocks may be exposed in the cores of the anticlines. As hornblende gneisses are more abundant in the lower part of the Kisseynew, and as the underlying Archaean rocks are derived largely from greenstones, the presence of favourable anticlinal structures may, in general, be detected from the relative abundance of dark rocks, especially the thicker bands. As geological information about most of the area of Kisseynew gneisses is fragmentary, prospecting might well commence with a search for favourable structures.

#### ACKNOWLEDGMENTS

The writer is indebted to Mr. Eldon L. Brown, General Manager, and Mr. W. J. Farley, Chief Geologist, of Sherritt Gordon Mines, Limited, for co-operation during the field work. The description of the Sherritt Gordon ore deposits is based largely on an examination of the geological plans of the mine and mine models prepared by Mr. Farley. Mr. Farley's geological work at the mine resulted in the discovery of the Bob Lake deposit, which lies entirely beneath the lake. The geology of the southern part of Sherridon map-area was mapped by J. M. Harrison of the Geological Survey.