O Orga (bog bog bog bog bog bog bog bog bog bog	anic g)  anic g)  anic g)  rmokarst  uvial odplain  luvial rrace  ciolacustrine in	Dominantly moderately decomposed fen peat derived from sedge, tamarack, and mosses  Dominantly moderately decomposed forest and/or undecomposed sphagnum peat derived from black spruce, cladonia, feather - mosses, ericaceous and/or sphagnum vegetation  Dominantly moderately decomposed forest and/or undecomposed sphagnum peat derived from black spruce, cladonia, feather - mosses, ericaceous and/or sphagnum vegetation  Gravel, sand, and silt; textures vary with the dominant material indicated first  Gravel, sand, and silt; textures vary with the dominant material indicated first  Colluvium over bedrock (see note 6 for lithology is shown asy a fraction- adventure for surficial deposits  Colluvium over bedrock (see note 6 for lithology indicated first  Sand, mainly fine to mdeium grained  Sand, mainly fine to mdeium grained	ESTIMATED THICKNESS  2-3 m  1.5-7 m  1-8 m  1-30 m	Flat to very gently sloping, some with reticulate network of low (<1 m) ridges (patterned fen)  Flat to gently sloping areas with scattered mounds (average relief 1 m, rarely to 6 m); numerous steepsided depressions and trenches  Flat to gently sloping areas with mounds (average relief 1 m, rarely to 6 m); numerous exposed peat scarps, depressions, and trenches  Floodplain and low bordering terraces; floodplains within mountains commonly scarred by braided channels; floodplains within plains region commonly with meander scars  Terraces with relief intermediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping (10-80)fans and coalescent fans  Gently to steeply sloping irregular surfaces; Cx1 <50, Cx2 50-200	DRAINAGE PATTERN  No organized drainage; water at surface throughout summer months  Depressions interconnected by seepage channels; drainage continually being modified  In braided areas intermittent drainage through channels; in meander scar areas no integrated drainage system, impeded by meander scroll ridges  Surface drainage without integrated drainage system  One or more shifting streams commonly present; downslope seepage in poorly defined runs	GROUND ICE (NOTE 2)  Unfrozen to at least 3 m  Frozen at 0.3-0.5 m. Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions commonly thawed to at least 1 m. Ice-wedge polygons present in zone 3  Frozen at 0.3-0.5 m. Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions, and scarps commonly thawed to at least 1 m  Ground ice in gravel, sand, and silt of low terraces in zone 2; Not observed elsewhere. Permafrost and segregated ice also present in areas where bog is more than 1.5 m thick  Frozen groundwater in gravel, sand, and silt of terraces, up to 30 m thick, in zone 2. Not observed elsewhere  None observed  Silty clayey colluvium contains disseminated ice crystals to seams 1 m thick in zones 2 and 6N	UNIFIED SOIL CLASSIFICATION (NOTE 5)  Pt  Pt  SM-GM to ML  Variable	PREDOMINANT SOIL*  Typic Mesisol (P-VP)  Mesic Organic Cryosol (W-I) Fibric Organic Cryosol (W-I)  Typic Fibrisol (P) Typic Mesisol (P)  Mesic Organic Cryosol (W-I)  Fibric Organic Cryosol (W-I)  Typic Fibrisol (P)  Typic Fibrisol (P)  Typic Mesisol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Brunisolic Turbic Cryosol (P)  Eluviated Eutric Brunisol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Gleyed Regosol (I) Gleyed Regosol (I) Gleyed Regosol (I) Gleyed Regosol (I) Gleyed Regosol (I) Gleyed Regosol (I) Gleyed Regosol (I) Gleyed Eutric Brunisol (W)	6S 10 10 5 5	6N 5 10 10 7 7 7 3 3 4 - 3 3 6 6 1 1 3 3 - 6 - 2 - 2	10 1 7 1 3 3 4 3 5 6 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	3 2 0 - 0 10 - 10 - 10 - 10	STABLE  Sedge-Bi or Sedge-Bi-tL in all zones  bS-Fm-Er in zones 6S and 6N; bS-lichen-Er in zones 4 and 5; lichen-bS-Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  bS-Fm-Er in zones 6S and 6N; bS-lichen-Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  wS-wB, bPo-wS-Eq, wS-bPo in zones 4, 5, 6S, and 6N  wS-wB, wS-bPo in zones 4, 5, and 6S  wS-wB, wS-bPo, tA in zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, wi, tA-wS in zones 4, 5, 6N, and 6S; wS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, wS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-Lichen-Er in zone 2	AFTER FIRE  Same, probably would not burn  Sphagnum, Er  Er, Fm, bS  Sphagnum  Wi, Al, bPo  WB, tA, bPo  WB, tA, bPo	Poor drainage, plus high compressibility and low strength of the material make it unsuitable for any type of construction  Alternation of permanently frozen peat mounds and thawed depressions and water bodies presents serious problems in construction of roads, pipelines, etc.; material highly compressible when thawed  Alternation of permanently frozen peat mounds, some actively degrading, and thawed depressions and water bodies presents serious problems in construction of roads, pipelines, etc.; material highly compressible when thawed  Subject to periodic flooding; floodplains within the mountains are potential sources of aggregate; extraction of aggregate may cause deleterious changes in stream course and downstream changes in stream regimen  Good construction sites and aggregate source where material is coarse
O Orga (bog bog ther start of the start of t	anic g)  anic g)  rmokarst  uvial odplain  luvial plex  ciolacustrine in	moderately decomposed fen peat derived from sedge, tamarack, and mosses  Dominantly moderately decomposed forest and/or undecomposed sphagnum peat derived from black spruce, cladonia, feather mosses, ericaceous and/or sphagnum vegetation  Dominantly moderately decomposed forest and/or undecomposed sphagnum peat derived from black spruce, cladonia, feather mosses, ericaceous and/or sphagnum vegetation  Gravel, sand, and silt; textures vary with the dominant material indicated first  Gravel, sand, and silt; textures vary with the dominant material indicated first  Colluvium over bedock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Si, dominant lithology indicated first  Sand, mainly fine to model my fine to mode in the symbol for the	1.5-7 m  1.5-7 m  1-8 m  1-6 m  Variable	some with reticulate network of low (<1 m) ridges (patterned fen)  Flat to gently sloping areas with scattered mounds (average relief 1 m, rarely to 6 m); numerous steepsided depressions and trenches  Flat to gently sloping areas with mounds (average relief 1 m, rarely to 6 m); numerous exposed peat scarps, depressions, and trenches  Floodplain and low bordering terraces; floodplains within mountains commonly scarred by braided channels; floodplains within plains region commonly with meander scars  Terraces with relief intermediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping interrupted by shallow channels and low terraces  Gently to steeply sloping irrequiar surfaces; Cx1<50,	Depressions interconnected by seepage channels  Depressions interconnected by seepage channels; drainage continually being modified  In braided areas intermittent drainage through channels; in meander scar areas no integrated drainage system, impeded by meander scroll ridges  Surface drainage without integrated drainage system  One or more shifting streams commonly present; downslope seepage in	Frozen at 0.3-0.5 m. Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions commonly thawed to at least 1 m. Ice-wedge polygons present in zone 3  Frozen at 0.3-0.5 m. Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions, and scarps commonly thawed to at least 1 m  Ground ice in gravel, sand, and silt of low terraces in zone 2; Not observed elsewhere. Permafrost and segregated ice also present in areas where bog is more than 1.5 m thick  Frozen groundwater in gravel, sand, and silt of terraces, up to 30 m thick, in zone 2. Not observed elsewhere	Pt Pt SM-GM SM-GM to ML	Mesic Organic Cryosol (W-I) Fibric Organic Cryosol (W-I) Fibric Organic Cryosol (W-I)  Typic Fibrisol (P) Typic Mesisol (P)  Mesic Organic Cryosol (W-I) Fibric Organic Cryosol (W-I)  Typic Fibrisol (P) Typic Mesisol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Gleyed Eutric Brunisol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Gleyed Regosol (I)  Gleyed Regosol (I)	10 5 5 5	10 10 7 7 7 3 3 3 - 4 - 3 3 6 6 6 3 3 3 - 6 - 2 - 2	10 1 3 3 4 3 6 - 1 1 - 3 3 6 6 1	0 10	bS-Fm-Er in zones 6S and 6N; bS- lichen-Er in zones 4 and 5; lichen-bS- Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  bS-Fm-Er in zones 6S and 6N; bS- lichen-Er in zones 6S and 6N; bS- lichen-Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  wS-wB, bPo-wS- Eq, wS-bPo in zones 4, 5, and 6S  wS-wB, wS-bPo, tA-wB, bPo-Al in zones 4, 5, and 6S  wS-wB, wS-bPo, tA-wB, bPo, tA in zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2	Sphagnum, Er  Sphagnum, Er  Er, Fm, bS  Sphagnum  Wi, Al, bPo  WB, tA, bPo  wB, Wi, tA, bPo	high compressibility and low strength of the material make it unsuitable for any type of construction  Alternation of permanently frozen peat mounds and thawed depressions and water bodies presents serious problems in construction of roads, pipelines, etc.; material highly compressible when thawed  Alternation of perman- ently frozen peat mounds, some actively degrading, and thawed depressions and water bodies presents serious problems in construc- tion of roads, pipelines, etc.; material highly compressible when thawed  Subject to periodic flooding; floodplains within the mountains are potential sources of aggregate; extraction of aggregate may cause deleterious changes in stream course and downstream changes in stream regimen  Good construction sites and aggregate source where material is
Ok Orga (bog ther)  As,siAp Allu floo  As,siAt Allu tery  Coll comp  i,sLp Glac plain  i,sLp plain	anic g)  anic g)  rmokarst  uvial podplain  luvial rrace  ciolacustrine in	Dominantly moderately decomposed forest and/or undecomposed sphagnum peat derived from black spruce, cladonia, feather - mosses, ericaceous and/or sphagnum vegetation  Dominantly moderately decomposed forest and/or undecomposed sphagnum peat derived from black spruce, cladonia, feather - mosses, ericaceous and/or sphagnum vegetation  Gravel, sand, and silt; textures vary with the dominant material indicated first  Gravel, sand, and silt; textures vary with the dominant material indicated first  Colluvium over bedocok (see note 6 for lithology) is shown as a fraction- ated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	1.5-7 m  1-8 m  1-30 m  Variable	with scattered mounds (average relief 1 m, rarely to 6 m); numerous steep- sided depressions and trenches  Flat to gently sloping areas with mounds (average relief 1 m, rarely to 6 m); numerous exposed peat scarps, depressions, and trenches  Floodplain and low bordering terraces; floodplains within mountains commonly scarred by braided channels; floodplains within plains region commonly with meander scars  Terraces with relief inter- mediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping irregular surfaces; Cxl 50,	Depressions interconnected by seepage channels; drainage continually being modified  In braided areas intermittent drainage through channels; in meander scar areas no integrated drainage system, impeded by meander scroll ridges  Surface drainage without integrated drainage system  One or more shifting streams commonly present; downslope seepage in	Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions commonly thawed to at least 1 m. Ice-wedge polygons present in zone 3  Frozen at 0.3-0.5 m. Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions, and scarps commonly thawed to at least 1 m  Ground ice in gravel, sand, and silt of low terraces in zone 2; Not observed elsewhere. Permafrost and segregated ice also present in areas where bog is more than 1.5 m thick  Frozen groundwater in gravel, sand, and silt of terraces, up to 30 m thick, in zone 2. Not observed elsewhere	Pt  SM-GM  SM-GM to ML	Typic Fibrisol (P) Typic Mesisol (P)  Mesic Organic Cryosol (W-I) Fibric Organic Cryosol (W-I) Fibric Organic Cryosol (W-I)  Typic Fibrisol (P) Typic Mesisol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Brunisolic Turbic Cryosol (P)  Gleyed Eutric Brunisol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Gleyed Regosol (W)  Gleyed Regosol (I)	5 - - 4 3 3 - - - - -	3 3 7 - 4 - 3 3 6 6 6 3 3 3 - 6 - 2 - 2	3 7 3 4 3 - 3 6 - 1 1 - 3 6 6	- 10	6S and 6N; bS- lichen-Er in zones 4 and 5; lichen-bS- Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  bS-Fm-Er in zones 6S and 6N; bS- lichen-Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  wS-wB, bPo-wS- Eq, wS-bPo in zones 4, 5, 6S, and 6S  wi, wi-Al, bPo-Al in zones 4, 5, and 6S  wS-wB, wS-bPo, tA-wB, bPo, tA in zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, wi, tA-wS in zones 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-Lichen-	Sphagnum, Er  Er, Fm, bS  Sphagnum  Wi, Al, bPo  WB, tA, bPo  WB, Wi, tA, bPo	Alternation of permanently frozen peat mounds and thawed depressions and water bodies presents serious problems in construction of roads, pipelines, etc.; material highly compressible when thawed  Alternation of permanently frozen peat mounds, some actively degrading, and thawed depressions and water bodies presents serious problems in construction of roads, pipelines, etc.; material highly compressible when thawed  Subject to periodic flooding; floodplains within the mountains are potential sources of aggregate; extraction of aggregate may cause deleterious changes in stream course and downstream changes in stream regimen  Good construction sites and aggregate source where material is
s,siAp Allu floo  s,siAt Allu terr  f Allu fan  Coll comp	anic g) rmokarst  uvial odplain  luvial rrace  ciolacustrine in	mosses, ericaceous and/or sphagnum vegetation  Dominantly moderately decomposed forest and/or undecomposed sphagnum peat derived from black spruce, cladonia, feather mosses, ericaceous and/or sphagnum vegetation  Gravel, sand, and silt; textures vary with the dominant material indicated first  Gravel, sand, and silt; textures vary with the dominant material indicated first  Colluvium derived from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	1-8 m  1-30 m  1-6 m  Variable	with mounds (average relief 1 m, rarely to 6 m); numerous exposed peat scarps, depressions, and trenches  Floodplain and low bordering terraces; floodplains within mountains commonly scarred by braided channels; floodplains within plains region commonly with meander scars  Terraces with relief intermediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping (10-80) fans and coalescent fans  Gently to steeply sloping irregular surfaces; Cxl <50,	nected by seepage channels; drainage continually being modified  In braided areas intermittent drainage through channels; in meander scar areas no integrated drainage system, impeded by meander scroll ridges  Surface drainage without integrated drainage system  One or more shifting streams commonly present; downslope seepage in	thawed to at least 1 m. Ice-wedge polygons present in zone 3  Frozen at 0.3-0.5 m. Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions, and scarps commonly thawed to at least 1 m  Ground ice in gravel, sand, and silt of low terraces in zone 2; Not observed elsewhere. Permafrost and segregated ice also present in areas where bog is more than 1.5 m thick  Frozen groundwater in gravel, sand, and silt of terraces, up to 30 m thick, in zone 2. Not observed elsewhere	SM-GM SM-GM to ML  Variable	Typic Mesisol (P)  Mesic Organic Cryosol (W-I) Fibric Organic Cryosol (W-I) Fibric Organic Cryosol (W-I)  Typic Fibrisol (P) Typic Mesisol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Brunisolic Turbic Cryosol (W)  Gleyed Eutric Brunisol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Orthic Regosol (W)  Gleyed Regosol (I)	4 3 3 6 - 1	7 - 4 - 3 3 6 6 6 3 3 - 6 - 2 - 2	7 3 4 3 6 - 1	- 10 	sphagnum-bS in zones 4, 5, 6S, and 6N  bS-Fm-Er in zones 6S and 6N; bS-lichen-Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  wS-wB, bPo-wS-Eq, wS-bPo in zones 4, 5 and 6S  wis-wB, wS-bPo, tan zones 4, 5, and 6S  wS-wB, wS-bPo, tan zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tan zones 4, 5, 6N, and 6S; wS-bPo, ws-bPo in zone 2  bPo-ws, wS-bPo, wi, tan zone 2  bPo-ws, wS-bPo, wi, tan zone 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-Lichen-	Er, Fm, bS  Sphagnum  Wi, Al, bPo  WB, tA, bPo  WB, Wi, tA, bPo	highly compressible when thawed  Alternation of permanently frozen peat mounds, some actively degrading, and thawed depressions and water bodies presents serious problems in construction of roads, pipelines, etc.; material highly compressible when thawed  Subject to periodic flooding; floodplains within the mountains are potential sources of aggregate; extraction of aggregate may cause deleterious changes in stream course and downstream changes in stream regimen  Good construction sites and aggregate source where material is
s,siAp Allufloo  Allufan  Coll comp  SLp Glac plain	uvial podplain  uvial rrace  uvial plex	moderately decomposed sand/or undecomposed sphagnum peat derived from black spruce, cladonia, feather mosses, ericaceous and/or sphagnum vegetation  Gravel, sand, and silt; textures vary with the dominant material indicated first  Gravel, sand, and silt; textures vary with the dominant material indicated first  Colluvium derived from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	1-8 m  1-30 m  1-6 m  Variable	with mounds (average relief 1 m, rarely to 6 m); numerous exposed peat scarps, depressions, and trenches  Floodplain and low bordering terraces; floodplains within mountains commonly scarred by braided channels; floodplains within plains region commonly with meander scars  Terraces with relief intermediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping (10-80) fans and coalescent fans  Gently to steeply sloping irregular surfaces; Cxl <50,	nected by seepage channels; drainage continually being modified  In braided areas intermittent drainage through channels; in meander scar areas no integrated drainage system, impeded by meander scroll ridges  Surface drainage without integrated drainage system  One or more shifting streams commonly present; downslope seepage in	Segregated ice content commonly 60-80% in peat; typically 10-100 cm thick. Segregated ice in mineral soil below. Peat in wet depressions, and scarps commonly thawed to at least 1 m  Ground ice in gravel, sand, and silt of low terraces in zone 2; Not observed elsewhere. Permafrost and segregated ice also present in areas where bog is more than 1.5 m thick  Frozen groundwater in gravel, sand, and silt of terraces, up to 30 m thick, in zone 2. Not observed elsewhere  None observed elsewhere	SM-GM SM-GM to ML  Variable	Typic Fibrisol (P) Typic Mesisol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Brunisolic Turbic Cryosol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleyed Eutric Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Orthic Regosol (W)  Gleyed Regosol (I)	4 3 3 6 - 1	3 - 4 - 3 3 6 6 6 3 3 3 - 6 - 2 - 2	4 3 - 3 6 - 1		6S and 6N; bS-lichen-Er in zones 2 and 3  Sphagnum or sphagnum-bS in zones 4, 5, 6S, and 6N  wS-wB, bPo-wS-Eq, wS-bPo in zones 4, 5 and 6S  Wi, Wi-Al, bPo-Al in zones 4, 5, and 6S  wS-wB, wS-bPo, tA in zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, wi, tA-wS in zones 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-Lichen-	Sphagnum  Wi, Al, bPo  Wi, Al, bPo  wB, tA, bPo	ently frozen peat mounds, some actively degrading, and thawed depressions and water bodies presents serious problems in construc- tion of roads, pipelines, etc.; material highly compressible when thawed  Subject to periodic flooding; floodplains within the mountains are potential sources of aggregate; extraction of aggregate may cause deleterious changes in stream course and downstream changes in stream regimen  Good construction sites and aggregate source where material is
sLp Glac sLpv Glain	uvial podplain  uvial rrace  uvial luvial plex	Gravel, sand, and silt; textures vary with the dominant material indicated first  Gravel, sand, and silt; textures vary with the dominant material indicated first  Mostly gravel, some sand  Colluvium derived from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	1-30 m  1-6 m  Variable	terraces; floodplains within mountains commonly scarred by braided channels; floodplains within plains region commonly within plains region commonly with meander scars  Terraces with relief intermediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping (10-80)fans and coalescent fans  Gently to steeply sloping irregular surfaces; Cxl <50,	mittent drainage through channels; in meander scar areas no integrated drainage system, impeded by meander scroll ridges  Surface drainage without integrated drainage system  One or more shifting streams commonly present; downslope seepage in	sand, and silt of low terraces in zone 2; Not observed elsewhere. Permafrost and segregated ice also present in areas where bog is more than 1.5 m thick  Frozen groundwater in gravel, sand, and silt of terraces, up to 30 m thick, in zone 2. Not observed elsewhere  None observed  Silty clayey colluvium contains disseminated ice crystals to seams 1 m thick in zones 2	SM-GM to ML Variable	Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Brunisolic Turbic Cryosol (W)  Gleyed Eutric Brunisol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Gleyed Regosol (W)  Gleyed Regosol (I)	3 3 6 - 1 3	- 3 3 6 6 6 1 1 1 3 3 - 6 - 2 - 2	3 6 - 1 - 3 6 6	6 1 - 3	wS-wB, bPo-wS-Eq, wS-bPo in zones 4, 5 and 6S  Wi, Wi-Al, bPo-Al in zones 4, 5, and 6S  wS-wB, wS-bPo, tA in zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, wi, tA-wS in zones 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-Lichen-	WB, tA, bPo  WB, Wi, tA, bPo	Subject to periodic flooding; floodplains within the mountains are potential sources of aggregate; extraction of aggregate may cause deleterious changes in stream course and downstream changes in stream regimen  Good construction sites and aggregate source where material is
Allufan  Coll comp  SLp Glac sLpv plait	uvial luvial plex ciolacustrine in	Mostly gravel, some sand  Colluvium derived from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	3-25 m  1-6 m  Variable	Terraces with relief intermediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping (10-80)fans and coalescent fans  Gently to steeply sloping irregular surfaces; Cxl <50,	Surface drainage with- out integrated drainage system  One or more shifting streams commonly present; downslope seepage in	where bog is more than 1.5 m thick  Frozen groundwater in gravel, sand, and silt of terraces, up to 30 m thick, in zone 2. Not observed elsewhere  None observed  Silty clayey colluvium contains disseminated ice crystals to seams 1 m thick in zones 2	to ML Variable	Gleysolic Turbic Cryosol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Brunisolic Turbic Cryosol (W)  Gleyed Eutric Brunisol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Orthic Eutric Brunisol (W)  Gleyed Regosol (I)	3	6 6 6 1 1 1 3 3 - 6 - 2 - 2	6 - 1 - 3 6	6 - 1 - 3	in zones 4, 5, and 6S  wS-wB, wS-bPo, tA-wB, bPo, tA in zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, wi, tA-wS in zones 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bs-Lichen-	wB, tA, bPo  wB, Wi, tA, bPo	deleterious changes in stream course and downstream changes in stream regimen  Good construction sites and aggregate source where material is
Allu fan  Coll comp  Aeol depo	uvial luvial plex ciolacustrine in	Mostly gravel, some sand  Colluvium derived from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	3-25 m  1-6 m  Variable	mediate between terraces associated with Ap and Gp or Gt; level to slightly sloping surfaces; some interrupted by shallow channels and low terraces  Gently to moderately sloping (10-80)fans and coalescent fans  Gently to steeply sloping irregular surfaces; Cxl 50,	One or more shifting streams commonly present; downslope seepage in	Silty clayey colluvium contains disseminated ice crystals to seams 1 m thick in zone 2	to ML Variable	Orthic Eutric Brunisol (W)  Brunisolic Turbic Cryosol (W)  Gleyed Eutric Brunisol (I)  Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Gleyed Regosol (W)  Gleyed Regosol (I)	3	1 1 1 3 3 3 - 6 - 2 - 2	1 - 3	- 6 1 - 3 - 6	tA-wB, bPo, tA in zones 4, 5, 6N, and 6S; wS-wB, bS-bPo, tA-bS in zone 2  bPo-wS, wS-bPo, Wi, tA-wS in zones 4, 5, 6N, and 6S; bS-bPo, wS-bPo in zone 2  bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bS-Lichen-	wB, Wi, tA, bPo	and aggregate source where material is
Aeol depo	luvial plex lian osits	Colluvium derived from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	1-6 m Variable	Gently to steeply sloping irregular surfaces; Cxl 50,	streams commonly present; downslope seepage in	Silty clayey colluvium contains disseminated ice crystals to seams 1 m thick in zones 2	GC-GP	Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Orthic Eutric Brunisol (W)  Gleyed Regosol (I)	-	3 3 - 6 - 2 - 2		- 1 3 - 6	and 6S; bS-bPo, wS-bPo in zone 2 bS-Fm, bS-Fm-Er, bS-sphagnum in zones 4, 5, 6N, and 6S; bs-Lichen-	Wi	
Aeol depo	luvial plex lian osits	Colluvium derived from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	1-6 m Variable	Gently to steeply sloping irregular surfaces; Cxl 50,	streams commonly present; downslope seepage in	Silty clayey colluvium contains disseminated ice crystals to seams 1 m thick in zones 2	GC-GP	Cumulic Regosol (W)  Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)  Orthic Regosol (W)  Orthic Eutric Brunisol (W)  Gleyed Regosol (I)		- 6 - 2 - 2		- 3	and 6S; bs-Lichen-		/ / / / / / / / / / / / / / / / / / / /
Aeol depo	rlian osits	from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	Variable	irregular surfaces; Cx1 <50,		contains disseminated ice crystals to seams 1 m thick in zones 2	to	Rego Gleysol (P) Gleysolic Turbic Cryosol (P) Orthic Regosol (W) Orthic Eutric Brunisol (W) Gleyed Regosol (I)	-	- 2	2	1	Mainly wB, tA, Wi, and Al in zones 2,	Wi, Al, wB	Generally offer well drained building sites,
Aeol depo	rlian osits	from entire range of surficial deposits  Colluvium over bedrock (see note 6 for lithology) is shown as a fractionated symbol, e.g., Cx Si, Sh; dominant lithology indicated first  Sand, mainly fine to mdeium grained	Variable	irregular surfaces; Cx1 <50,		contains disseminated ice crystals to seams 1 m thick in zones 2	to	Orthic Eutric Brunisol (W) Gleyed Regosol (I)			2 2	- 2		tA, wB at low	but sudden channel shifting may cause damage; good source of aggregate
Lp Glac Lpv plain	ciolacustrine in	Sand, mainly fine to mdeium grained	1-20 m	Constitution of the second			14/200			2/16		4/1	tA, WS, WB at low elevations and stunted Alpine fir or wS, Er, lichen at high elevations in zone 5; bS, wB at low elevations and Er, lichen at high elevations; cryoturbated surfaces are unvegetated in zones 2	elevations	Irregularity of topography and slope instability present major problems for any type of construction texture prefix included in map unit if known
Lp Glac Lpv plain	ciolacustrine in	fine to mdeium grained	1-20 m					Rego Gleysol (P) Gleysolic Turbic Cryosol (P)		- 3		3 3	and 3  Sphagnum, bS- sphagnum-Er in zones 2, 3, and 5	Sphagnum, Er	
Lh Hummn glac	in		The state of the s	sEr, dune ridges, usually parallel to subparallel; sEh, irregularly shaped dunes, no apparent pattern	Mainly subsurface seepage	None observed, probably no ice present	SP-SW	Eluviated Eutric Brunisol (W)  Gleyed Eluviated Eutric Brunisol (	5				P, tA-wB in zones 5 and 6S tA-bS-Wi in zones 5 and 6S	P, tA	Subject to wind erosion when vegetation mat is removed
Lh Hummn glac	in			VE SUBSE			MA	Rego Gleysol (P) Gleysolic Turbic Cryosol (P)	3	- 3	-		bS-Fm, bS- sphagnum-Er in zones 5 and 6S	Wi	
glac		Mainly silt and fine sand; locally includes gravel (g) and clay (c); dominant material indicated first	si,sLp 1.5-50 m si,sLpv 0.5-1.5 m	Flat to gently sloping	Surface seepage through organic-filled depressions and downslope seepage in shallow subparallel runs	Commonly 10-50% segregated ice as thin (1 mm - 2 cm) seams parallel with bedding, segregated ice in a reticulate network up to 60% by volume, and thick	CL-ML to SM	Brunisolic Grey Luvisol (W) Orthic Grey Luvisol (W)	3	2 -			wS, wS-tA in zones 5 and 6S; - wS-tA, bS-wB- tA in zone 6N	tA	Failure common along scarps; generally unsuitable for location of structures because of poor bearing capacity and drainage character-
glac	11/10000	The second secon				tabular bodies in zones 2 and 6N; discontinuous zones of segregated ice as seams, reticulated net- works, and tabular bodies (up to 3 m thick) in northern part of zone		Gleyed Grey Luvisol (I)  Luvisolic Turbic Cryosol (I)  Rego Gleysol (P)	- 5	2 -	-		bS-wB-tA, bS- Wi-Al in zones 5, 6S, and 6N bS-Fm-Er, bS- tL-wB in zones	Wi, Al, wB	istics
	mocky ciolacustrine	Mainly silt and fine sand	2-5 m	Low hummocks up to 2 m; individual hummocks up to	Deranged	6S; no ice recorded in southern part of zone 6S except below p0  None observed	CL-ML	Gleysolic Turbic Cryosol (P)  Brunisolic Grey Luvisol (W)	4	6 5	N		wS, wS-tA in zones 5 and 6S;	tA	Failure common along scarps; generally
depo	osits	THE SUITO		5 m relief; slopes to 200				Orthic Grey Luvisol (W)  Gleyed Grey Luvisol (I)  Luvisolic Turbic Cryosol (I)  Rego Gleysol (P)	- 2 - 5	2 - 2 2	-		wS-tA, bS-wB- tA in zone 6N bS-wB-tA, bS- Wi-Al in zones 5, 6S, and 6N bS-Fm-Er,bS-tL-	tA, wB Wi, Al, wB	unsuitable for location of structures because of poor bearing capacity and drainage characteristics
	ches	Mainly gravel with minor sand; locally include silt(si); dominant material indicated	g,sLpbx 1.5-2 m g,sLpbxv 0.5-1.5 m g,sLpbv 0.5-1.5 m	g,sLpbx and g,sLpbxv, parallel to subparallel beach ridges arranged in belts; up to 60 slopes; g,sLpbv, beach material without distinct ridges forming	Drainage mainly subsurface	None observed	SP-GP	Gleysolic Turbic Cryosol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Gleyed Eutric Brunisol (I)	5		-		P-wB-Wi, bS-wB-Wi, tA-bS-Wi	P, tA, wB	Good construction sites and aggregate source where material is coarse; beaches at 800 -900 foot (240-275 m)
plain	ciofluvial in	Gravel, sand and silt; textures vary	g,s,siGp 1-30+ m	belts up to 6+ km wide  Flat to gently sloping	Drainage mainly subsurface	Frozen groundwater in gravel, sand, and silt in	GP-GW	Rego Gleysol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)	3		-	-   -	bS-Fm-Er p-wB-tA, P in zones 6S and 6N	Wi P, tA	elevations offer the best potential  Good construction sites and aggregate source
siGt Glaci terra	ciofluvial	with the dominant material indicated first	g,s,siGpv 0.5-1.5 m g,s,siGt 1-30+ m			zone 2. Not observed elsewhere. Where overlying bog is more than 1.5 m thick perma- frost may be present below	SP-SW	Gleyed Eutric Brunisol (I)  Rego Gleysol (P)	3		-		bS-wB-Wi,P-wB- Wi,tA-bS-Wi in zones 6S and 6N bS-Fm-Er in zones 6S and 6N	P, tA, Wi, wB	— where material is coarse
glac depo: Ridge glac	ciofluvial osits	Mainly gravel with sand	1-10 m	Hummocks with local relief up to 10 m Long, sinuous esker ridges, up to 30 m high	Drainage mainly subsurface	No segregated ice in well drained sites, but segregated ice may be present in association with silt layers beneath depressions in zones 2,	GP-GW to SP-SW	Gleysolic Turbic Cryosol (P)  Eluviated Eutric Brunisol (W) Orthic Eutric Brunisol (W)  Gleyed Eutric Brunisol (I)		3 -	5		tA-bS-wB, P, wS-tA-wB	P, tA, wB	Good construction sites and aggregate
incl	ludes ers and er complexes aine in	Moderately to strongly calcareous glacial till, typically clay,	tMp 1.5-50 m	Flat to uniformly sloping; tMp <sup>1</sup> , tMv <sup>1</sup> : slope 2°- 5° tMp <sup>2</sup> , tMv <sup>2</sup> : slope 5°-15°; map symbol may be suffixed by one or more r,	Downslope seepage in subparallel runs Downslope seepage in shallow channels	Locally 5-40% segregated ice as thin (1 mm-2 cm) horizontal layers or in a reticulated net-	CL	Gleysolic Turbic Cryosol (P)  Brunisolic Grey Luvisol (W)  Orthic Grey Luvisol (W)	4	- 3 3	3 - 3		bS-Fm-Er, bS-lichen-Er wS-tA-wB, tA- P-wB in zones 4, 5, 6S, and 6N; wS-wB, bS-wB in	Wi tA, P, wB wB, tA	Failure common along scarps; poor source of aggregate; up to 10% unmapped p0 and/or f0;
	45	silt, and minor sand with 5% pebbles and boulders; pre- fixes g, s, si, or c, indicate lenses of gravel, sand, silt, or	55	m, s, d, or h (see below) indicating the mapped area in part consists of one or more of these landform units		work in zones 2, 3, and 6N; presence controlled by exposure, elevation, drainage, and/or organic cover		Brunisolic Turbic Cryosol (W)  Gleyed Grey Luvisol (I)  Gleyed Brunisolic Grey Luvisol (I)	2	- 3	2	4 5	wS-tA-wB, wS- tA-Wi in zones 4, 5, 6S, and 6N;	tA, wB, Wi,	because of drainage by numerous subparallel runs, roads, or berms normal to slope direction, numerous culverts required to avoid impoundment of surface water; slopes susceptible to soil
		clay within the till. If till is thin (tMv), the nature of the underlying bedrock is shown by a fractionated symbol (Note 6 gives bedrock	ċΜv <1.5 m		Downslope seepage in subparallel runs		7	Brunisolic Turbic Cryosol (I)  Rego Gleysol (P)	4	2 -		1 1	bS-Wi-Al, bS-wB in zones 2 and 3	Wi, Al	creep and channelling
Flut	tings and	Tithology); dominant lith- ology indicated first  Moderately to strongly calcareous	2-30 m	Area consisting largely of parallel drumlins and/or	Trellis pattern or deranged drainage in	Segregated ice may be present in some drum-	CL	Gleysolic Turbic Cryosol (P)  Brunisolic Grey Luvisol (W)	4		5	5 4	- 6S; bS-Fm-lichen- Er in zones 4, 5, and 6N; bS-lichen- Er in zones 2 and 3 P-tA, tA-P-wB, wS-tA-wB in	tA, P, wB	Poor source of aggregate except where
	mlins tings	glacial till; pre- fixes g, s indicate lenses of gravel or sand within the till		flutings	drumlin areas to parallel seepage or streams in fluted areas	lins and flutings in zones 2, 3, and 6N, although none has been observed; intervening depressions in the same zones likely contain segregated ice; presence con-		Orthic Grey Luvisol (W)  Brunisolic Turbic Cryosol (W)  Gleyed Grey Luvisol (I)	2	- 2	2	4 4	zones 4, 5, and 6S; wS-wB, bS- wB-Wi in zones 2 and 3	Wi, wB, tA	gravel is present; some drumlins and most flutings are bedrock cored; crests of drumlins and flutings typically well drained; intervening depressions poorly drained
		Will Company				trolled by exposure, elevation, drainage, and/or organic cover		Gleyed Brunic Grey Luvisol (I)  Brunisolic Turbic Cryosol (I)			-	2 2	tA-wB in zones 4, 5, and 65; bS-Wi-A1, bS- wB in zones 2 and 3	AT	
							1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Rego Gleysol (P) Gleysolic Turbic Cryosol (P)	4	- 4	5	4 4	bS-Fm-Er in zone 6S; bS-Fm-lichen- Er in zones 4 and 5; bS-lichen-Er in zones 2 and 3	Wi, Al	
Mir Creve fill ridge mora	aine	Moderately to strongly cal- careous glacial till, gravel, and sand; textures vary with the dominant mate-	1-10 m	Individual, parallel to sub- parallel, straight to sinuous ridges within a moraine plain; 0.5-5 m relief; slopes 50-300	Integrated, weakly developed drainage controlled by ridge pattern	None observed, probably no ice present	CL	Brunisolic Grey Luvisol (W) Orthic Grey Luvisol (W) Brunisolic Turbic Cryosol (W)	6	- 2	6	10 11	P-tA, tA-P-wB, wS-tA-wB in zones 4, 5, and 6S; wS-wB in zones 2 and 3	P, tA, wB	Crests of prominent ridges offer restricted but good construction sites
		dominant mate- rial indicated first	7				CL to GP-SP	Rego Gleysol (P) Gleysolic Turbic Cryosol (P)	4			4 4	bS-Fm-Er in zone -6S; bS-Fm-lichen- Er in zones 4 and 5; bS-lichen-Er in	Wi, Al	Vote 1 State 1
sMh Hummo mora	aine	Moderately to strongly cal- careous glacial till, gravel, and sand; textures	1-20 m	Individual to coalescent hummocks; slopes to 20°	Deranged	Crests of prominent hummocks well drained and ice free; lower slopes and depressions commonly have 5-40%	CL	Brunisolic Grey Luvisol (W) Orthic Grey Luvisol (W) Brunisolic Turbic Cryosol (W)	6	- 6 -	6		P-tA, tA-P-wB, wS-tA-wB in zones 4, 5, and 6S; wS-wB, bS- wB in zones 2	tA, P, wB	Crests of prominent hummocks offer restricted but good construction sites
X3-1/15 A		vary with the dominant mate- rial indicated first				segregated ice as thin (1 mm-2 cm) horizontal layers or in a recticulated network in zones 2, 3, and 6N; presence controlled by exposure, elevation, drainage,	CL to GP-SP	Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)	4	4	4	.   -	bS-Fm-Er in zone 6S; bS-Fm- lichen-Er in zones 4 and 5; bS-lichen-	Wi, Al	
and	mocky rolling	Moderately to strongly cal- careous glacial till	5-30 m	Subdued hummocks and rolling terrain; slopes 50-300	Deranged	well drained sites ice free; lower slopes and depressions commonly have 5-40% segregated	CL	Brunisolic Grey Luvisol (W) Orthic Grey Luvisol (W)	6	- 6 -	6		P-tA, tA-P-wB, wS-tA-wB in zones 4, 5, and 6S; wS-wB, bS-	tA, P, wB	Summits of broad hum- mocks typically well drained and offer restricted but good
		19				ice as thin (1 mm-2 cm) horizontal layers or in a reticulate network in zones 2, 3, and 6N; presence controlled by exposure, elevation, drainage, and/or organic cover	N. Comments	Brunisolic Turbic Cryosol (W)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)	4	1/3	- 4	6 6	wB in zones 2 and 3 bS-Fm-Er in zone 6S; bS-Fm- lichen-Er in zones	Wi, Al	construction sites
	lanche;	Material derived mainly from glacio- lacustrine silts	Variable	Debris avalanches commonly occur as thin narrow tongues; earthflows and mudflows as		Fine grained material may contain segregated ice in zones 2, 3, 6N,	Variable	Cumulic Regosol (W)		- 4		4 4	4 and 5; bS-lichen- Er in zones 2 and 3 bS-wB-Wi-fireweed or unvegetated in	Firewood, Wi	Shale bedrock commonly fails as debris avalanches in mountainous regions;
iS, eart iSx, mudf flow & sl	thflow, flow, wslide,	lacustrine silts and clays, till. Material derived mainly from shale bedrock		earthflows and mudflows as bulbous masses; and slump deposits as blocks		ice in zones 2, 3, 6N, and northern part of 6S; presence controlled by exposure, elevation, drainage, and/or organic cover		Gleyed Cumulic Regosol (I)  Rego Gleysol (P)  Gleysolic Turbic Cryosol (P)	-	- 4		5 5	zones 2, 3, and 5 Wi, bS-Wi in zones 2, 3, and 5	Wi	in mountainous regions; large-scale failures are common along major river scarps in fine textured glacial deposits and in shale bedrock
W.		200	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	DESCRIPTION OF MA	AP UNITS FOR BEDE	ROCK AND MOUNTAI	IN TERRAIN						HELL		

MAP SYMBOL	NAME	TOPOGRAPHY	MILITA	PRINCIPAL BEDROCK UNITS**	SURFICIAL DEPOSITS	CODE	MICRORELIEF	SOILS AND VEGETATION ***	ENGINEERING COMMENTS
in -1i	Mountains developed in carbonate rockslimestone and dolomite with minor shale	Mountains up to 1500 m with sharp ridge crests and steep slopes. Ram Plateau area forms a level surface, dissected by deep canyons	L	includes Headless, Landry, and Nahanni formations in Mackenzie Mountains and Flett Formation, in Silent Hills and Liard Range includes Arnica, Manetoe, and Sombre formations in	Bare rock with discontinuous patches of limestone and dolomite rubble, bedrock colluvium with a matrix of fines and sand and/or reworked till on crests and steep slopes; variable thicknesses	GP-GC	Well developed stone polygons, stripes, and nets on flat to sloping ground. Numerous small solifluction lobes in McConnell Range of Franklin Mountains	Steep slopes: no soil development; lichen on bare surfaces. Base of slopes: Orthic Eutric Brunisol or Brunisolic Turbic Cryosol; above timberline Brunisolic Turbic Cryosol; lichen, ericaceous plants above timberline; white spruce, trembling aspen below	Steep slopes and high relief present serious difficulties to engineæring activities such as road, pipelime, and related construction; limestone and dolomite are highly resistant and could be used for construction material; coarse deposits of rubble make suitable
			canyons	L,D D,L L,Sh	Mackenzie Mountains includes Whittaker and Sunblood formations in Mackenzie Mountains includes Delorme Formation in Mackenzie Mountains and Camsell Formation in Camsell Range includes Funeral Formation in Mackenzie Mountains	of bedrock rubble and colluvium and/or reworked till at base of slopes; discontinuous veneer of till on plateau areas			timberline. Ram Plateau: Orthic Eutric Brunisol, Brunisolic Turbic Cryosol, or Orthic Regosol; lichen, ericaceous plants, some white spruce
,Sh ,Si,Sh M	Mountains developed on sandstone with shale, siltstone, or limestone combinations	Mountains up to 1600 m with rounded summits, long moderately steep slopes, and extensive debris mantles	S,Sh S,Si,Sh	includes Mount Clark Formation in Franklin Range and an unnamed Proterozoic sandstone in Moose Prairie area Mattson Formation includes Trout River and Redknife formations on Mackenzie Plain	Bare rock with discontinuous patches of sandstone, and limestone rubble, bedrock colluvium with a matrix of fines and sand and/or reworked till on crests and steep slopes; variable thicknesses of bedrock rubble and colluvium and/or reworked till at base of slopes	GP-GC	Stone polygons, stripes, and nets developed on flat to sloping ground	Steep slopes: no soil development; lichen on bare surfaces. Base of slopes: Orthic Eutric Brunisol, Brunisolic or Regosolic Turbic Cryosol, Orthic Regosol; above timberline Brunisolic Turbic Cryosol; lichen, ericaceous plants above timberline, white spruce, trembling aspen below timberline	Steep solopes and high relief present serious difficulties to engineering activities such as road, pipeline, and related construction; sandstone and limestone are stable and resistant and could be used for construction materials; shale is unstable and is subject to mass wasting; detachment slides and rotational slumping commonly occur when organic cover and/or vegetation are removed or altered; fine grained colluvium contains ice
Sh H	High hills developed on sandstone and shale with minor limestone member	Rounded summits and moderately steep slopes off the lower areas of the main mountain ranges	S,Sh S,Si,Sh	Mattson Formation includes Trout River and Redknife formations on Mackenzie Plain	Discontinuous patches of clayey, silty, sandy colluvium, sandstone and limestone rubble, reworked till, and/or bare rock on slopes and summit areas; variable thicknesses of bedrock rubble and colluvium and/or reworked till at base of slopes	GC-GP	Stone polygons on flat surfaces. Terracettes and poorly developed stripes on slopes	Orthic Eutric Brunisol, Brunisolic or Regosolic Turbic Cryosol; trembling aspen, white birch, white spruce	Shale iis highly unstable and is subject to mass wasting; detachment slides and rotational slumping commonly occur when organic cover and/or vegetattion are removed or altered; fin grained colluvium contains ice; sandstone and limestone rubble could be source of aggregate
h,Si,S h h,S h,S,Cg h,L,Si	High hills developed on shale with limestone and siltstone members	Well rounded summits with moderate slopes, forming much of the lower areas between Franklin Mountains and Front Ranges of Mackenzie Mountains to the west	Sh,Si,S Sh Sh,S Sh,S,Cg Sh,L,Si	Fort Simpson Formation includes Horn River and Klassen formations Fort St. John Formation and Cretaceous shale and siltstone of the Interior Plains Buckinghorse Formation Upper Devonian shale, limestone, and siltstone of the Interior Plains	Discontinuous patches of clayey, silty, sandy colluvium, sandstone, and limestone rubble, reworked till and/or bare rock on slopes of summit areas; variable thicknesses of bedrock rubble and colluvium, and/or reworked till at base of slopes	CL-ML	Stone polygons on flat surfaces. Terracettes and poorly developed stripes on slopes	Brunisolic or Regosolic Turbic Cryosol; trembling aspen, white birch, white spruce	Shale iis highly unstable and is subject omasss wasting; detachment slides and rotational slumping commonly occur when organic cover and/or vegetation are remmoved or altered; fine grained colluvium probably contains ice; sandstone and limestone rubble could be source of aggregate
,Si,Sh	Low hills developed on sandstone and shale with a small area of limestone	Moderate to gentle slopes and rounded summits containing little or no glacial drift indicate bedrock control of topography in formerly glaciated areas	S,Si,Sh	includes Trout River Formation	Veneer of clayey, silty, sandy colluvium; some with till and/or rubble	GC-GP to ML		Orthic Eutric Brunisol, Brunisolic or Regosolic Cryosol, Orthic Regosol; trembling aspen, white birch, white spruce	Shale iis highly unstable and is subject to mass wasting; detachment slides and rotational slumping commonly occur when organic cover and/or vegetation are removed or altered; fine grained colluvium may contaim ice in northern areas or at higher elevation

LEGEND

	ls shown in the legend may not aped on the face of the map and may		oundaries	
Rock outcrop			×	
Geological boundary (defined or a	pproximate, assumed or transition	na1)		
End or lateral moraine				
Crevasse fillings, moraine ridges			er 1/1	
Drumlinoid ridges, striae, flutin	gs (direction of ice movement kn	own, unknown)	F # S *	
Ice gouge			0	
Esker (direction of flow known, u	nknown)		« »«»«	
Beach ridge (depositional, erosio	mal)		-	
Meltwater channel (large, small).			HHHH+	
Landslide scar				
Escarpment			TITITITITITITITI	
Sinkhole				
Area of potential slumping			·······	
Patterned ground	1 33-92			
Rock glacier				
	NOTE 1			
	MAP UNIT DESIGNATION			
map unit, e.g. gAf. The upport of the letter (s) that for front of the genetic classif	used to designate each map unit of er case letter indicates the broad ollows indicates morphology; the ication describes texture. A num tMp' is used for ground moraine	ad genetic class; the lower case letter mber superscript is	ne (s) in	
Textural Modifier (placed in front of genetic category)	Genetic Classification		gic Modifier genetic category)	
r - rock and rubble	0 - organic	p - plain	d - drumlinoid	
t - till	A - alluvial	b - beach	c - channelled	
g - gravel	C - colluvial	r - ridged	h - hummocky	
s - sand	E - eolian	m - rolling	k - thermokarst	
si - silt	L - lacustrine	t - terraced	e - eroded	
c - clay	G - glaciofluvial	x - complex	f - fan	
p - bog	M - morainal	s - striated	v - veneer	
f - fen	S - slump	(i.e., flutings)	(<1.5m thick)	
	Number Supersoriet			

1- gentle slope (2°-5°); 2-steep slope (5°-15°)
- slope is up to 20° in Cx units


Complex Units

- a dash (-) means "with 16-49%" of the following unit (e.g., tMp-p0, p0 constitutes 16-49% of the area)
- a slash (/) signifies that the following unit comprises 5-15% of the total area within the boundaries of that complex unit
- a plus (+) indicates the presence of an unknown percentage of a second unit; less than 5% of a second unit is ignored
- fractionated units indicate a veneer of one unit over a thicker zone of another unit: (e.g., s.silpv a flat-surfaced veneer of sandy, silty lacustrine material over till)

- brackets () separate groups of units (e.g., (p0-f0)-tMp means p0 containing 16-49% f0, and 16-49% of the total area is tMp

Number Superscript

N.B. - one textural modifier signifies the dominant material
 - commas used between textural modifiers for distinct lithologies: e.g., g,s,siAp - floodplain consisting predominantly of gravel and sand with minor silt



SOILS
(by C. Tarnocai)

\*Drainage characteristics of soils

W = well

I = imperfect

P = poor

VP = very poor

Soils developed on si,sLh
In climatic zone 6S, 50% of soil cover is either Rego Gleysol or Cryic Rego Gleysol,
20% Gleyed Grey Luvisol, and 30% Brunisolic Grey Luvisol
In climatic zones 4, 3, and 2, there are either no exposures of si,sLh, or not enough mappable exposures to warrant generalisations about the soil type

NOTE 4

VEGETATION (by C. Tarnocai) Vegetation: species abbreviation bS - black spruce (Picea mariana) Sedge - Carex sp. wS - white spruce (Picea glauca) Cottongrass - Eriophorum Sp. wB - white birch (Betula neoalaskana) Lichen - Cladonia Sp., Cetraria Sp. Bi - dwarf birch (Betula glandulosa) Sphagnum - Sphagnum Sp. Er - Ericaceae (Ledum, Chamaedaphne, Kalmia, etc.) tL - tamarack (Larix laricina) Fm - feathermosses Wi - willow (Salix Sp.) P - pine (*Pinus banksiana* and *Pinus contorta* var. *latifolia*) Al - alder (Alnus sp.)

tA - trembling aspen (Populus tremuloides)

bPo - balsam poplar (Populus balsamifera)

Eq - horsetails (Equisefum sp.)

Major Group divisions Symbol					Typical Names	Classification criteria for coarse grained soils						
Coarse grained soils (more than half of material is larger than No.200 sieve size)	fraction size)	vith fines uble f fines)	GW		Well graded gravels, gravel-sand mixtures, little or no fines	O sieve size), GW, GP, SW, requiring	$c_u = D_{60}/D_{10} > 4$ $c_r = 1 < D^2 30/D_{10} > 4$	D <sub>60</sub> <3				
	Gravels (more than half of coarse firs larger than No.4 sieve s	Gravels with fine (appreciable amount of fines)	GP		Poorly graded gravels, gravel-sand mixtures, little or no fines	. 200 sieve 5% - GW, GF ases requir	Not meeting all gr requirements for					
		gravels le or no	GM	d	Silty gravels, gravel-sand-silt mixtures	of fines (fraction smaller than No. 200 classified as follows: Less than 5% - GC, SM, SC; 5 to 12% - Borderline cases	Atterberg limits below A line or 1 <sub>p</sub> <4	Above A line with				
		Clean g (little fines)	GC Clayey gravels, gravel-sand-	tion small follows: to 12% - Bo	Atterberg limits re	borderline cases requiring use o dual symbols						
	Sands han half of coarse fraction ler than No.4 sieve size)	ands or no	SW		Well graded sands, gravelly sands, little or no fines	fines (frac ssified as SM, SC; 5 t	$c_u = D_{60}/D_{10} > 6$ $c_r = 1 < D^2_{30}/D_{10} \times 6$	D <sub>60</sub> < 3				
		Clean sands (little or n fines)	SP		Poorly graded sands, gravelly sands, little or no fines	ntages of 1s are clas	Not meeting all gr requirements for					
		h fines ble fines)	SM	d	Silty sands, sand-silt mixtures	Depending on percentages coarse grained soils are SP; More than 12% - GM, 6 dual symbols	Atterberg limits below A line or 1 <sub>p</sub> <4	Limits plotting i hatched zone with 4 < 1p < 7				
	(more than is smaller	Sands with fappreciable	SC	2	Clayey sands, sand-clay mixtures	Dependin coarse g SP; More dual sym	Atterberg limits above A line with 1 <sub>p</sub> > 7	are borderline cases requiring use of dual symbols				
sieve size)	nd clays				ML		Inorganic silts and very fine sands, rock flour, silty or clayey fine sands, or clayey silts with slight plasticity		ATAI PRINT			
Fine grained soils half of material is smaller than No.200 s			CL		Inorganic clays of low to medium plasticity, gravelly clays, sandy clays, silty clays, lean clays	1000						
	Sil	Silts (liquid			Organic silts and organic silty clays of low plasticity							
	ys	>50)	МН		Inorganic silts, micaceous or diatomaceous fine sandy or silty soils, elastic silts							
	Silts and clays		СН		Inorganic clays of high plasticity, fat clays							
an			THOUGH T	77	REPORTED THE PROPERTY OF THE PARTY OF THE PA	20,000						

BEDROCK AND MOUNTAIN TERRAIN MAP SYMBOL DESIGNATION

TOPOGRAPHIC SYMBOL

BEDROCK

SLOPE ANGLE

MORPHOLOGIC MODIFIER

M Mountain: local relief greater than 450 m

High Hill: local relief 150-450 m

L Limestone

L Low Hill: local relief 30-150 m

S Sandstone

S Sandstone

S Siltstone

Mudstone

Sh Shale

Ch Chert

Cg Conglomerate

Texture - see NOTE 5

Organic clays of medium to high plasticity, organic silts

EXAMPLE

(Bedrock) (Texture - surficial)
D, L GC-GP

M

M

discontinuous residual or colluvial veneer of clayey gravels, gravel-sand-silt mixtures, and poorly graded gravels, with slopes generally 150-350, in places > 350, but locally 50-150 order of abundance)