


Magnetic contour intervals (total field)
 500 gammas
 100 gammas
 20 gammas
 10 gammas
 Magnetic depression contour
 Flight line
 Flight altitude: 1,000 feet above ground level

MAP 37G
AMOS
 ABITIBI COUNTY
 QUEBEC

Scale: One Inch to One Mile = $\frac{1}{63,360}$ Miles



Magnetic survey, August 1948, by Geophysics Division, Geological Survey of Canada; Department of Mines and Technical Surveys, in collaboration with the Royal Canadian Air Force.

No correction has been made for regional variation; this increases at the rate of 4.5 gammas per mile from east to west and 3.0 gammas per mile from south to north.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below, the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentine, which have a relatively high iron content; but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced by the geologist into, or across, areas of few or no outcrops. In many instances, however, no present interpretation of particular anomalies may be possible.