
GEOLOGICAL SURVEY OF CANADA



## Isomagnetic lines (total field) 500 gammas 100 gammas 20 gammas 10 gammas Hagnetic depression contour Flight line

Flight altitude: 1,000 feet above ground level

INDEX MAP

## DESKENATLATA LAKE SOUTH

DISTRICT OF MACKENZIE

NORTHWEST TERRITORIES

		Scale :	1:50 000 -	Échelle 1/50	000		
Kilometres	1	0	11	2	3	4	Kilomètre
Universal Transverse Mercator Projection © Crown copyrights reserved			Projection transverse universelle de Mercator © Droits de la Couronne réservés			r	
	thro		onal Air Pho		may be obtain ary, Topograph		

Magnetic Survey, August, 1950, by Geophysics Section, Geological Survey of Canada, Department of Mines and Technical Surveys.

No correction has been made for regional variation; this increases at the rate of 2.5 gammas per mile from west to east and 4.0 gammas per mile from north

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below, the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentine, which have a relatively high iron content; but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced by the geologist into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.

GEOPHYSICS PAPER 107

DESKENATLATA LAKE SOUTH

NORTHWEST TERRITORIES