

QUEENS AND SHELBURNE COUNTIES NOVA SCOTIA

20 gammas

10 gammas

Flight line

Magnetic depression

Flight altitude: 1000 feet above ground level

Scale: One Inch to One Mile = $\frac{1}{63,360}$

Air photographs covering this map-area may be obtained through the National Air Photographic Library, Topographical Survey, Ottawa, Ontario.

Magnetic Survey, September and October, 1956, by Geophysics Division, Geological Survey of Canada, Department of Mines and Technical Surveys. No correction has been made for regional variation; this increases at the rate of 5.4 gammas per mile from

east to west and 6.2 gammas per mile from south to

This map has been reprinted from a scanned version of the original map Reproduction par numérisation d'une carte sur papier

contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below, the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentine, which have a relatively high iron content; but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible.

> GEOPHYSICS PAPER 609 PORT MOUTON NOVA SCOTIA SHEET 20 P