
SHEET 21 G



OUEBEC Rimouski Gulf of St Lawrence Edmundston Newcastle Egrence Edmundston Newcastle Egrence Edmundston St Lawrence Edmundston Sydney Moncton Sydney Moncton Sydney Moncton Sydney Moncton Sydney Moncton St Lawrence Edmundston Sydney Moncton Syd

INDEX MAP

AEROMAGNETIC SERIES

Isomagnetic lines (total field)
500 gammas
100 gammas
20 gammas
10 gammas
Magnetic depression contour
Flight line
Flight aktitude: 500 feet above ground level

## CANTERBURY

YORK AND CARLETON COUNTIES
NEW BRUNSWICK

		Scale	1:50,000		- 12°
Kilometres 1	0	1	2	3	4 Kilometres
Miles 1		0	1		2 Miles
		ational Air Pl	this map-area notographic Lib		

Magnetic Survey, January and March 1950, by Geophysics Section, Geological Survey of Canada, Department of Mines and Technical Surveys

No correction has been made for regional variation; this increases at the rate of 3.0 gammas per mile from east to west and 2.5 gammas per mile from south to north

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below, the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentine, which have a relatively high iron content; but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced by the geologist into, or across, areas of few or no outcrops. In many instances, however, no present interpretation of particular anomalies may be possible.