


The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below, the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentine, which have a relatively high iron content; but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced by the geologist into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.

Airborne Magnetic Survey, September 1953,
by Geophysics Section, Geological Survey of Canada,
Department of Mines and Technical Surveys.

No correction has been made for regional variation;
this increases at the rate of 4.0 gammas per mile from
east to west and 2.5 gammas per mile from south
to north.

Air photographs covering this map area may be obtained
through the National Air Photographic Library, Topographical
Survey, Ottawa, Ontario.



G
34101
-C93
1948
G4
omvsc
DAG
1955

612 ca f ~~sc~~

This document was produced
by scanning the original publication.
Ce document est le produit d'une
numérisation par balayage
de la publication originale.



11 ^F/₄