



ISOMAGNETIC LINES (absolute total field)

500 gammas

100 gammas

50 gammas

Magnetic depression

Bathymetric contours in feet below sea-level

200

The ship magnetometer survey was conducted between May and October 1961, and 1962 using Newfoundland Decca Chain 6 as a navigational aid. No correction has been made for regional variation.

The bathymetric contours were obtained from charts published by the Canadian Hydrographic Service, Marine Sciences Branch.

Drafting by Lockwood Survey Corporation Ltd.

MAP 232 G 2ND EDITION

LOUISBOURG

CAPE BRETON ISLAND

NOVA SCOTIA

Scale: One Inch to One Mile = $\frac{1}{63,360}$ Miles

This map has been reprinted from a scanned version of the original map. Reproduction par numérisation d'une carte sur papier.

ISOMAGNETIC LINES (total field)

500 gammas

100 gammas

20 gammas

10 gammas

Magnetic depression

Flight line

The airborne magnetic survey was conducted during September 1953 by the Geological Survey of Canada at a flight altitude of 1000 feet above ground level. No correction has been made for regional variation.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentinite, which have a relatively high iron content, but in special instances may be due, or partly due, to concentrations of magnetic minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.