


Joins Map 2507G, "Broken Lake"

Joins Map 2510G, "Roblin River"

Joins Map 2517G, "Gilliam Island"

Joins Map 2508G, "Panco Lake"

PUBLISHED, 1964



ISOMAGNETIC LINES (total field)

500 gammas . . . . .

100 gammas . . . . .

20 gammas . . . . .

10 gammas . . . . .

Magnetic depression . . . . .

Flight lines . . . . .

Flight altitude: 1000 feet above ground level.

MAP 2509G

# DEER ISLAND

MANITOBA

Scale: One Inch to One Mile =  $\frac{1}{63,360}$  Miles

Airborne Magnetic Survey, June to November, 1963, by Canadian Aero Service Ltd.

No correction has been made for regional variation.

The planimetry for this map was obtained from the topographical map sheet published at a scale of one inch to four miles.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below, the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentinite, which have a relatively high iron content; but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced by the geologist into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.