

A L A S K A Steephology WHITEHORSE OF A L A S K A Haines OF A L A S K A JUNEAU OF A L A S K A JUNEAU OF A L A S K A JUNEAU JUNE

ISOMAGNETIC LINES (absolute total field)

500 gammas
100 gammas
20 gammas
10 gammas
Magnetic depression

SEAGULL CREEK

MAP 1306G

YUKON TERRITORY

Scale: One Inch to One Mile = $\frac{1}{63,360}$ Miles

1 1/2 0 1 2

Air photographs covering this map-area may be obtained

through the National Air Photographic Library, Topographical

Survey Ottawa, Ontario.

Airborne Magnetic Survey, June to September, 1961, by Aero Surveys Ltd.

No correction has been made for regional variation.

The planimetry for this map was obtained from topographical map sheets, published by the Department of Mines and Technical Surveys.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentine, which have a relatively high iron content, but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.

PUBLISHED, 1962

GEOPHYSICS PAPER 1306

SEAGULL CREEK
YUKON TERRITORY

SHEET 105 B