


MAP 4179G

# LUNDAR MANITOBA

Scale: One Inch to One Mile =  $\frac{1}{63,360}$  Miles



The planimetry for this map was obtained from topographical map sheets published by the Department of Energy, Mines and Resources, Ottawa.

Airborne Magnetic Survey, March 1967 to October 1968 by Spartan Air Services Ltd.


No correction has been made for regional variation.

Where the survey aircraft traversed large areas of water and swamp, Doppler navigation was utilized to direct the course of the aircraft and the Doppler output was recorded on an incremental X, Y recorder for compilation purposes.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentinite, which have a relatively high iron content, but in special instances may be due, or partly due, to concentrations of magnetic minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.

GEOPHYSICS PAPER 4179

LUNDAR  
MANITOBA  
SHEET 62 <sup>J</sup>/<sub>9</sub>



INDEX MAP