



ISOMAGNETIC LINES (absolute total field)

500 gammas

100 gammas

20 gammas

10 gammas

Magnetic depression

Flight lines

Flight altitude 300 metres above ground level.

1 gamma = 10⁻⁹ tesla in SI units

MAP 8348G
46 ^L/₅ and 46 ^L/₆
DISTRICT OF KEEWATIN
NORTHWEST TERRITORIES

SCALE 1:50,000

Air photographs covering this map area may be obtained through the National Air Photographic Library, Topographic Survey, Ottawa, Ontario.

COPIES OF THIS MAP MAY BE OBTAINED FROM THE DIRECTOR-GENERAL, GEOLOGICAL SURVEY OF CANADA, OTTAWA.

Airborne magnetic survey, June and July 1976
by Geotrex Ltd., Survoir Ltd., Northway Survey Corporation Ltd.

No correction has been made for regional variation.

The topography for this map was reproduced from 1:250,000 topographical map sheets published by the Department of Energy, Mines and Resources, Ottawa.

Where the survey aircraft traversed large areas of water and ice, Doppler navigation was utilized to direct the course of the aircraft and the Doppler output was recorded on an incremental X, Y recorder for compilation purposes.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentinite, which have a relatively high iron content, but in special instances may be due, or partly due, to concentrations of magnetic minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced into, or across, areas of low or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.