



ISOMAGNETIC LINES (absolute total field)

1000 gammas . . . . .

100 gammas . . . . .

20 gammas . . . . .

10 gammas . . . . .

Magnetic depression . . . . .

Flight lines . . . . .

Flight altitude: 500 feet above ground level.

MAP 958 G

# SHERIDAN LAKE

KENORA DISTRICT  
ONTARIO

Scale: One Inch to One Mile =  $\frac{1}{63,360}$  Miles

Air photographs covering this map-area may be obtained through the National Air Photographic Library, Topographical Surveys, Ottawa, Ontario.

Airborne Magnetic Survey, May, 1959 to March, 1960, by Spartan Air Services Ltd.

No correction has been made for regional variation.

The planimetry for this map was obtained from a provisional map prepared on contract by Hunting Survey Corporation Limited.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below, the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentine, which have a relatively high iron content; but in special instances may be due, or partly due, to concentrations of magnetic ore minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced by the geologist into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.