


ISOMAGNETIC LINES (total field)

500 gammas
 100 gammas
 20 gammas
 10 gammas
 Magnetic depression

Flight lines
 Flight altitude 1000 feet above ground level

MAP 3902G
43 F/2
 DISTRICT OF KENORA
 ONTARIO

Scale: One Inch to One Mile = $\frac{1}{63,360}$
 Miles



Airborne Magnetic Survey, September 1966 to September 1968
 by Lockwood Survey Corporation Ltd.

The planimetry for this map was obtained from
 topographical map sheets published by the
 Department of Energy, Mines and Resources, Ottawa.

No correction has been made for regional variation.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentinite, which have a relatively high iron content, but in special instances may be due, or partly due, to concentrations of magnetic minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.