

## NOMANSLAND POINT

MAP 2332G

KEEWATIN DISTRICT – COCHRANE DISTRICT
NORTHWEST TERRITORIES – ONTARIO

Scale: One Inch to One Mile =  $\frac{1}{63,360}$ Miles

2

Airborne Magnetic Survey, May 1963 to April 1964 by Spartan Air Services Ltd.

No correction has been made for regional variation.

The planimetry for this map was obtained from topographical map sheets published by the Department of Mines and Technical Surveys and the Ontario Department of Lands and Forests.

The magnetic data on this map were compiled from information recorded along the flight lines shown. The anomalies expressed by the magnetic contours are dependent on the variable magnetic intensities of the underlying rocks, and may be due to conditions near, or at unknown depths below the surface. High magnetic anomalies normally indicate the presence of basic rocks, such as diabase, gabbro, or serpentinite, which have a relatively high iron content, but in special instances may be due, or partly due, to concentrations of magnetic minerals. By means of the magnetic anomalies, various rock bodies or structural features, such as faults or folds, may be traced into, or across, areas of few or no outcrops. In many instances, however, no interpretation of particular anomalies may be possible without further geological information.

SHEET 43A3

81°00'

GEOPHYSICS PAPER 2332

NOMANSLAND POINT
NORTHWEST TERRITORIES - ONTARIO

SHEET 43A