

BJARNI FORMATION: WELL TOPS AND THICKNESS

Well	Formation ¹ Top (BKB; m)	Thickness ¹ (m)	Net Sand ² (m)	Net Shale ² (m)	Net Other (m)	Sand/ Shale	% Sand	% Shale
Hare Bay E-21	NP ³	0.0	0.0	0.0	0.0	-	-	-
Verrazano L-77	NP	0.0	0.0	0.0	0.0	-	-	-
Freydis B-87	1788.0	117.0	40.0	75.0	2.0	0.53	34.19	64.10
Leif M-48	NP	0.0	0.0	0.0	0.0	-	-	-
Leif E-38	U ⁴	U	U	U	U	U	U	U
Indian Harbour M-52	NP	0.0	0.0	0.0	0.0	-	-	-
North Leif I-05	2820.0	574.5	90.0	470.5	14.0	0.19	15.68	81.88
Cartier D-70	NP	0.0	0.0	0.0	0.0	-	-	-
Roberval C-02	NP	0.0	0.0	0.0	0.0	-	-	-
Roberval K-92	3080.0	107.0	44.0	63.0	0.0	0.69	41.12	58.88
Gudrid H-55	NP	0.0	0.0	0.0	0.0	-	-	-
Bjarni O-82	2285.0	365.0 ⁺	313.0 ⁺	52.0	0.0	6.02	85.75	14.25
Bjarni H-81	2150.5	105.0	99.0	6.0	0.0	6.50	94.29	5.71
Herjolf M-92	2614.0	1137.0	1039.0	84.0	14.0	12.37	91.38	7.39
North Bjarni F-06	2423.0	389.0 ⁺	349.0 ⁺	40.0	0.0	8.70	89.72	10.28
Tyrk P-100	1195.0	328.0	74.0	251.0	3.0	0.29	22.56	76.52
South Labrador N-79	3496.0	52.0	10.0	42.0	0.0	0.24	19.23	80.77
South Hopedale L-39	1920.0	88.0	0.0	88.0	0.0	-	0.00	100.00
Hopedale E-33	1920.0	60.0	23.0	37.0	0.0	0.62	38.33	61.67
Corte Real P-85	U	U	U	U	U	U	U	U
Snorri J-90	3027.0	121.5	26.0	68.5	27.0	0.38	21.40	56.38
Ogmund E-72	1712.0	1372.0	1211.0	137.0	24.0	9.18	88.26	9.98
Skolp E-07	2472.0	495.5	246.5	218.0	31.0	1.13	49.75	44.00
Pothurst P-19	U	U	U	U	U	U	U	U
Karlsefni A-13	NP	0.0	0.0	0.0	0.0	-	-	-
Gilbert F-53	NP	0.0	0.0	0.0	0.0	-	-	-
Rut H-11	U	U	U	U	U	U	U	U

¹ Formation top and thickness determined by P. N. Moir.

² Lithology net thickness extracted from master logs of P. E. Miller and G. J. D'Eon (1987).

³ NP = not present

⁴ U = unknown

FORMATION DE BJARNI: TOITS DES Puits DE FORAGE ET EPAISSEURS

Puits	Formation ¹ Toit (BKB; m)	Epaisseur ¹ (m)	Sable Pur ² (m)	Argilite Pure ² (m)	Autres Purs (m)	Sable/ Argilite	% Sable	% Argilite
Hare Bay E-21	NP ³	0.0	0.0	0.0	0.0	-	-	-
Verrazano L-77	NP	0.0	0.0	0.0	0.0	-	-	-
Freydis B-87	1788.0	117.0	40.0	75.0	2.0	0.53	34.19	64.10
Leif M-48	NP	0.0	0.0	0.0	0.0	-	-	-
Leif E-38	U ⁴	U	U	U	U	U	U	U
Indian Harbour M-52	NP	0.0	0.0	0.0	0.0	-	-	-
North Leif I-05	2820.0	574.5	90.0	470.5	14.0	0.19	15.68	81.88
Cartier D-70	NP	0.0	0.0	0.0	0.0	-	-	-
Roberval C-02	NP	0.0	0.0	0.0	0.0	-	-	-
Roberval K-92	3080.0	107.0	44.0	63.0	0.0	0.69	41.12	58.88
Gudrid H-55	NP	0.0	0.0	0.0	0.0	-	-	-
Bjarni O-82	2285.0	365.0 ⁺	313.0 ⁺	52.0	0.0	6.02	85.75	14.25
Bjarni H-81	2150.5	105.0	99.0	6.0	0.0	6.50	94.29	5.71
Herjolf M-92	2614.0	1137.0	1039.0 ⁺	84.0	14.0	12.37	91.38	7.39
North Bjarni F-06	2423.0	389.0 ⁺	349.0	40.0	0.0	8.70	89.72	10.28
Tyrk P-100	1195.0	328.0	74.0	251.0	3.0	0.29	22.56	76.52
South Labrador N-79	3496.0	52.0	10.0	42.0	0.0	0.24	19.23	80.77
South Hopedale L-39	1920.0	88.0	0.0	88.0	0.0	-	00.00	100.00
Hopedale E-33	1920.0	60.0	23.0	37.0	0.0	0.62	38.33	61.67
Corte Real P-85	U	U	U	U	U	U	U	U
Snorri J-90	3027.0	121.5	26.0	68.5	27.0	0.38	21.40	56.38
Ogmund E-72	1712.0	1372.0	1211.0	137.0	24.0	9.18	88.26	9.98
Skolp E-07	2472.0	495.5	246.5	218.0	31.0	1.13	49.75	44.00
Pothurst P-19	U	U	U	U	U	U	U	U
Karlsefni A-13	NP	0.0	0.0	0.0	0.0	-	-	-
Gilbert F-53	NP	0.0	0.0	0.0	0.0	-	-	-
Rut H-11	U	U	U	U	U	U	U	U

¹ Toit et épaisseur des formations par P. N. Moir.

² L'épaisseur des formations lithologiques provient des enregistrements de P. E. Miller et G. J. D'Eon (1987).

³ NP = absent

⁴ U = inconnu