

**KENAMU FORMATION: WELL TOPS AND THICKNESS**

Well	Formation Top <sup>1</sup> (BKB; m)	Thickness <sup>1</sup> (m)	Net Sand <sup>2</sup> (m)	Net Shale <sup>2</sup> (m)	Net Other <sup>2</sup> (m)	Sand/ Shale	% Sand	% Shale
Hare Bay E-21	1938.0	1107.0	45.0	1055.0	7.0	0.04	4.65	95.30
Verrazano L-77	NP <sup>3</sup>	0.0	0.0	0.0	0.0	-	-	-
Freydis B-87	860.0	527.0	104.0	407.0	16.0	0.25	19.73	77.23
Leif M-48	1222.3	443.4	56.0	382.4	5.0	0.14	12.63	86.24
Leif E-38	U <sup>4</sup>	U	U	U	U	U	U	U
Indian Harbour M-52	2002.5	910.5	11.0	869.5	30.0	0.01	1.20	95.49
North Leif I-05	1476.0	665.0	46.0	603.0	16.0	0.08	6.92	90.67
Cartier D-70	1275.3	437.7	77.0	353.7	7.0	0.22	17.59	80.81
Roberval C-02	1690.5	506.0	46.0	454.0	6.0	0.10	9.09	89.72
Roberval K-92	1815.0	541.0	4.0	520.0	17.0	0.01	0.73	96.12
Gudrid H-55	1679.0	501.0	28.0	470.0	3.0	0.06	5.59	93.81
Bjarni O-82	1360.0	520.0	28.0	486.0	6.0	0.06	5.38	93.46
Bjarni H-81	1334.0	486.0	6.0	476.0	4.0	0.01	1.23	97.94
Herjolf M-92	1398.0	565.0	32.0	520.0	13.0	0.06	5.66	92.03
North Bjarni F-06	1480.5	578.5	46.0	509.5	23.0	0.09	7.95	88.07
Tyrk P-100	762.0	322.0	4.0	315.0	3.0	0.01	1.24	97.83
South Labrador N-79	1447.0	985.0	9.0	967.0	9.0	0.01	0.91	98.17
South Hopedale L-39	1062.0	399.0	0.0	396.0	3.0	0.00	0.00	99.29
Hopedale E-33	982.0	611.0	30.0	564.0	17.0	0.05	4.91	92.31
Corte Real P-85	3379.0	1172.0	49.0	1117.0	6.0	0.04	4.18	95.31
Snorri J-90	1715.0	776.0	42.0	721.0	13.0	0.06	5.41	92.91
Ogmund E-72	1121.5	336.5	12.0	322.5	2.0	0.04	3.56	95.84
Skolp E-07	855.0	119.0	40.0	79.0	0.0	0.51	33.61	66.39
Pothurst P-19	3184.0	792.0	184.0	608.0	0.0	0.30	23.23	76.77
Karlsefni A-13	2191.0	845.0	166.0	678.0	1.0	0.25	19.64	80.24
Gilbert F-53	1527.0	542.0	4.0	530.0	8.0	0.01	0.74	97.77
Rut H-11	1322.0	794.0	8.0	784.0	2.0	0.01	1.01	98.74

<sup>1</sup> Formation top and thickness determined by P. N. Moir.

<sup>2</sup> Lithology net thickness extracted from master logs of P. E. Miller and G. J. D'Eon (1987).

<sup>3</sup> NP = not present

<sup>4</sup> U = unknown

**FORMATION DE KENAMU: TOITS DES PUIITS DE FORAGE ET EPAISSEURS**

Puits	Formation Toit <sup>1</sup> (BKB; m)	Epaisseur <sup>1</sup> (m)	Sable Pur <sup>2</sup> (m)	Argilite Pur <sup>2</sup> (m)	Autres Pur <sup>2</sup> (m)	Sable/ Argilite	% Sable	% Argilite
Hare Bay E-21	1938.0	1107.0	45.0	1055.0	7.0	0.04	4.65	95.30
Verrazano L-77	NP <sup>3</sup>	0.0	0.0	0.0	0.0	-	-	-
Freydis B-87	860.0	527.0	104.0	407.0	16.0	0.25	19.73	77.23
Leif M-48	1222.3	443.4	56.0	382.4	5.0	0.14	12.63	86.24
Leif E-38	U <sup>4</sup>	U	U	U	U	U	U	U
Indian Harbour M-52	2002.5	910.5	11.0	869.5	30.0	0.01	1.20	95.49
North Leif I-05	1476.0	665.0	46.0	603.0	16.0	0.08	6.92	90.67
Cartier D-70	1275.3	437.7	77.0	353.7	7.0	0.22	17.59	80.81
Roberval C-02	1690.5	506.0	46.0	454.0	6.0	0.10	9.09	89.72
Roberval K-92	1815.0	541.0	4.0	520.0	17.0	0.01	0.73	96.12
Gudrid H-55	1679.0	501.0	28.0	470.0	3.0	0.06	5.59	93.81
Bjarni O-82	1360.0	520.0	28.0	486.0	6.0	0.06	5.38	93.46
Bjarni H-81	1334.0	486.0	6.0	476.0	4.0	0.01	1.23	97.94
Herjolf M-92	1398.0	565.0	32.0	520.0	13.0	0.06	5.66	92.03
North Bjarni F-06	1480.5	578.5	46.0	509.5	23.0	0.09	7.95	88.07
Tyrk P-100	762.0	322.0	4.0	315.0	3.0	0.01	1.24	97.83
South Labrador N-79	1447.0	985.0	9.0	967.0	9.0	0.01	0.91	98.17
South Hopedale L-39	1062.0	399.0	0.0	396.0	3.0	0.00	0.00	99.29
Hopedale E-33	982.0	611.0	30.0	564.0	17.0	0.05	4.91	92.31
Corte Real P-85	3379.0	1172.0	49.0	1117.0	6.0	0.04	4.18	95.31
Snorri J-90	1715.0	776.0	42.0	721.0	13.0	0.06	5.41	92.91
Ogmund E-72	1121.5	336.5	12.0	322.5	2.0	0.04	3.56	95.84
Skolp E-07	855.0	119.0	40.0	79.0	0.0	0.51	33.61	66.39
Pothurst P-19	3184.0	792.0	184.0	608.0	0.0	0.30	23.23	76.77
Karlsefni A-13	2191.0	845.0	166.0	678.0	1.0	0.25	19.64	80.24
Gilbert F-53	1527.0	542.0	4.0	530.0	8.0	0.01	0.74	97.77
Rut H-11	1322.0	794.0	8.0	784.0	2.0	0.01	1.01	98.74

<sup>1</sup> Toit et épaisseur des formations par P.N. Moir.

<sup>2</sup> L'épaisseur des formations lithologiques provient des enregistrements de P. E. Miller et G. J. D'Eon (1987).

<sup>3</sup> NP = absent

<sup>4</sup> U = inconnu