



MAP UNIT*	NAME	MATERIAL	TYPICAL THICKNESS (m)	LANDFORM	GENERAL COMMENTS
FO	ORGANIC DEPOSITS	Fen, locally includes marsh, swamp, and shallow water classes of Tarnocel (1980)	0.4-1.5	Flat to gently sloping except for low hummocks and ridges	Occurrence of permafrost sporadic, in general more prevalent with increasing elevation; in some areas mapped as FO, organic accumulation is less than 0.4 m and hence does not meet the thickness criterion for organic deposits and hence should be considered as an organic soil horizon
		Alluvial plain, thermokarst alluvial plain	5-10	Flat to gently irregular; floodplains of larger streams commonly have distinct meander scrolls and oxbow lakes; thermokarst alluvial plains have irregular thermokarst ponds and depressions 2 to 3 m below the alluvial plain surface	Flooding common to infrequent; permafrost absent beneath active channels of permanent streams but common elsewhere; permafrost occurs throughout thermokarst alluvial plains except beneath ponds and wet depressions; with segregated ground ice to 50% by volume
Apt	ALLUVIAL DEPOSITS	Alluvial plain and alluvial terrace, undivided	5-10	Flat to gently irregular	Flooding common to infrequent on floodplain, rare on bordering terrace; permafrost common
Al		Alluvial fan or fan apron	2-20	Gently to steeply sloping fans occur where high gradient tributary streams meet lower gradient trunk streams; fans aprons are formed by coalescence of small fans associated with intermittent streams and rills	Streams on fans are subject to sudden and damaging changes in course (avulsions); fans comprising mainly gravel and coarse sand may be free of permafrost, or if perennially frozen are thaw-stable; fans comprising mainly silt are typically perennially frozen; the silt may be loesslike and unstable when thawed
At		Alluvial complex, combinations of Ap, Apt, and Al	2-20	Various	See Ap, Apt, Al as applicable
Lp, Lpk	GLACIOLACUSTRINE DEPOSITS	Glaciolacustrine plain	2-20	Generally flat but commonly with 25% or more of area occupied by thermokarst ponds or depressions 1 to 4 m below the general surface (Lpk)	Permafrost present throughout except beneath thermokarst ponds; 15 to 50% segregated ice by volume; highly unstable when thawed
Lb, Lbg		Glaciolacustrine blanket, gullied glaciolacustrine blanket	2-5	Gently to moderately sloping, commonly blanketing lower slopes of valleys and wedging out upslope	Mostly perennially frozen; limited data suggest that content of segregated ice is typically low, but high ice contents are possible; highly susceptible to erosion
Lv		Glaciolacustrine veneer	0-2	This layer conforming to surface of subjacent unit (commonly Mb)	Permafrost common, ice content variable
Gp, Gpc, Gpd	GLACIOFLUVIAL DEPOSITS	Glaciofluvial plain, channelled glaciofluvial plain (McConnell age, Reid age)	5-30	Flat to very gently sloping, commonly with shallow anastomosing channels Gp-c	Permafrost not common, but where present, deposits are mostly thaw-stable; constitute main source of aggregate for construction purposes; good drainage and soil stability make the units suitable for location of most types of facilities; areas of Gp, Gd are preferred for airstrips; or other installations requiring large areas of well-drained stable terrain; generally high permeability permits use of septic disposal systems
Gt, Gtr		Glaciofluvial terrace (McConnell age, Reid age)	2-20	As for Gp, but in terrace position adjacent to a major stream	
Gh, Ghr		Hummocky or ridged glaciofluvial deposits	5-30	Hummocky (including kames) or ridged (including eskers and esker complexes)	
Gi, Gir		Glaciofluvial delta, glaciofluvial fan	2-30	Glaciofluvial deltas have typical delta form with flat top and steep outer slope; glaciofluvial fans have characteristic fan form, with moderate slope	
Gx, Gxr	GLACIOFLUVIAL COMPLEXES OF Gp, Gt, Gh, G, Or (McConnell age, Reid age)	Glaciofluvial complex, combinations of Gp, Gt, Gh, G, Or (McConnell age, Reid age)	2-30	Various, includes areas that would be classed as Gp or Gt except for presence of kettles	
Md		Drumlinoid or fluted till plain	2-50	Till plain with individual drumlins and/or distinct glacial fluting	Permafrost sporadic; commonly drumlin crests and elevated parts of flutings are permafrost free or have a thick active layer, whereas associated silt and/or organic deposits are likely to be perennially frozen and may be ice-rich; (Construction of linear facilities such as roads would be much easier parallel to rather than across the grain of the topography)
Mvd		Drumlinoid moraine veneer	0-2	Till veneer over bedrock, with drumlinoid or crag-and-tail topography	As for Mv (below)
Mb, Mbr, Mb-g, Mb-g		Moraine blanket, channelled moraine blanket, gullied moraine blanket (McConnell age, Reid age)	2-10	Gently to moderately sloping, conforming broadly to topography of subjacent bedrock	Permafrost sporadic; common on northerly facing slopes; suitable for conventional cut-and-fill road construction where permafrost is lacking
M, Mh, Mv, Mv-c	GLACIAL DEPOSITS	Moraine-colluvial blanket	2-10	Till and colluvium	As for Mb
Mv, Mv-c		Moraine veneer, channelled moraine veneer	0-2	As for Mb; bedrock common in channels in Mv-c	Permafrost sporadic; engineering capabilities and limitations determined in large part by the lithology and topography of the subjacent bedrock (Mv-c)
M, Cv		Moraine-colluvial veneer	0-2	Till and colluvium	As for Mb
Mh, Mr, Mf, Mf-g	GLACIAL DEPOSITS	Hummocky moraine, ridged moraine (McConnell age, Reid age)	2-50	Hummocks and ridges with slopes to 30° and relief to 20 m (exceptionally 40 m) superposed on flat to moderately sloping surfaces; locally, ridges included in Mh are lowes of large moulton flows formed during early retreat stage of McConnell Glaciation by slope failures where lateral moraines formed embankments on steep slopes	Few data on distribution of permafrost; vegetation cover suggests general absence of permafrost or presence of unusually thick active layer; locally efforts well drained road locations sites and constitutes source of common fill
Mh, Mf, Mf-g		Moraine complex, combinations of Mh, Mr, Mf, Mf-g (McConnell age, Reid age)	2-50	As for Mh; up to 2 m silt and/or organic deposits common in depressions	
Mm, Mm-g		Rolling moraine (McConnell age, Reid age)	5-50	Broad hummocks 10-30 m high and 100 to 300 m across, slopes to 12°	No data on distribution of permafrost or prevalence of ground ice, probably similar to Mh, Mb
Cb	COLLUVIAL DEPOSITS	Colluvial blanket	2-5	Any of the deposits described above, plus bedrock detritus, modified and/or interstratified as a result of downslope movement of material; texture ranges from coarse blocky bedrock detritus of mountain fans to clayey or silty diamict of some lower slopes, locally includes talus and/or moulton deposits, on many slopes there is a downslope transition from material that has been moved by gravity (Cb) to material that has been transported by water. Extensive slopes have therefore been designated as Cb/W or Af/Cb	Permafrost sporadic; widespread on northerly facing slopes and on high plateaus and periglacial features, including small solifluction lobes, sorted polygons, and felsite areas; conspicuous and widespread above treeline (1280 to 1370 m, 4200 to 4500 ft.)
C, Mb		Colluvial-moraine blanket	2-5	Colluvium and till	As for Cb
Cv		Colluvial veneer	0-2	Gently to highly irregular, conforming to topography of subjacent material (usually bedrock)	As for Cb; engineering properties and limitations determined in large part by the character of subjacent material (usually bedrock)
C, Mv		Colluvial-moraine veneer	0-2	Colluvium and till	As for Cv
Cz	Rockslides	Blocky bedrock detritus	10-50	Hummocky	The few occurrences in the area are associated with bedrock slopes that are locally oversteeped by glacial erosion or deep incision of ice marginal channels
R, Rc	ROCK	Bedrock, channelled bedrock	Various	Various	Permafrost sporadic; ground ice low to completely lacking
Rc		Bedrock, unchannelled bedrock	Various	Various	

*The most commonly occurring units are shown above; for others, refer to Explanation of Map Unit Designations, coloured legend blocks indicate primary map units that appear on this map

EXPLANATION OF MAP UNIT DESIGNATIONS

SIMPLE MAP UNITS
A simple map unit designation consists of a generic symbol (upper case letter) followed by one or more morphologic descriptors (lower case letters). The range of material textures to be expected within a map unit is indicated under "MATERIAL". Where the texture of the material is known more specifically, it is indicated by one or more textural prefixes (lower case letter).

MIXED UNITS
Two types of mixed units are used:
a. Combinations of the form "Gp/Mb" indicate that two distinct types of deposit are distinguishable within the boundaries of the unit, but cannot be differentiated because of map scale. The first named deposit type occupies more than 50% of the delineated area, the second named less than 50%, but more than 10%. Deposit types that constitute less than 10% of the delineated area are ignored.
b. Combinations of the form "M, Cv", "M, C, L" etc., indicate that two or more distinct generic classes of deposit are known to occur or suspected to occur within a delineated area. The respective classes of deposit lack distinctive landforms that would permit differentiation by aphotographic interpretation, and differentiation on the ground has not been undertaken. The order in which the respective classes are listed indicates the likely prevalence by area of each class within the delineated area.

TEXTURAL MODIFIERS
f - fan
c - site
h - hill
s - sand
g - gravel

GENERIC CATEGORIES
O - Organic deposits
A - Alluvial deposits
C - Glaciofluvial deposits
L - Glaciolacustrine deposits
M - Moraine deposits
R - Bedrock

MORPHOLOGIC MODIFIERS
b - blanket (generally 2m thick)
d - drumlinoid
f - fanlike
h - hummocky
i - irregular
m - miring
r - ridge
t - terrace
v - veneer (generally 2m thick)
w - complex combinations of permafrost
x - xeric

EXCESSIVE MODIFIERS
c - channelled
g - gullied
k - kettle in the form
p - pond
s - scarp

REFERENCES
Tarnocel, C. 1980. Canadian wetland registry. In Proceedings of a Workshop on Canadian Wetlands (Saskatoon 1979). C.D.A. Rubec and F.C. Pollett (ed.), Environment Canada, Land Directorate, Ecological Land Classification, Series 12.

MAP 22-1987
SURFICIAL GEOLOGY
STEVENS LAKE
YUKON TERRITORY

Scale 1:100 000 - Échelle 1/100 000

UNIVERSAL TRANSVERSE MERCATOR PROJECTION
Projection transverse universelle de Mercator
© Crown copyrights reserved / © Droits de la Couronne réservés

Geology by O.L. Hughes, 1966, 1967, 1979, based mainly on a photore interpretation with limited ground checking

Thematic information on this map is in part reproduced directly from author's copy

Geological cartography by the Geological Survey of Canada

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Library / Bibliothèque
DEC 12 1989
GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA

Base map assembled by the Geological Survey of Canada from maps published at 1:50 000 scale by the Surveys and Mapping Branch in 1968, 1969, 1985

Copies of the topographical editions covering this map area may be obtained from the Canada Map Office, Department of Energy, Mines and Resources, Ottawa, Ontario, K1A 0E9

Mean magnetic declination 1988, 29°20' East, decreasing 15.6' annually. Readings vary from 29°02' E in the SW corner to 29°38' E in the NE corner of the map

Elevations in feet above sea level

22-1987

