MINISTÈRE DE L'ÉNERGIE. DES MINES ET DES RESSOURCES

LEGEND

SOUTHWEST OF TINTINA FAULT

MID-CRETACEOUS

Kg resistant, grey weathering, locally foliated, biotite + hornblende diorite, granodiorite, and granite

Between St.Cyr and Tintina Faults

TERTIARY

Tv, undivided; Tv1, small stocks and necks of white weathering, flow-banded, rhyolitic, quartz-sanidine porphyry; Tv2, laminated rhyolitic ash-flow tuffs and flows; Tv3, dark grey weathering, locally amygdaloidal, dark grey-green basalt necks and flows; Tv4. nassive quartz-feldspar porphyry

Ts recessive, thick bedded to massive, pebble to boulder chert-quartz conglomerate, chert sandstone and thin bedded, dark brown siltstone and shale

CARBONIFEROUS

Csl recessive, dark grey weathering, thin bedded, interlaminated, buff-yellow siltstone and brown argillite, strongly bioturbated brown argillite, strongly bioturbated

MISSISSIPPIAN

Mtf resistant, rusty-orange weathering, thin bedded, apple green and dark grey chert and cherty tuff

DEVONIAN AND MISSISSIPPIAN

DMs recessive, black weathering with rusty streaks, thin bedded, black siliceous slate and minor interbedded, chert wacke and chert-granule grit

DMcsi buff-orange weathering <u>phyllite</u>, calcareous phyllite and phyllitic, platy <u>limestone</u>; minor buff weathering dolomitic <u>siltstone</u>, black very finely crystalline limestone, <u>black</u> pyritic slate, and fine grained quartz arenite; DMcsl1, blue-grey weathering, calcareous, medium grained quartz arenite

uDcg resistant, grey-orange weathering, massive, limestone conglomerate (not shown on stratigraphic cross-section; relations to other map units uncertain)

ORDOVICIAN TO DEVONIAN

ODSI moderately resistant, black weathering, siliceous, graphitic, black siliceous and pyritic slate

CAMBRO-ORDOVICIAN

buff, orange and orange-brown weathering, thinly interlaminated calcareous shale and silty limestone

Southwest of St. Cyr Fault

UPPER DEVONIAN AND? MISSISSIPPIAN

DMs recessive, black weathering with rusty streaks, thin bedded, black siliceous slate and minor interbedded chert wacke and chert-granule grit

SILURIAN AND DEVONIAN

ASKIN GROUP (SDdq, Sst)

SDdq resistant, medium grey to buff weathering, medium to thick bedded dolomite, sandy dolomite and quartz arenite

Sst tan weathering, thin bedded, dolomitic, platy siltstone

CAMBRO-ORDOVICIAN

KECHIKA GROUP (OSs. uEOsl)

PRECAMBRIAN AND LOWER CAMBRIAN

OSs black weathering, thin bedded, black graptolitic shale (only preserved locally)

ucosi grey-orange weathering, finely interlaminated, calcareous phyllite and grey limestone;

abundant quartz-carbonate veins and pods

PEp dark brown weathering, non-calcareous, dark blue-grey slate, siltstone and minor limestone; metamorphosed equivalents near Glenlyon Batholith are biotite-muscovite-quartz schist and minor marble; Pep1, buff-orange to light grey weathering, thin bedded, argillaceous limestone and calcareous phyllite

_____ Limit of outcrop _____ Geological boundary (defined, approximate, assumed, extrapolated beneath overburden where exposure warrants) + + + - Bedding (horizontal, inclined, vertical, overturned, tops unknown) Foliation (inclined, vertical) ← ← Wrinkle lineation, axis of small scale fold (inclined, horizontal) Fault, steeply dipping (defined, approximate, assumed, extrapolated beneath overburden; barb on downthrown side) _____ overburden, overturned; teeth on upper plate)

Fault, thrust (defined, approximate, assumed, extrapolated beneath Fault, transcurrent (defined, approximate, assumed, extrapolated beneath overburden; arrows indicate slip) Anticline (defined, approximate, assumed, extrapolated beneath

Syncline (defined, approximate, assumed, extrapolated beneath overburden) Anticline, syncline (overturned) Mineral occurrence (showing, work target)

→ Fossil locality

1) contacts are extrapolated, where exposure warrants, on basis of assumed simple structure 2) mineral occurrence numbers follow convention in Yukon Exploration 1987, Exploration and

NOTES

Outcrop not present, map unit inferred (italic map unit symbols)

Geological Services Division, Dept. Indian and Northern Affairs, Yukon

3) Jones Lake, Mt. Christie, Prevost, Portrait Lake, Steel, Gull Lake, Narchilla and Yusezyu formations and Hyland Group are new lithologic names approved and accepted for publication by the Geological Survey (Gordey, in press).

LEGEND FOR OPEN FILES 2249, 2250, 2251

NORTHEAST OF TINTINA FAULT

TERTIARY

Tv, undivided; Tv1, small stocks and necks of white weathering, flow-banded, rhyolitic, quartz-sanidine porphyry; Tv2, laminated rhyolitic ash-flow tuffs and flows; Tv3, dark grey weathering, locally amygdaloidal, dark grey-green basalt necks and flows; Tv4, massive quartz-feldspar porphyry

recessive, thick bedded to massive, pebble to boulder chert-quartz conglomerate, chert sandstone and thin bedded, dark brown siltstone and shale

MID-CRETACEOUS

South Fork Volcanics: dark brown weathering, locally columnar jointed, massive, densely welded, biotite-quartz-hornblende-feldspar crystal tuff

Selwyn Plutonic Suite: grey weathering, resistant, medium- to coarse grained, locally megacrystic (K-spar), biotite ± hornblende ± muscovite granite, quartz monzonite and granodiorite; Ks1, plutons without hornblende; Ks2, plutons with hornblende; Ks3, porphyritic biotite ± hornblende granite characterized by large smokey grey quartz phenocrysts and locally K-feldspar phenocrysts; Ks4, fine grained, mafic free, granite with disseminated pyrite

ALLOCHTHONOUS ASSEMBLAGES

PENNSYLVANIAN AND PERMIAN

Anvil Allochthonous Assemblage: CPav, resistant, dark weathering dark grey-green basalt, tuff, and breccia; CPat, thin bedded, grey-green, jasper-red and apple-green chert and siliceous tuff, and minor quartz-chert sandstone and shale; CPal, light grey weathering, massive, finely crystalline, dark grey limestone; CPaub, recessive, green weathering serpentinite

CARBONIFEROUS TO TRIASSIC

Nisutlin Allochthonous Assemblage: CTnm, grey weathering, muscovitic, quartz blastomylonite; recessive, muscovitic quartzite and quartz-muscovite-biotite+ glaucophane schist with local pods of eclogite; Ckncg, resistant, massive, poorly sorted, conglomerate with pebble to cobble size clasts of basalt, chert, mylonite, and

ALLOCHTHONS EMPLACED IN JURA-CRETACEOUS

AUTOCHTHON

Jones Lake Formation: brown weathering, medium- to thick-bedded, calcareous siltstone, sandstone and shale, ripple cross-laminated; massive light grey weathering, finely crystalline, dark grey limestone:

Pc Mount Christie Formation, lesistant, orange medium-bedded, light grey-green to black chert Mount Christie Formation: resistant, orange to buff weathering, thin- to

DEVONO-MISSISSIPPIAN

EARN GROUP (Mc, Ms, DMp, Dp, DMe)

DMe undivided Mc, DMp, minor Dp

recessive, dark brown weathering, thin- to medium-bedded, calcareous, dark grey to brown siltstone, sandstone and shale; thin to thick interbeds of finely crystalline, dark grey limestone; local light grey weathering, thick bedded to massive, dark grey,

Crystal Peak Formation: resistant, dark grey weathering, massive chert-pebble conglomerate and chert quartz sandstone; minor brown weathering, dark blue-grey

Prevost Formation: recessive, brown weathering, thin bedded, laminated, dark blue-grey to black slate and thin to thickly interbedded fine- to medium-grained chert-quartz arenite and wacke, and chert-pebble conglomerate; DMp1, resistant, coarse grained quartz sandstone

Portrait Lake Formation: black, gun-blue or silvery white weathering, thin bedded, siliceous, black siltstone, slate and chert; minor quartz arenite

SILURIAN AND DEVONIAN

Dc1, massive, medium grained <u>quartz arenite</u>; Dc2, light grey weathering, massive to thick bedded, finely crystalline <u>limestone and dolostone</u>, locally cherty

Ssp tan weathering, thin bedded, dolomitic, platy siltstone

ORDOVICIAN AND SILURIAN

ROAD RIVER GROUP (Ss. OSd)

OSr undivided Duo Lake and Steel formations (may include infolds of €Or and Dp)

Steel Formation: orange weathering, thin bedded, burrowed, dolomitic, grey-green mudstone, siltstone and chert; thin bedded black chert; rare black graptolitic shale

Duo Lake Formation: resistant, grey weathering, thin- to medium-bedded, light grey to black chert; recessive, gunsteel weathering, black graptolitic shale

CAMBRO-ORDOVICIAN

resistant, dark weathering, massive, locally pillowed, dark grey-green basalt, tuff and

resistant, dark grey weathering, massive to laminated, blocky, white to light grey quartzose siltstone and chert and rare black slate; strikingly laminated, very fine grained tuffaceous siltstone and chert; minor grey phyllitic limestone, calcareous phyllite, and greenstone

Rabbitkettle Formation: grey-buff weathering, laminated to thin bedded, locally nodular, shaly limestone to calcareous phyllite (includes tuffaceous phyllite and greenstone on south flank of Anvil Batholith)

LOWER CAMBRIAN

Gull Lake Formation: recessive, brown weathering, non-calcareous, dark grey to black slate and siltstone; metamorphosed equivalents near Anvil and Orchay batholiths include quartz-muscovite-biotite schist (+garnet, +sillimanite, +staurolite, +andalusite) and minor marble


PRECAMBRIAN AND LOWER CAMBRIAN

HYLAND GROUP (PCn. Pv)

Narchilla Formation: recessive, maroon weathering, interbedded maroon and apple-green slate; grey-brown weathering, medium- to thick-bedded quartz sandstone and quartz- pebble conglomerate


Yusezyu Formation: grey brown weathering, thin- to thick bedded, interbedded, quartz sandstone, local quartz pebble conglomerate, and grey-green to dark grey slate;
Py1, grey-white weathering, finely crystalline, dark grey limestone

STRATIGRAPHIC RELATIONS AND GEOLOGIC HISTORY



ACKNOWLEDGMENTS

The author thanks G. Jilsen and L. Pigage of Curragh Resources, Whitehorse, who discussed freely their knowledge of the geology near Faro. J.G. Abbott (Exploration and Geological Services, Indian and Northern Affairs Canada, Whitehorse), M.P. Cecile and L.E. Jackson provided discussion and some logistical coordination. T. Frakes (1982), D. Thorkelson (1983), B. Thomas (1985, 1986), S. Gareau (1986) and D. Rhys (1987) assisted in the field. Superb helicopter service was provided by John Witham (Trans North Air, Ross River). Paleontologic control was provided through fossil identifications by E.W. Bamber, B.S. Norford, and M.J. Orchard. This Open File benefited from review by C. Roots.



LOCATION

REFERENCES

Abbott, J.G., 1983, Geology of the Macmillan fold belt; Exploration and Geological Services Division, Department of Indian and Northern Affairs, Whitehorse, Y.T., Open File (1:50,000, 3 maps and legend) Campbell, R.B., 1967, Glenlyon map area; Geological Survey of Canada, Memoir 352 Cecile, M.P., ----, Geology of Niddery Lake map area (1050); Geological Survey of Canada, Open Files 1006 (1984), 1118 (1984), 1242 (1986), and 1326 (1986)(1:50,000 maps and legends for 1050/3 to 16) Gordey, S.P., 1981, Geology of Nahanni map area (1051), Yukon Territory and District of Mackenzie; Geological Survey of Canada, Open File 780 (1:50,0000 & 1:125,000 maps and legend)

Gordey, S.P., 1983, Thrust faults in the Anvil Range and a new look at the Anvil Range Group; Geological Survey of Canada, Current Research, Part A, Paper 83-1A, p. 225-227 Gordey, S.P., 1988, The South Fork Volcanics: mid-Cretaceous caldera fill tuffs in east-central Yukon; Geological Survey of Canada, Current Research, Part E, Paper 88-1E, p. 13-18

Gordey, S.P., in press, Evolution of the northern Cordilleran miogeocline, Nahanni map area (1051), Yukon Territory and District of Mackenzie: Geological Survey of Canada, Memoir 482 Gordey, S.P. and Irwin, S.E.B., 1987, Geology of Sheldon Lake (105J) and Tay River (105K) map areas, Yukon Territory; Geological Survey of Canada Map 19-1987 Hughes, J.D. and Long, D.G.F., 1980, Geology and coal resource potential of early Tertiary strata along

Tintina Trench, Yukon Territory; Geological Survey of Canada, Paper 79-32

INAC, 1988, Yukon exploration 1987; Exploration and Geological Services Division, Yukon, Indian and Northern Affairs, Canada Jackson, L.E., Jr., and Morrison, S.E., 1984, Terrain inventory, Sheldon Lake (105J), Yukon Territory;

Geological Survey of Canada, Open File 1033 (1:125,000 map and notes) Jackson, L.E., Jr., 1986, Terrain inventory, Tay River (105K), Yukon Territory; Geological Survey of Canada, Open file 1380 (1:125,000 map and notes) Jennings, D.S. and Jilson, G.A. 1986, Geology and sulphide deposits of the Anvil Range, Yukon; in Mineral

Deposits of the Northern Canadian Cordillera, Morin, J.A. (ed.), Canadian Institute of Mining and Metallurgy, Special Volume 37, p. 319-361. Pigage, L.C. and Anderson, R.G., 1985, The Anvil plutonic suite, Faro, Yukon Territory; Canadian Journal of Earth Sciences, v. 22, p. 1204-1216

Roddick, J.A. and Green, L.H., 1961a, Geology, Tay River, Yukon (105K); Geological Survey of Canada, Map Roddick, J.A. and Green, L.H., 1961b, Geology, Sheldon Lake, Yukon (105J); Geological Survey of Canada, Map 12-1961

Tempelman-Kluit, D.J., 1972, Geology and origin of the Faro, Vangorda and Swim concordant zinc-lead deposits, central Yukon Territory; Geological Survey of Canada, Bulletin 208

Tempelman-Kluit, D.J., 1979, Transported cataclasite, ophiolite and granodiorite in the Yukon: Evidence of arc-continent collision; Geological Survey of Canada, Paper 79-14

Tempelman-Kluit, D.J., 1977, Geology of Quiet Lake and Finlayson Lake map-areas, Yukon Territory (105F and G); Geological Survey of Canada, Open File 486 (1:250,000 maps and legend)

Wood, D.H. and Armstrong, R.L., 1982, Geology, chemistry and geochronometry of the Cretaceous South

Fork Volcanics, Yukon Territory; Geological Survey of Canada, Paper 82-1A, p. 309-316

COMPOSITE LEGEND FOR GSC OPEN FILES 2249, 2250, AND 2251

Open File 2249: Geology of Tenas Creek (105K/1), Swim Lakes (105K/2), and Faro (105K/3) map areas, Yukon Territory

Open File 2250: Geology of Mount Atherton (105K/4), Rose Mountain (105K/5), and Mount Mye (105K/6) map areas, Yukon Territory

Geology of Blind Creek (105K/7), Teddy Creek (105K/10), and Barwell Lake (105K/11) map areas, aYukon Territory

S.P. Gordev

(Nine 1:50,000 scale maps for the Tay River area (105K); information on these maps locally differs from and supercedes that on the preliminary 1:250,000 scale map of the Tay River area (Gordey and Irwin, 1987))

June, 1990

SHEET 1 OF 4 (\$12.00/SET)