



GEOLOGICAL SURVEY OF CANADA
 GEOSCIENCE BRANCH
 1972

**AIRBORNE RADIOACTIVITY MAP
 MONT-LAURIER
 QUEBEC**

INTERVAL (0.41-2.81 MW)

The gamma and beta activity shown on this map was obtained from an aerial survey of the Mont-Laurier region in Quebec, Canada, during the summer of 1972. The data were obtained from a low-altitude survey aircraft flying at an altitude of 2000 feet and equipped with a sodium iodide crystal detector. The data were processed by a computer and plotted on a map. The map shows the distribution of radionuclides in the environment. The data were obtained from a low-altitude survey aircraft flying at an altitude of 2000 feet and equipped with a sodium iodide crystal detector. The data were processed by a computer and plotted on a map. The map shows the distribution of radionuclides in the environment. The data were obtained from a low-altitude survey aircraft flying at an altitude of 2000 feet and equipped with a sodium iodide crystal detector. The data were processed by a computer and plotted on a map. The map shows the distribution of radionuclides in the environment.

Airborne radioactivity survey 1972
 A. G. Mackay, R. L. Gray, K. A. McPherson
 Geological Survey of Canada

Scale 1:50,000
 0 500 1000 1500 2000 2500 3000 3500 4000 4500 5000
 METERS

OPEN FILE
 1110 1972
 GEOLOGICAL SURVEY OF CANADA