NOTE: This legend is common to Open File 405, 406 MIDDLE ORDOVICIAN 35 TRENTON GROUP: limestone 34 BLACK RIVER GROUP: limestone 33 ROCKCLIFFE: sandstone, shale, limestone LOWER ORDOVOCIAN 32 OXFORD: dolomite 31 MARCH: sandstone, dolomite **30** NEPEAN: sandstone UPPER CAMBRIAN 29 THERESA: sandstone, dolomite HELIKIAN 25 Slate 28 28a Syenite 28b Hornblende syenite 24 Marble 28c Nepheline syenite 23 Conglomerate

27 27a Granite 27b Potassic granite 22 Andesite 21 Rhyolite

20 20a Amphibolite 20b Hornblende gneiss 19 Paragneiss 18 Migmatite, granitic gneiss

27c Biotitic potassic granite 27d Hornblende potassic granite

17 Amphibole-pyroxene gneiss

10 Hornblende gneiss **9** Biotitic quartzofeldspathic gneiss 8 Biotitic migmatite **7** Amphibole-hypersthene gneiss 6a Potassic granite
6b Biotitic potassic granite | Paragneiss

APHEBIAN OR HELIKIAN

16 16a Granodiorite 16b Granite

15a Migmatite 15b Biotitic migmatite 15c Biotitic quartzofeldspathic gneiss

14 Quartzofeldspathic gneiss, leucogranulite

13 Biotite gneiss
13b Hornblende-biotite gneiss

12 Hornblende-garnet-biotite gneiss

11 Sillimanite-garnet-biotite gneiss

3 Hornblende gneiss 2 Migmatite, granitic gneiss 1 Amphibole-hypersthene gneiss

Geology derived from the map 1334-A, Rivière Gatineau at the scale of 1: 1,000,000. Compiled by A.J. Bear, W.H. Poole and B.V. Sandford, 1971

Geological cartography by the Geological Survey of Canada

Base-map at the same scale published by the Mapping and Charting Establishment, M.C.E. 1958-61

Mean magnetic declination 1977, 11043.8' West decreasing 0.1' annually. Readings vary from 12033' in the S.E corner to 10°58.8' in the NW corner of the map area

Elevation in feet above mean sea-level

Geochemical Symbol and Data Presentation


The concentration of an element at a sample site is graphically represented as one of 15 symbols, if a sample was collected but there is no data available a dot is plotted. The symbols are symmetrically arranged so that they first increase in size to the eighth symbol and then increase in blackness to the fifteenth. The two small crosses at the low end of the scale are used to respectively denote concentrations below the analytical detection limit, or, in the data group containing the detection limit. The data are grouped on a semi-logarithmic scale, i.e. 1,2,5,10,20,50,100 etc. Five decades can be spanned and this arbitrary division has been chosen for the continuing Canada wide series of maps constituting the National Geochemical Reconnaissance. The choice of symbols and the data groups they represent for any specific element

is based on the histogram and cumulative frequency plot for the total survey data from one, or more contiguous, open file sheets covered in one field season. The eighth symbol is used for the model group as defined by the histogram, this group usually includes the median of the data as defined by the 0.5 (50%) point on the cumulative frequency plot. Some, or all, of the remaining 14 symbols are chosen so as to achieve an appropriate graphical impact. An example of all 15 symbols is given

The symbol maps, being based on the total survey data distributions, are unaffected by the availability of ever increasing levels of knowledge in bedrock and surficial geology, and other environmental factors. Therefore, the raw data symbol maps are only intended to assist the rapid inspection of the data for gross regional features. To fulfil the needs of a more specific and thorough interpretation, the raw symbol maps should be modified using the field and analytical data provided in the data listings and any other knowledge available. To assist in the appraisal and modification of the data in terms of the symbol map bedrock geology, a table of summary statistics and proposed threshold values for each mapped bedrock unit, or broad lithologic unit, again based on the total survey data, is presented below the histogram. This table can be used along, or in conjuction, with the sample location map and data listings to indicate above threshold samples where they occur on the map. In many instances, the table will also illustrate, more clearly than the map, the dependence of mean geochemical levels on bedrock type. It may often be also observed that whilst the total data appears to approximate a log-normal distribution the data for individual map or lithologic units appears to approximate a normal distribution. The proposed thresholds presented are believed to be useful in interpreting the data from a mineral exploration viewpoint. Locations of samples with concentrations in excess of the threshold for the rock unit they appear to be derived from, should be studied carefully. The above threshold concentration can be due to a wide range of geological and environmental factors, but one of these could be the presence of abnormal concentrations of the element in a form of interest to the mineral explorationist.


To comprehensively study an area, all available geological, environmental and recorded data should be utilized. The data separation by bedrock type can often be improved by constructing new data subsets and deriving local threshold levels based on the most detailed and up-to-date knowledge available.


The term reliability factor and value that appears below the table is an estimate of the reliability of the geochemcial map. On the basis of duplicate sampling 5% of all lakes sampled it can be stated that there is a 95% chance that if any lake is resampled and identical methods of sample preparation and analysis are used the new value will lie between X + RF and X x RF where X is the original value obtained. This factor takes into account variability due to both heterogeneity of the centre-lake bottom sediments and sample preparation and analytical causes.




Province of Ontario Ministry of Natural Resources Geological Branch


Department of Energy, Mines and Resources Geological Survey of Canada







NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 2-1976 MANGANESE IN LAKE SEDIMENTS CANADA-ONTARIO SUBSIDIARY AGREEMENT ON MINERAL EXPLORATION AND DEVELOPMENT









NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 2-1976 OPEN FILE 406

Resource Geophysics and Geochemistry Division

Geological Survey of Canada, Ottawa

Geochemistry and Federal-Provincial coordination by E.H.W. Hornbrook Field operation supervised by C.C. Durham Analytical chemistry by J.J. Lynch

Data monitoring by R.G. Garrett, N.G. Lund and D. Ellwood

Ontario, Geological Branch

Federal-Provincial coordination by K,D. Card

Contractors Sample preparation by Golder Associates Chemical analyses by Chemex Labs Ltd.

This map forms one of a series of 28 sheets released under Geological Survey of Canada, Open Files 405-406. The open files consist of data for 12 elements each, per cent loss on ignition and sample site location.

The data are also available in digital form. For further information please contact:

> The Director, Computer Science Centre, Department of Energy, Mines and Resources, Ottawa, Ontario K1A OE8