

Published, 1977

EQUIVALENT URANIUM (eU)
 MAP 35764G
BIG SAND LAKE
 MANITOBA

Uranium Reconnaissance Program Airborne Gamma-Ray Spectrometer Survey, 1976, flown and compiled by the consortium of Terra Surveys Ltd., consortium directors, Kenting Earth Sciences Ltd., and Northway Survey Corporation Ltd.

The topography for this series of maps was reproduced from (1:250,000) topographical map sheets published by the Department of Energy, Mines and Resources, Ottawa.

This map was compiled from airborne gamma-ray spectrometer data recorded digitally along the flight lines shown. The spectrometer, with 50 litres of sodium iodide (NaI(Tl)) detectors, recorded gamma radiation in four channels, with the following energy ranges:

- Channel 1 2.42 - 2.82 MeV
- Channel 2 1.66 - 1.86 MeV
- Channel 3 1.36 - 1.56 MeV
- Channel 4 0.40 - 2.85 MeV



Channels 1, 2 and 3 were centered on the 2.62 MeV ^{238}Pu photo peak, the 1.76 MeV Bi^{214} photo peak and on the 1.46 MeV K^{40} photo peak, respectively. Counts were accumulated in these channels and recorded at one second intervals. The terrain clearance was averaged and recorded at one second intervals. The detectors were thermally stabilized to minimize spectrum shift. The survey aircraft were flown at a planned survey altitude of 400 feet and at a ground speed between 150 mph and 240 mph.

The data were corrected for dead time, atmospheric changes in temperature, background radiation, spectral scattering and deviations of terrain clearance from the planned survey altitude. Corrected count rates from channels 1, 2 and 3 were converted to concentrations of equivalent thorium, equivalent uranium, and potassium, using conversion factors determined for each gamma-ray spectrometer used in the survey. The total count rates from channel 4 were converted to units of radioelement concentration. The conversion factors which differed among the 3 aircraft used, are approximately those listed below:

- Channel 1 1 ppm eTh = 6 cps
- Channel 2 1 ppm eU = 8 to 10 cps
- Channel 3 1 %K = 70 to 80 cps
- Channel 4 1 %K = 140 to 160 cps

Data were smoothed using 40 data points along the flight lines (rejecting values over water), gridded at 2.2 kilometer intervals along track and 5 kilometer intervals across track, and contoured.

The contoured values are surface radioelement concentrations averaged over areas of approximately 700,000 square meters. These areas generally include some outcrop, overburden, swamps and small bodies of water. Consequently the concentrations indicated by the contour map are generally lower than the concentration in bedrock.



COPIES OF THIS MAP MAY BE OBTAINED FROM THE DIRECTOR GENERAL, GEOLOGICAL SURVEY OF CANADA, OTTAWA.

This map has been reprinted from a scanned version of the original map
 Reproduction par numérisation d'une carte sur papier

