

**LEGEND**  
Note: This legend is common to National Geochemical Reconnaissance Map 5-1976, Open File 409; Map 6-1976, Open File 410 and Map 7-1976, Open File 411.

**QUATERNARY**

- 8 Glacial, lacustrine, and fluviatile gravel, sand, silt and clay
- 7 Plateau basalts, olivine basalts
- 6 Volcanic flow rocks with interbedded sedimentary rocks; 6a, conglomerate
- 5 CORYELL: alkalic plutonic rocks; porphyritic granite and rhyolite

**JURASSIC □ CRETACEOUS**

- 4 NELSON and VALHALLA: granitic plutonic rocks

**JURASSIC**


- 3 Mafic and ultramafic intrusive rocks, pyroxenite, hornblende

**PALEOZOIC (INCLUDING UPPER PROTEROZOIC AND TRIASSIC)**

- 2 Basaltic and andesitic; lavas, greenstone, tuff, quartzite, limestone, phyllite, sandstone and conglomerate
- 1 Argillite; 2a, quartzite, argillite, limestone, slate, schist, phyllite, sandstone and conglomerate

**PROTEROZOIC (SHUSWAP TERRANE)**

- 1 Gneiss, minor schist, limestone, marble, dolomite, slate, phyllite; 1a, schist, quartzite, limestone, slate, argillite



Legend modified and geology compiled for the geochemical map by T.E. Kalnins from maps 1059A, by H.M.A. Rice 1945, 1946, and A.G. Jones 1947, 1951.

Geological cartography by the Geological Survey of Canada

Base-map at the same scale published by the Mapping and Charting Establishment, M.C.E., 1966. Additional drainage obtained from Department of Lands, Forests and Water Resources, British Columbia Land Use maps, 1:125,000 scale.

Mean magnetic declination 1977, 2309.2' East decreasing 4.9' annually. Readings vary from 21049.2' in the SE corner to 23044.2' in the NW corner of the map area.

Elevation in feet above mean sea-level

**Geochemical Symbol and Data Presentation**

The concentration of an element at a sample site is graphically represented as one of 15 symbols, if a sample was collected but no data available a dot is plotted. The symbols are symmetrically arranged so that the first increase in size to the eighth symbol and then increase in blackness to the fifteenth. The two small crosses at the end of the scale are used to respectively denote concentrations below the analytical detection limit, or, in the data group containing the detection limit. The data are grouped on semi-logarithmic scales (e.g. 1, 2, 5, 10, 20, 50, 100 etc. Five decades can be spanned and this arbitrary division has been chosen for the continuing Canada wide series of maps constituting the National Geochemical Reconnaissance.

The choice of symbols and the data groups they represent for any specific element is based on the histogram and cumulative frequency plot for the total survey data. Eight symbols are used for the data group containing one in field season. The eighth symbol is used for the model group as defined by the histogram, this group usually includes the median of the data as defined by the 0.5 (50%) point on the cumulative frequency plot. Some or all, of the remaining 14 symbols are chosen so as to achieve an appropriate graphical impact. An example of all 15 symbols is given below.



The symbol maps, being based on the total survey data distributions, are unaffected by the availability of ever increasing levels of knowledge in bedrock and surficial geology, and other environmental factors. In preference, the raw data symbol maps are only intended to assist the rapid inspection of the data for broad regional features. To fulfill the needs of a more specific and thorough interpretation, the symbol maps should be modified using the field and analytical data provided in the notes listing any other knowledge available. To assist in the appraisal and modification of the data items of the symbol map bedrock geology, table of summary statistics and proposed threshold values for each lithologic unit, or broad lithologic unit, again based on the total survey data, is presented for each geological unit. This table can be used along, or in conjunction, with the sample location map and data listings to indicate above threshold samples where they occur on the map. In many instances, the table will also illustrate, more clearly than the map, the dependence of mean geochemical concentration on type. It may often be observed that whilst the total data appears to approximate a log-normal distribution the data for individual map or lithologic units appears to approximate a normal distribution. The proposed thresholds presented are believed to be useful in interpreting data from a mineral exploration viewpoint. Locations of samples with concentrations in excess of the threshold for the rock unit they appear to be derived from, should be studied carefully. The data may show abnormal concentrations can be due to a wide range of geological and environmental factors, but one of these can be the presence of abnormal concentrations of the element in a form of interest to the mineral explorationist.

To comprehensively study an area all available geological, environmental and recorded data should be utilized. The data separation by bedrock type can often be improved by constructing new data subsets and deriving local threshold levels based on the most detailed and up-to-date knowledge available.

The objective of the survey is to outline broad areas of increased mineral potential worthy of further study leading to the identification of exploration targets. Individual samples with high metal concentrations may not be automatically regarded as finite exploration targets. It is recommended that the data levels vary across the survey area with local geology and surficial environments.

The dispersion of elements in stream sediments is controlled by both mechanical and chemical processes. An insight into the relative importance of these processes can be obtained by a study of host topography, bedrock and surficial geology particularly in terms of host minerals and the chemical properties of each element. The field observations on sediment composition and sample site environment recorded in the data listings can yield information on the relative importance of clastic versus chemical dispersion.

The uranium data for stream sediments were obtained by the total analysis method of neutron activation delayed neutron counting. Therefore, where resistive minerals containing uranium e.g., monzonite, allanite, sphene etc. are present these will lead to acid digestions. As a result care should be taken in interpreting data where streams could derive part or all of their sediment load from crystalline rocks containing such minerals, which at the present have little economic significance as a uranium resource.



NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 6-1976  
URANIUM IN STREAM SEDIMENTS  
CANADA-BRITISH COLUMBIA AGREEMENT ON A URANIUM RECONNAISSANCE PROGRAM

Scale 1:250,000  
Kilometres 6 0 6 12 18 Kilometres  
Miles 4 0 4 8 Miles  
Universal Transverse Mercator Projection  
© Crown Copyrights reserved

92 P 92 M 92 N  
92 L 92 K  
92 H 92 E 92 F  
Map 6-1976  
Open File 410  
Map 5-1976  
Open File 411  
Map 7-1976  
Open File 412




Table of Thresholds for Major Geological Units

Lithology	No. of Samples	Mean	S.D.	C.V.%	Threshold
8 TILL	405	7.1	6.8	96	35
7 OLVB	197	4.4	10.9	249	15
6 COLM	23	5.7	2.7	47	15
6 ANDS	118	7.8	9.7	124	15
5 SYNT	146	17.9	35.7	199	15
4 ANT	966	11.6	17.4	149	50
3 UNFC	3	1.7	0.6	89	60
2a QRTZ	50	5.7	5.0	37	25
2 GRNS	321	3.6	3.4	92	13
1a SCST	241	3.1	1.6	51	15
1 GNSS	1087	9.3	9.5	102	30

Data units are ppm

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 6-1976  
OPEN FILE 410

Resource Geophysics and Geochemistry Division

Geological Survey of Canada, Ottawa

Geochemistry and field operations supervised by S.B. Ballantyne  
Federal-Provincial coordination by E.H.W. Hornbrook  
Analytical chemistry by J.J. Lynch  
Data monitoring by R.G. Garrett, N.C. Lund and D.J. Ellwood

British Columbia, Mineral Resources Branch

Federal-Provincial coordination by A. Sutherland Brown, N.C. Carter and P.A. Christopher  
Field operation assistance by T.E. Kalnins

Contractors:  
Sample collection staff and vehicles supplied by Stokes Exploration Management Co. Ltd.  
Sample preparation by Golder Associates  
Uranium analyses by Atomic Energy of Canada Ltd., Commercial Products Division by delayed neutron activation

This map forms one of a series of 39 sheets released under Geological Survey of Canada, Open Files 409, 410, 411. The Open Files consists of data for 10 elements each for stream sediments, two elements for stream waters and sample site location. The data listing of each Open File includes pH data.

The data are also available in digital form. For further information please contact:

The Director,  
Computer Science Centre,  
Department of Energy, Mines and Resources  
Ottawa, Ontario K1A 0E8

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 6-1976  
OPEN FILE 410  
SOUTH EASTERN BRITISH COLUMBIA, 1976  
URANIUM IN STREAM SEDIMENTS