

Note: This legend is common to National Geochemical Reconnaissance Map 5-1976, Open File 409; Map 6-1976, Open File 410 and Map 7-1976, Open File 411

8 Glacial, lacustrine, and fluvial gravel, sand, silt and clay

7 Plateau basalts, olivine basalts

- 6 Volcanic flow rocks with interbedded sedimentary rocks; 6a, conglomerate, sandstone, shale and tuff

- 5 CORYELL: alkalic plutonic rocks; porphyritic granite and rhyolite

4 NELSON and VALHALLA: granitic plutonic rocks

3 Maffic and ultramafic intrusive rocks, pyroxinite, hornblendite
serpentinite

- PALEOZOIC (including UPPER PROTEROZOIC and TRIASSIC)

- 2 Basaltic and andesitic lavas, greenstone, tuff, quartzite, limestone and argillite; 2a, quartzite, argillite, limestone, slate, schist, phyllite, sandstone and conglomerate

1 Gneiss, minor schist, limestone, marble, dolomite, slate, phyllite;
1a, schist, quartzite, limestone, slate, argillite

Geological context

Geological contact.....
Fault.....
Dike.....

Dyke.....
Mineral occurrence..... $Zn \times$

Legend modified and geology compiled for the geochemical map by T.E. Kalnins from maps 1059A, by H.M.A. Rice 1945, 1946, and A.G. Jones 1947, 1951

Geological cartography by the Geological Survey of Canada

Base-map at the same scale published by the Mapping and Charting Establishment, M.C.E., 1966. Additional drainage obtained from Department of Lands, Forests and Water Resources, British Columbia Land Use maps, 1:125,000 scale

Mean magnetic declination 1977, 23°07.2' East decreasing 4.9' annually.
Readings vary from 21°49.2' in the SE corner to 23°04.2' in the NW corner of the map area

Elevation in feet above mean sea-level

Geochemical Symbol and Data Presentation

The concentration of an element at a sample site is graphically represented as one of 15 symbols, if a sample was collected but there is no data available a dot is plotted. The symbols are symmetrically arranged so that they first increase in size to the eighth symbol and then increase in blackness. The two lowest small crosses at the low end of the scale are used to respectively denote concentrations below the analytical detection limit, or, in the data group containing the detection limit. The data are grouped on a semi-logarithmic scale, i.e., 1,2,5,10,20,50,100 etc. Five decades can be spanned by this arbitrary division has been chosen for the cre-



The choice of symbols and the data groups they represent for any specific element is based on the histogram and cumulative frequency plot for the total survey data from one, or more contiguous, open file sheets covered in one field season. The usually includes the median of the model group as defined by the histogram, this group eighth symbol is used for the data as defined by the 0.5 (50%) point on the group cumulative frequency plot. Some, or all, of the remaining 14 symbols are chosen so as to achieve an appropriate graphical impact. An example of all 15 symbols is given below.

The symbol maps, being based on the total survey data distributions, are unaffected by the availability of ever increasing levels of knowledge in bedrock and surficial geology and environmental factors. Therefore, the raw data symbol maps are only intended to assist in the inspection of the data for gross regional features. To fulfill the needs of a more specific and thorough inspection, the symbol maps should be modified using the field and analytical data provided in the data tables and listings and the sampling and analysis data provided in the data tables of the data in terms of the symbol maps. To assist in the appraisal and modification of the data, the symbol maps for bedrock and surficial geology, and the data tables and proposed threshold values for each mapped bedrock unit, and the data tables and listings for the total survey data, is presented below the histogram. This table can be used along with the symbol maps and the sample location map and data listings to indicate above threshold samples where they occur. The symbol maps, data tables, and the table will also illustrate, more clearly than the map, the dependence of the total data upon the bedrock type. It may often be also observed that whilst the total data appear to be log-normally distributed, the data for individual lithologic units appears to approximate a normal distribution. The data for individual thresholds presented are believed to be useful in interpreting the data from a mineral threshold for the rock unit they appear to be in excess of. The above threshold concentration can be due to a wide range of geological elements, one of these could be the presence of abnormal concentrations of the element in a rock type.

To comprehensively study an area, the following geological, environmental and recorded data should be utilized. The data separation by bedrock type is often best improved by constructing new data subsets and deriving local threshold levels based on the more detailed and up-to-date knowledge available.

The objective of the survey is to outline broad areas of increased mineral potential worthy of further study for identification of exploration targets. Individual data samples with high metal contents should not be automatically regarded as finite areas with local geology and surficial environment.

The dispersion of elements in stream sediments is controlled by both mechanical and chemical processes. An insight into the relative importance of these processes can be gained from a study of local topography, bedrock and surficial geology particularly in terms of host minerals and the chemical properties of each element. The field observations on sediment composition and sample site environment recorded in the data listings can yield information on the relative importance of clastic versus chemical dispersion.



Province of British Columbia
Department of Mines and Petroleum Resources
Mineral Resources Branch

Canada
Department of Energy, Mines and Resources
Geological Survey of Canada






Table of Thresholds for Major Geological Units

Lithology	No. of Samples	Mean	S.D.	C.V. %	Threshold
8 TILL	405	1.6	1.3	80	5
7 OLVB	197	1.5	1.4	90	6
6a CYLM	23	1.3	0.7	55	7
6 ANR	118	2.7	1.7	65	5
5 SALT	146	1.6	2.2	132	5
4 GRNT	966	2.0	4.3	217	7
3 UWS	3	2.7	2.7	88	5
2a ORTZ	50	1.4	0.8	56	7
GRNS	321	2.3	3.6	154	5
1a SCOT	181	1.2	1.8	69	5
1 GNSS	1087	1.7	4.8	274	6

Data units are ppm

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 6-1976
OPEN FILE 410

Resource Geophysics and Geochemistry Division

Geological Survey of Canada, Ottawa

Geochemistry and field operations supervised by S.B. Ballantyne
Federal-Provincial coordination by E.H.W. Hornbrook
Analytical chemistry by J.J. Lynch
Data monitoring by R.G. Garrett, N.G. Lund and D.J. Ellwood

British Columbia, Mineral Resources Branch

Federal-Provincial coordination by A. Sutherland Brown, N.C. Carter
and P.A. Christopher
Field operation assistance by T.E. Kalnins

Contractors
Sample collection staff and vehicles supplied by Stokes
Exploration Management Co. Ltd.
Sample preparation by Golder Associates
Chemical analysis by Chemex Labs Ltd.

This map forms one of a series of 39 sheets released under Geological Survey of Canada, Open Files 409, 410, 411. The Open Files consists of data for 10 elements each for stream sediments, two elements for stream waters and sample site location. The data listing of each Open File includes pH data.

The data are also available in digital form. For further information please contact:

The Director,
Computer Science Centre,
Department of Energy, Mines and Resources
Ottawa, Ontario K1A 0E8

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 6-1976
OPEN FILE 410
SOUTH EASTERN BRITISH COLUMBIA, 1976
MOLYBDENUM IN STREAM SEDIMENTS