

## LEGEND

Note: This legend is common to National Geochemical Reconnaissance Map 5-1976, Open File 409; Map 6-1976, Open File 410 and Map 7-1976, Open File 411

## QUATERNARY

8 Glacial, lacustrine, and fluvial gravel, sand, silt and clay

## TERTIARY

7 Plateau basalts, olivine basalts

6 Volcanic flow rocks with interbedded sedimentary rocks; 6a, conglomerate, sandstone, shale and tuff

5 CORYELL: alkaline plutonic rocks; porphyritic granite and rhyolite

## JURASSIC - CRETACEOUS

4 NELSON and VALHALLA: granitic plutonic rocks

## JURASSIC

3 Maffic and ultramafic intrusive rocks, pyroxinite, hornblende serpentinite

## PALEOZOIC (including UPPER PROTEROZOIC and TRIASSIC)

2 Basaltic and andesitic lavas, greenstone, tuff, quartzite, limestone and argillite; 2a, quartzite, argillite, limestone, slate, schist, phyllite, sandstone and conglomerate

## PROTEROZOIC (SHUSWAP TERRANE)

1 Gneiss, minor schist, limestone, marble, dolomite, slate, phyllite; 1a, schist, quartzite, limestone, slate, argillite

Geological contact.....  
Fault.....  
Dyke.....  
Mineral occurrence.....

Legend modified and geology compiled for the geochemical map by T.E. Kalnins from map 48-1963, by R.B. Campbell and map 12-1964 by J.O. Wheeler

Geological cartography by the Geological Survey of Canada

Base map modified by the Geological Cartography Unit from map published at same scale by Surveys and Mapping Branch, 1965. Additional drainage obtained from Department of Lands, Forests and Water Resources, British Columbia Land Use maps, 1:250,000 scale

Mean magnetic declination 1977, 23°07.2' East decreasing 4.9' annually.  
Readings vary from 22°30' in the SE corner to 23°45' in the NW corner of the map area

Elevation in feet above mean sea-level

## Geochemical Symbol and Data Presentation

The concentration of an element at a sample site is graphically represented as one of 15 symbols, if a sample was collected but there is no data available a dot is plotted. The symbols are symmetrically arranged so that they first increase in size to the eighth symbol and then increase in blackness to the fifteenth. The two small crosses at the low end of the scale are used to respectively denote concentrations below the analytical detection limit, or, in the data group containing the detection limit. The data are grouped on a semi-logarithmic scale, i.e. 1,2,5,10,20,50,100 etc. Five decades can be spanned and this arbitrary division has been chosen for the continuing Canada wide series of maps constituting the National Geochemical Reconnaissance.

The choice of symbols and the data groups they represent for any specific element is based on the histogram and cumulative frequency plot for the total survey data from one, or more contiguous, open file sheets covered in one field season. The eighth symbol is used for the model group as defined by the histogram, this group usually includes the median of the data as defined by the 0.5 (50%) point on the cumulative frequency plot. Some, or all, of the remaining 14 symbols are chosen so as to achieve an appropriate graphical impact. An example of all 15 symbols is given below.

The symbol maps, being based on the total survey data distributions, are unaffected by the availability of ever increasing levels of knowledge in bedrock and surficial geology, and other environmental factors. Therefore, the raw data symbol maps are only intended to assist the rapid inspection of the data for gross regional features. To fulfill the needs of a more specific and thorough interpretation, the raw symbol maps should be modified using the field and analytical data provided in the data listings and any other knowledge available. To assist in the appraisal and modification of the data in terms of the symbol map bedrock geology, a table of summary statistics and proposed threshold values for each mapped bedrock unit, or broad lithologic unit, again based on the total survey data, is presented below the histogram. This table can be used along, or in conjunction, with the sample location map and data listings to indicate above threshold samples where they occur on the map. In many instances, the table will also illustrate, more clearly than the map, the dependence of mean geochemical levels on bedrock type. It may often be also observed that whilst the total data appears to approximate a log-normal distribution the data for individual map or lithologic units appears to approximate a normal distribution. The proposed thresholds presented are believed to be useful in interpreting the data from a mineral exploration viewpoint. Locations of samples with concentrations in excess of the threshold for the rock unit they appear to be derived from, should be studied carefully. The above threshold concentration can be due to a wide range of geological and environmental factors, but one of these could be the presence of abnormal concentrations of the element in a form of interest to the mineral explorationist.


To comprehensively study an area, all available geological, environmental and recorded data should be utilized. The data separation by bedrock type can often be improved by constructing new data subsets and deriving local threshold levels based on the most detailed and up-to-date knowledge available.

The objective of the survey is to outline broad areas of increased mineral potential worthy of further study leading to the identification of exploration targets. Individual samples with high metal contents should not be automatically regarded as finite exploration targets. It is recommended that the data levels vary across the survey area with local geology and surficial environment.

The dispersion of elements in stream sediments is controlled by both mechanical and chemical processes. An insight into the relative importance of these processes can be gained from a study of local topography, bedrock and surficial geology particularly in terms of host minerals and the chemical properties of each element. The field observations on sediment composition and sample site environment recorded in the data listings can yield information on the relative importance of clastic versus chemical dispersion.

Province of British Columbia  
Department of Mines and Petroleum Resources  
Mineral Resources Branch






Canada  
Department of Energy, Mines and Resources  
Geological Survey of Canada



NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 7-1976  
MANGANESE IN STREAM SEDIMENTS  
CANADA-BRITISH COLUMBIA AGREEMENT ON A URANIUM RECONNAISSANCE PROGRAM

Scale 1:250,000

Kilometers 6 0 6 12 18 Kilometers  
Miles 4 0 4 8 Miles  
Universal Transverse Mercator Projection  
© Crown Copyrights reserved



## Table of Thresholds for Major Geological Units

Lithology	No. of Samples	Mean	S.D.	C.V. %	Threshold
8 TILL	405	455	472	104	1500
7 OLVB	197	622	857	138	1800
6a CGLM	23	576	851	148	1600
6 ANDS	118	502	932	186	1500
5 SYNT	146	449	393	87	2000
4 GRNT	966	454	424	93	1500
3 UMFC	3	463	274	59	1500
2a QRTZ	50	412	155	37	1500
2 GRNS	321	522	463	89	2000
1a SCST	241	632	1560	247	2000
1 GNSS	1087	394	398	101	1600

Data units are ppm

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 7-1976  
OPEN FILE 411

Geological Survey of Canada, Ottawa

Geochemistry and field operations supervised by S.B. Ballantyne  
Exploration Management Co. Ltd.  
Analytical chemistry by J.J. Lynch  
Data monitoring by R.G. Garrett, N.G. Lund and D.J. Ellwood

British Columbia, Mineral Resources Branch

Federal-Provincial coordination by A. Sutherland Brown, N.C. Carter  
and P.A. Christopher  
Field operation assistance by T.E. Kalnins

Contractors  
Sample collection staff and vehicles supplied by Stokes  
Exploration Management Co. Ltd.  
Sample preparation by Golder Associates  
Chemical analysis by Chemex Labs Ltd.

This map forms one of a series of 39 sheets released under Geological Survey of Canada, Open Files 409, 410, 411. The Open Files consists of data for 10 elements each for stream sediments, two elements for stream waters and sample site location. The data listing of each Open File includes pH data.

The data are also available in digital form. For further information please contact:

The Director,  
Computer Science Centre,  
Department of Energy, Mines and Resources  
Ottawa, Ontario K1A 0E8

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 7-1976  
OPEN FILE 411  
SOUTH EASTERN BRITISH COLUMBIA, 1976  
MANGANESE IN STREAM SEDIMENTS