LEGEND

Note: This legend is common to National Geochemical Reconnaissance Maps 96-1976, Open File 370, Map 97-1976, Open File 371 and Map 98-1976, Open File 372

QUATERNARY

Q — Unconsolidated glacial and alluvial deposits

PROTEROZOIC

M — Sedimentary, shale and conglomerate; ITA, ADAMS MINE FORMATION: sandstone, shale and conglomerate; ITB, ANNIE MINE FORMATION: conglomerate and sandstone; Ds, ARVID BAY FORMATION: shale and siltstone

Mv — Metamorphosed basaltic greenstone, chlorite schist, commonly porphyritic and amygdaloidal; locally pillowed

Mi — Diorites and gabbros

Mf — Marbled green and maroon shale and brown siltstone

M — HOOK HILL QUARTZITE: massive quartzites, minor slates and phyllite

ARCHEAN

AM — LAKE ELOISE division: argillite, slates, phyllite and minor quartzite, limestone and shale

NEVADIAN

Mn — EAGLE FORMATION: limestone and shaly, sandstone clastics and carbonates

PERMIAN

Pc — CROCKETT FORMATION: grey shale and limestones; Ds, ANNIE CREEK FORMATION: sandstone, shale, carbonates and clastics, undivided

CARBONIFEROUS TO PERMIAN

Pc — CROCKETT FORMATION: shale and limestones; HART RIVER FORMATIONS: shale, siltstone and limestones, Siltarrow Group undivided; Cbs, limestone, black shale, clastic rocks and conglomerate, argillaceous

DEVONIAN

Dd — DUO CREEK FORMATION: shales, unnamed conglomerates

SILURIAN

Sd — SILURO-DEVONIAN TO CARBONIFEROUS

SDG — GOSSAN FORMATION: black shale; BATTEN RIVER FORMATION: shale conglomerate and chert-quartz sandstone; shale, argillite, quartzite, dolomites and minor chert-pebble conglomerate and quartzite

LOWER TO MIDDLE ORDOVICIAN

Ol — OLIVER FORMATION: limestone; CHARKARTA FORMATION: limestone; MITCHELL FORMATION: limestone and shaley unnamed limestone, Siltarrow Group undivided; undivided black shale units

ORDOVICIAN AND SILURIAN

OSE — ROAD RIVER FORMATION: shale, limestone, black chert and argillite; minor quartzite and chert-pebble conglomerate

Ol — Dolomite and limestone; argillaceous limestone and dolomites

CAMBRIAN

A — Limestone and dolomite; minor red phyllic unnamed clastics; Ou, OLDFIELD FORMATION: limestone and dolomite; Ca, CARBONATE FORMATION: quartz, flags and cliffs; Bm, JAMES RIVER: limestone, unnamed dolomite and limestone

PRECAMBRIAN AND/OR CAMBRIAN

Ca — Dark argillite, phyllite, slates, gneiss, tuff, agglomerate, chert, quartzite, limestone and limestone

Ca — Quartzites, conglomerates, quartz-pebble conglomerate, marbles and green phyllite; CRYSTAL RIVER: quartzitic schists, phyllites, limestone and dolomite

ARCHEAN

A — Unnamed carbonates and clastics

A — WHITE GRANITE: muscovite, limestone, fine formation and dolomite

BELTON

Ca — ALPHABEL FORMATION: sandstone and dolomite; TALPEYE FORMATION: sandstone and dolomite, unnamed carbonates, shale and gypsum

Ca — BELTON RIVER GROUP: undivided sandstone, limestone, minor basalt, quartzite, conglomerate, limestone, black shale, argillite and siltstone

BELTON AND ITI UNDIVIDED

S — Argillite, slate, pyritic quartzite, dolomite, conglomerate and siltstone; limestone

Drift-covered areas.....

Geological contact.....

Fault.....

Geology generalized for geochemical map by W.D. Goodfellow from Wind River (1961) and Hart River (1964) by D.K. Norris (G.S.C. Open File 274, 1973), and Eagle River (1963) manuscript unpublished map by D.K. Norris.

Geological cartography by the Geological Survey of Canada

Base-map assembled by the Geological Survey of Canada from maps published at the same scale by the Surveys and Mapping Branch in 1966, 1969

Mean magnetic declination 1977, 30°02'-1° East, decreasing 2.4' annually. Readings vary from 30°26.0' in the SE corner to 29°33.0' in the NW corner of the map-area

Elevations in feet above mean sea-level

GEOCHEMICAL SYMBOLS AND DATA PRESENTATION

The absolute background and anomalous concentrations, and the contrast between them will vary regionally depending on factors such as the physiography, geology, the sample media and the chemistry of the elements determined.

Regional variations in background are well illustrated by the Wernecke, Ogilvie and Richardson Mountains, and the Eagle Plains and Bonnet Plume basin. Most of the sediment in streams intersecting mountain terrain has been derived by the mechanical breakdown of the adjacent rocks. Chemical weathering is relatively unimportant and the heavy spring run-off in these sediments range in size from fine silts to boulders and gravels in a relatively wetter present. By contrast, streams intersecting the Eagle Plains and Bonnet Plume basin are commonly discontinuous, flow at low velocities, and deposit organic-rich sediment. Under these conditions, organic matter may be expected to play a significant role in the dispersion of certain elements (e.g. Cu, Ni, Co, Zn, Cu, Cr, Cu, Ni, Mo, etc.) and may therefore produce spurious anomalies. It is suggested that field observations, such as sediment composition, present in the data will be helpful in the interpretation of the regional variations in background.

The geology of the survey area is represented by sedimentary, volcanic and intrusive rocks that range in age from Lower(?) Proterozoic to Upper Cenozoic, with Paleozoic rocks predominating. Rocks of the Richardson Mountains are chiefly clastics associated with the Proterozoic sedimentary and volcanic rocks of the Wernecke Mountains. Other geological environments that are considered to have a high to moderate potential for the occurrence of stratiform base metal sulphide deposits are Ordovician to Devonian shales and limestones of the Road and Hod Formations in the Richardson Mountains and Eagle Plains, and Cambrian rocks, such as the Eagle Plains and Bonnet Plume Basins and structures such as faults and unconformities which may serve as favorable traps for deposition of mineralization.

Because the chemical derivation of stream sediments from mountainous terrain, the geochemistry is strongly influenced by the chemistry of the underlying rocks. For example shales, or their metamorphic equivalents, which are common in Proterozoic, Paleozoic and Mesozoic rocks, are known to contain anomalous values for a number of trace elements determined which is reflected in the geochemistry of the stream sediments. Therefore, it is suggested that each stream system be evaluated in terms of the local geology before the significance of any one value is established.

The element associations expected for particular types of (and base metal) mineralization should be considered when evaluating geochemically anomalous stream sediments. For example, Ba, Cu, Co and to a lesser extent, Mn, Fe, and V are associated with the Proterozoic(?) occurrences whereas Mo, Y and to a lesser extent, Ni are associated with the Cambrian rocks of the Eagle Plains and Bonnet Plume Basins. Furthermore, element associations in stream sediments and waters will be useful in identifying regional geochemical trends that may be, at least in part, controlled by bedrock lithology. Once these trends have been established, local anomalies can then be evaluated for possible mineralization on the basis of not only the absolute concentrations but also element interrelations.

The distribution of each element in each sample site is graphically represented as one of 15 symbols. If a sample was collected but there is no data available a dot is plotted. The symbols are arranged so that they first increase in size to the eighth class and then increase in density for the remaining seven classes. The smallest symbol (the low end of the size) are used to respectively denote concentrations below the analytical detection limit, or in the data group containing the detection limit. The data are grouped on a semi-logarithmic scale, i.e. 1,2,5,10,20,50,100 etc. Five decades can be spanned and this arbitrary division has been chosen for the contouring of regional data. Other groupings may be more suitable for detailed investigations.

The choice of symbols and the data groups may also represent the statistical element is based on the histogram and cumulative frequency plot for the total survey area. Changes in symbol size normally occur at natural breaks in the frequency. The 4-inch symbol is used for the model group as defined by the histograms. This group usually includes the median of the data as defined by the 0.5 (50%) point on the cumulative frequency plot. Symbols 1 to 7 are used for the remaining 18 symbols to display anomalies of progressively greater geochemical impact.

The raw data symbol maps are only intended to assist the rapid inspection of the data for gross regional features. To fulfil the need for a more specific and thorough interpretation, the field and analytical data provided in the data listings should be considered. For the purpose of comparing the data with the available bedrock geology, a table of summary statistics for the drainage samples underlain by each bedrock unit, or broad lithologic unit, is presented in the data listings. Statistics are presented for each unit in the order of abundance on the map. The dependence of mean geochemical levels on bedrock type. It may also be observed that within the total data appears to approximate a log-normal distribution. The mean and standard deviation of the logarithms of the data approximate a normal distribution except where the concentration of an element at or below the detection limit for a large number of samples (e.g. U and F in waters) Ag, Mo and U in sediments). In such cases the mean and standard deviation of the logarithms are skewed and the mean will not represent the total population. Therefore, statistics must be corrected when using the table of summary statistics to establish background and anomalous concentration ranges for a given element.

To comprehensively study an area, all available geological, environmental and geochemical information must be used. The most significant anomalies can then be determined by constructing raw-data subsets and deriving local threshold levels based on the most detailed and up-to-date knowledge available.

EXAMPLE

Canada
Department of Energy, Mines and Resources
Geological Survey of Canada

Parts of 106 E, 5 and 116 B, 1