

Geochemical Symbol and Data Presentation



The concentration of an element at a sample site is graphically represented as one of 15 symbols. If a sample was collected but there is no data available a dot is plotted. The symbol and its graphical arrangement so that the first increase in size to the eighth symbol and then increase in blackness to the fifteenth. The two small crosses at the low end of the scale are used to respectively denote concentrations below the analytical detection limit, or, in the data group containing the detection limit. The data are recorded on a semi-logarithmic scale, i.e., 1, 2, 5, 10, 20, 50, 100, etc. decades can be spaced and this arbitrary division has been chosen for the continuing Canada wide series of maps constituting the National Geochemical Reconnaissance.

The choice of symbols and the data groups they represent for any specific element is based on the histogram and cumulative frequency plot for the total survey data from one, or more contiguous, open file sheets covered in one field season (above). The eighth symbol is used for the model group as defined by the histogram. This group usually includes the median of the data as defined by the 50% point of the distribution. Some or all of the remaining 14 symbols are chosen so as to achieve an appropriate graphical impact. An example of all 15 symbols is given below.

The symbol maps, being based on the total survey data distributions, are unaffected by the availability of ever increasing levels of knowledge in bedrock and surficial geology, and other environmental factors. Therefore, the raw data symbol maps are only intended to assist the rapid inspection of the data for gross regional features. To fulfill the needs of a more specific and thorough interpretation of the symbol maps, they should be modified using the field and analytical data provided in the data listings and any other knowledge available.

The data listings contain notes on survey and analytical methods, raw data listing with legend and statistics for total data as well as for data grouped on the basis of rock type.

To comprehensively study an area, all available geological, environmental and recorded data should be utilized. The data compilation of bedrock types can often be carried out using the data subsets and deriving local threshold levels based on the most detailed and up-to-date knowledge available.




Copies of map material and listings of field observations and analytical data from which the material was prepared may be available at users expense by application to:

K.G. Campbell Corporation
880 Wellington Street
Bay 238
Ottawa, Ontario
K1R 6X7

The data is also available in digital form. For further information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4



Elevations in feet above mean sea level


Mean magnetic declination 1978, 16° 11'.4" East,
decreasing 6.3' annually. Readings vary
from 16920.0' in the SE corner to
20911.4' in the NW corner of
the map-area

IRON (%) OPEN FILE 508

NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 19-1977 URANIUM RECONNAISSANCE PROGRAM

Scale 1:250,000
Kilometers 6 9 6 12 18 Kilometers
Miles 4 6 4 5 Miles
Universal Transverse Mercator Projection
© Crown Copyright reserved

Base-map assembled by the Geological Cartography Unit from maps published at the same scale by the Surveys and Mapping Branch in 1962, 1963



This map forms one of a series of 14 sheets released under the Geological Survey of Canada, Open File 508. The Open File consists of maps of 11 elements, each for lake sediments, 2 elements for lake waters and sample site location.

IRON (%)
OPEN FILE 508
NORTHEASTERN SASKATCHEWAN, 1977

IRON (%)
OPEN FILE 508
NORTHEASTERN SASKATCHEWAN, 1977

IRON (%)
OPEN FILE 508
NORTHEASTERN SASKATCHEWAN, 1977

64 L 64 M

This map has been reproduced from a scanned version of the original map.
Reproduction par numérisation d'une carte sur papier

64 L 64 M