

Geological
Survey
of Canada



Commission
Géologique
du Canada


Energy, Mines and Resources
Énergie, Mines et Ressources

SURFICIAL GEOLOGY AND GEOMORPHOLOGY

SOMERSET ISLAND

MAP 3

by J. A. Netterville, A. S. Dyke, R. D. Thomas
1976



UNCONTROLLED PHOTOMOSAIC BASE

EXPLANATION OF MAP SYMBOLS

DOMINANT TEXTURE

- r - rock rubble
- g - gravel
- s - sand
- f - silt and fine sand
- t - clay to fine sand
- l - till

BEDROCK TYPE

- g - granitic
- c - carbonate
- s - sandstone
- sh - shale
- cgl - conglomerate
- ES - Eureka Sound Formation

ORIGIN

- M - moraine
- I - ice-contact outwash
- P - preglacial outwash
- Re - marine
- D - deltaic
- A - alluvial (active)
- B - bedrock
- W - weathered product

MORPHOLOGY

- p - plain
- m - rolling
- h - hummocky
- r - ridged
- t - terraced
- k - kettled
- f - fan

MORPHOLOGIC MODIFIER

- D - dissected
- W - washed

RELIEF CLASS

- 1 - less than 5 metres
- 2 - 5 to 20 metres
- 3 - 20 to 50 metres
- 4 - greater than 50 metres

SLOPE CLASS

- 1 - less than 5 degrees
- 2 - 5 to 15 degrees
- 3 - 15 to 35 degrees
- 4 - greater than 35 degrees

- Glacial striae (ice direction known, not known)
- Drumlin, drumlinoid, fluting (ice direction indicated, not indicated)
- Crag - and - tail (ice movement in direction of arrow)
- Moraine ridge
- Esker (direction of flow assumed, uncertain)
- Meltwater channel (large, small)
- Abandoned beach ridge
- Escarpment
- Retrospective thaw-flow slide
- Ice wedge polygons (areas known to contain ice wedges)
- Rock glacier

COMPOSITE UNITS

- / first of units covers more than 50% of total unit area
- first of units covers 60 - 80% of total unit area
- units are of roughly equal proportions

Notes: units which comprise less than 5% of the total area of another unit are not mapped


STRATIGRAPHY

Vener units (less than 1.5 metres thick) are indicated by the lower case letter "v" after the generic symbol. A horizontal line separates the vener unit symbols from the symbols representing the underlying unit.

EXAMPLE

origin stratigraphy
texture - morphology modifier
bedrock type - slope class
relief class
morphology

Describes an area which consists of gently rolling carbonate and shale bedrock with macroscopic relief of 20 - 50 m and slopes commonly less than 5°. Superimposed on this gently rolling landscape occur lower bedrock hills (relief 5-20 m) with steeper slopes of 5 - 15°. About 60 - 80% of the area is covered by a veneer (less than 1.5 m) of silt; the remainder (20 - 40% of the area) is covered by a veneer of silty rubble produced by weathering of the bedrock. The entire area was once below water. This has to some extent modified the original materials through sorting and/or redeposition.



OPEN FILE
DOSSIER PUBLIC
357
AUG 1976
GEOLOGICAL SURVEY
COMMISSION GÉOLOGIQUE
OTTAWA

This document was produced
by scanning the original publication.
Ce document est le produit d'une
numérisation par balayage
de la publication originale.

Refer to extended legend for additional information