LEGEND

DEVONIAN OR YOUNGER

7 Pink to buff, massive, very fine grained to fine grained, quartz-feldspar porphyritic dikes.

SILURIAN TO DEVONIAN

Burgeo Batholith (6a-6b)

- 6b Pink, protomylonitic to mylonitic, K-feldspar porphyroclastic granite.
- 6a Pink, very strongly foliated to protomylonitic, medium grained, equigranular, biotite granite with screens of migmatite (Unit 3).

North Bay Granite (subunits 5a-5i)

- 5i Pink to buff, masive to weakly cleaved, medium grained, K-feldspar porphyritic, biotite-muscovite granite.
- 5h Pink to buff, massive to weakly cleaved, medium to coarse-medium grained, K-feldspar porphyritic, biotite ± muscovite granite.
- 5g Pink to buff, massive to weakly cleaved, medium to coarse-medium grained, K-feldspar porphyritic, biotite granite; screens of foliated granite and migmatite common southwest of Dolland Brook.
- 5f White to buff, massive, coarse grained, equigranular, biotite-muscovite
- 5e Buff, massive, medium grained, equigranular, biotite granite; includes minor biotite-muscovite granite.
- 5d Buff, massive, medium grained, equigranular, biotite granodiorite and granite; granite becomes fine grained and muscovite-bearing southwest of Meelpaeg Lake.
- 5c Buff to gray, foliated, medium to coarse-medium grained, equigranular, biotite granodiorite and minor tonalite; locally intruded by garnet-muscovite aplite and pegmatite veins.
- 5b Pink to white, strongly foliated, medium to coarse grained, equigranular to pegmatitic, muscovite-garnet and muscovite-biotite granite; locally contains numerous xenoliths of migmatite, psammite, and foliated tonalite.
- 5a Pink to buff, foliated, medium grained, equigranular to porphyritic, biotite ± muscovite granodiorite and granite.

SILURIAN?

4 Interbedded sandstone, siltstone and pebble conglomerate; minor thickly bedded subangular cobble conglomerate.

LOWER TO MIDDLE ORDOVICIAN

Bay du Nord Group

3 Highly deformed, migmatitic metasediments cut by numerous granitoid veins, dikes and small granitoid plugs.

Baie d'Espoir Group (subunits 2a-2e)

- 2e Thinly bedded, locally highly folded, gray slate and sandstone; possibly equivalent to Salmon River Dam Formation.
- 2d Gray and green, thinly bedded, slate and minor sandstone.
- 2c Very thickly bedded, brown weathering, gray massive sandstone, minor interbeds of subunit 2b.
- 2b Thinly bedded, highly folded and quartz veined, black graphitic slate and siltstone.
- 2a Thickly bedded, schistose, quartz-crystal tuff and coarse tuffaceous

Spruce Brook Formation (subunits la-ld)

- ld Migmatite, psammite, and numerous granitoid dikes; high grade equivalent of la, lb.
- 1c Clast-supported orthoquartzite cobble metaconglomerate, with minor matrix-supported slate and quartzite breccia and coarse metasandstone.
- 1b Thinly bedded pelite and minor psammite and semipelite.
- la Dominantly biotite psammite and orthoquartzite with minor semipelite.

NOTE Legend is common to maps 12A/2E and 11P/15E; thus some units may not appear on both maps.

SYMBOLS

Geology by W.L. Dickson and P.W. Delaney, 1983. Released April, 1984.

Contribution to the Canada-Newfoundland Co-operative Minerals Program 1982-1984. Project funded by Geological Survey of Canada and the Newfoundland Department of Mines and Energy.

This preliminary map is subject to revision.

Base map at same scale published by Surveys and Mapping Branch, Department of Energy, Mines and Resources, Ottawa, 1973.

Approximate M-gnetic Declination 1984 for centre of map 12A/2 is 26°22'W, decreasing 3.3' annually.

Copies of this map may be obtained from the Publications and Information Section, Mineral Development Division, Department of Mines and Energy, P.O. Box 4750, St. John's, Newfoundland AlC 5T7, and from the Geological Survey of Canada, 601 Booth Street, Ottawa, Ontario KlA 0E8, and from K.G. Campbell Corporation, 880 Wellington St., Bay 238, Ottawa, Ontario KlR 6K7

G.S.C. Open File # 1028 Wolf Mountain.

