

Geological Survey of Canada
Resource Geophysics and Geochemistry Division
and
Manitoba Department of Energy and Mines
Mineral Resources Division

CONTRACTORS

Sample collection by Marshall Macklin Monaghan Ltd., Toronto
Sample preparation by Golder Associates, Ottawa



Sediment chemical analysis by Barringer Magenta Ltd., Rexdale, Ontario
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
880 Wellington St.
Bay 238
Ottawa, Ontario
K1R 6K7

The data are also available in digital form. For further information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4



Elevation in feet above mean sea level

Mean magnetic declination 1985, 9°08' East, decreasing 21.3' annually. Readings vary from 7°44' in the NE corner to 10°25' in the SW corner of the map area

MERCURY(ppb)
GSC OPEN FILE 1103
REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 68-1984
CANADA - MANITOBA
MINERAL DEVELOPMENT AGREEMENT (1984-89)
LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY
NORTH-WEST MANITOBA, 1984

Base map at the same scale published by the Surveys and Mapping Branch in 1963

LEGEND

Note: This legend is common for Regional Geochemical Reconnaissance Map 68-1984, Open File 1103.

- PROTEROZOIC (APHEBIAN)
- 31 GRANITIC INTRUSIVE ROCKS, POST-SICKLE (HUDSONIAN) (AH1a to AH1f)*
31a-granite (AH1a); 31b-granodiorite, tonalite; 31c-megacrystic granite; 31c-granite, granodiorite ± muscovite; 31d-leucogranite, tonalite; 31e-monzonite, syenite; 31f-pegmatite
 - 30 GRANITIC INTRUSIVE ROCKS, POST-SICKLE and remobilized PRE-SICKLE
30-granite, granodiorite (AH1g)
 - 29 INTERMEDIATE INTRUSIVE ROCKS, POST-SICKLE and remobilized PRE-SICKLE
29-tonalite, granodiorite, quartz diorite (AH1h); 29a-pyroxene tonalite (AH1p)
 - 28 MAFIC INTRUSIVE ROCKS, POST-SICKLE
28-gabbro, minor ultramafic rock (AH1r)
 - 27 BLACK TROUT INTRUSIVE SUITE
27-quartz diorite, diorite (AT1q)
- | SICKLE GROUP | SICKLE METAMORPHIC SUITE | SOUTHERN INDIAN GNEISS |
|--|---|---|
| 26 ARKOSIC METASEDIMENTARY ROCKS, DERIVED GNEISS
26a-conglomerate (ASac)
26b-arkosic sandstone (ASas) | 26c-sandstone-derived gneiss, migmatite (ASan) | 26d-felsic, minor mafic gneiss (age unknown) (AImf) |
| PRE-SICKLE INTRUSIVE ROCKS
25a-gabbro, norite, ultramafic rock (AP1r)
25b-tonalite, granodiorite, diorite (AP1t)
25c-granite (AP1g); 25d-gabbro-quartz diorite (AP1d) | | |
| 24 WASEKWAN OR SICKLE GROUP
24a-AMPHIBOLITE, CALC-SILICATE ROCK, METASEDIMENTARY ROCKS
24a-conglomerate, greywacke (AGmc); 24b-felsic gneiss (AGmf) | 24c-AMPHIBOLITE, CALC-SILICATE ROCK, METASEDIMENTARY ROCKS
24c-greyswacke (AGsm)
24c-conglomerate, greywacke (AGmc); 24d-felsic gneiss (AGmf) | 24d-amphibolite, tuff (A11ma)
24e-greyswacke-derived gneiss, migmatite (A11sw) |
| 23 WASEKWAN GROUP
23a-METASEDIMENTARY ROCKS
23a-greyswacke, conglomerate, mafic mudstone (AWsw) | 23b-METASEDIMENTARY ROCKS
23b-greyswacke-derived gneiss, migmatite (ABsw) | 23c-greyswacke-derived gneiss and migmatite (A11sw) |
| 22 FELSIC, INTERMEDIATE VOLCANICS
22a-dacite, rhyolite (AW1d) | 22b-METASEDIMENTARY ROCKS
22b-greyswacke, quartzite, marble (ABmn)
22b-greyswacke-derived gneiss, migmatite (ABmn) | |
| 21 MAFIC, INTERMEDIATE VOLCANICS
21a-basalt andesite (AW1va)
21b-basalt (AW1vb) | | |

Geological boundary (approximate, inferred).....
Fault approximate or inferred.....
Area of no outcrop.....
No analytical result.....

* A four character mnemonic name recorded rock type as part of the 1984 field observations

Provisional Compilation map by H.W. Zwanzig,
Manitoba Department of Energy and Mines

This map forms one of a series of maps released by the Geological Survey of Canada, Open File 1103 to 1105. Each Open File consists of maps of various geochemical variables: 16 for lake sediment, 3 for lake water and 1 sample site location