

## LEGEND

Note: This legend is common for Regional Geochemical Reconnaissance Map 69-1984, Open File 1104.

- A\* Metadiorite, hornblendite of possible Archean age
- 1 Amphibolite, volcanic derived with locally preserved pillows
- 2a Biotite-feldspar-quartz-paragneiss + garnet + granite ± muscovite
- 2b Biotite metatexite + garnet + granite (25-75% white granitic lit)
- 2c Biotite metatexite + garnet + cordierite
- 3a Light grey biotite (5-10%) quartz-feldspar-gneiss + magnetite + garnet with discontinuous diorite gneiss lenses
- 3b Light grey to dark-grey biotite (5-15%) quartz-feldspar-gneiss interlayered with thin layers of amphibolite and/or hornblende-biotite bearing layers
- 4 Calc-silicate rock
- 5 Amphibolite, metagabbro, locally agmatitic
- 6a Metaconglomerate
- 6b Thin interlayered amphibolite and hornblende biotite-bearing layers
- 6c Arkosic gneiss
- 6d Metavolcanic rocks
- 6e Metagreywacke
- 7 Gneissic diorite and leucodiorite
- 7a Biotite ± hornblende granodiorite gneiss with white granitic lit
- 7b Gabbro
- 8 Grey, medium to coarse grained biotite (5%) + magnetite-tonalite to quartz monzonite
- 8a Hybrid gneiss of grey biotite-quartz monzonite and gneissic diorite
- 9 Foliated quartz diorite + magnetite
- 10 Biotite (15-20%) - tonalite ± garnet
- 11a Megacrystic biotite-granodiorite
- 11b Megacrystic biotite-hornblende ± pyroxene-granodiorite
- 11c Coarse grained leucocratic granodiorite
- 12 White leucocratic medium grained to pegmatitic monzogranite ± garnet
- 13 Coarse grained to megacrystic-pyroxene-hornblende-monzonite to monzogranite with olive-brown feldspar
- 13a Anorthositic gabbro
- 13b Hornblende-biotite-monzonite to quartz monzonite with variegated olive-brown and pink feldspar
- 14 Megacrystic-biotite-magnetite quartz monzonite
- 15 Biotite ± hornblende coarse grained to megacrystic pink granite to quartz monzonite
- 15a Biotite-hornblende granite gneiss
- 15b Leucocratic megacrystic pink granite
- 15c Fine grained quartz monzonite
- 16 Magnetite-biotite-hornblende quartz monzonite
- 17 Granite pegmatite
- 18 Diabase

Pyrite, chalcopyrite, galena, sphalerite,  
Iron Formation ..... Py; ▲ Cp; ▲ Pb; ▲ Sp; ▲ I.F.  
Geological boundary (approximate, assumed, gradational) ....  
Drift covered .....


Provisional Compilation map by H.W. Zwanzig,  
Manitoba Department of Energy and Mines

\* A four character mnemonic name recorded rock type as part of the 1984 field observations

This map forms one of a series of maps released by the Geological Survey of Canada, Open File 1103 to 1105. Each Open File consists of maps of various geochemical variables: 16 for lake sediment, 3 for lake water and 1 sample site location

PB

PBM X TILE  
14 MAX  
6 95  
4 90  
3 80  
1 MIN  
965 SAMPLES



Geological Survey of Canada  
Resource Geophysics and Geochemistry Division

Manitoba Department of Energy and Mines  
Mineral Resources Division

CONTRACTORS

Sample collection by Marshall Macklin Monaghan Ltd., Toronto  
Sample preparation by Golder Associates, Ottawa




Sediment chemical analysis by Barringer Magenta Ltd., Rexdale, Ontario  
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation  
880 Wellington St.  
Bay 238  
Ottawa, Ontario  
K1A 6K7

The data are also available in digital form. For further information please contact:

The Director  
Computer Science Centre  
Department of Energy, Mines and Resources  
Ottawa, Ontario  
K1A 0E4



PROGLACIAL AND GLACIAL ENVIRONMENT

- GLACIOLACUSTRIAL DEPOSITS: beach and nearshore deposits: sand and gravel 1-4 m thick, forming distinct ridges
- GLACIOLACUSTRIAL DEPOSITS: deep basin deposits: silt, clay and sand, 1-30 m thick
- GLACIOFLUVIAL DEPOSITS: gravel, sand and silt, 1-100 m thick

GLACIAL ENVIRONMENT

- GLACIAL DEPOSITS: till 1-5 m thick, derived primarily from Precambrian bedrock


NONGLACIAL ENVIRONMENT

- BEDROCK
- ORGANIC DEPOSITS: marsh, fen, swamp and bog deposits up to 6 m thick, characterized by seasonal flooding

Striations .....

Flutings, drumlins, and drumlinoid ridges, oriented parallel to ice flow direction .....


Esker (flow direction known or inferred) .....



Elevation in feet above mean sea level  
GSC OPEN FILE 1104

REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 69-1984  
CANADA - MANITOBA  
MINERAL DEVELOPMENT AGREEMENT (1984-89)  
LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY  
NORTH-WEST MANITOBA, 1984

Scale 1:250 000  
Kilometres 5 0 5 10 15 20 Kilometres  
Universal Transverse Mercator Projection  
© Crown Copyrights reserved



This map has been reprinted from a scanned version of the original map.  
Reproduction par numérisation d'une carte sur papier