

- LEGEND
- Note: This legend is common for Regional Geochemical Reconnaissance Map 70-1984, Open File 1105
- A\* Metadiorite, hornblende of possible Archean age
  - 1 Amphibolite, volcanic derived with locally preserved pillows
  - 2a Biotite-feldspar-quartz-paragneiss + garnet + granite ± muscovite
  - 2b Biotite metatextite + garnet + granite (25-75% white granitic lit)
  - 2c Biotite metatextite + garnet + cordierite
  - 3a Light grey biotite (5-10%) quartz-feldspar-gneiss + magnetite + garnet with discontinuous diorite gneiss lenses
  - 3b Light grey to dark grey biotite (5-15%) quartz-feldspar-gneiss interlayered with thin layers of amphibolite and/or hornblende-biotite bearing layers
  - 4 Calc-silicate rock
  - 5 Amphibolite, metagabbro, locally agmatitic
  - 6a Metacglomerate
  - 6b Thin interlayered amphibolite and hornblende biotite-bearing layers
  - 6c Arkosic gneiss
  - 6d Metavolcanic rocks
  - 6e Metagreywacke
  - 7 Gneissic diorite and leucodiorite
  - 7a Biotite ± hornblende granodiorite gneiss with white granitic lit
  - 7b Gabbro
  - 8 Grey, medium to coarse grained biotite (5%) + magnetite-tonalite to quartz monzonite
  - 8a Hybrid gneiss of grey biotite-quartz monzonite and gneissic diorite
  - 9 Foliated quartz diorite + magnetite
  - 10 Biotite (15-20%) - tonalite ± garnet
  - 11a Megacrystic biotite-granodiorite
  - 11b Megacrystic biotite-hornblende ± pyroxene-granodiorite
  - 11c Coarse grained leucocratic granodiorite
  - 12 White leucocratic medium grained to pegmatitic monzogranite ± garnet
  - 13 Coarse grained to megacrystic-pyroxene-hornblende-monzonite to monzogranite with olive-brown feldspar
  - 13a Anorthostic gabbro
  - 13b Hornblende-biotite-monzonite to quartz monzonite with variegated olive-brown and pink feldspar
  - 14 Megacrystic-biotite-magnetite quartz monzonite
  - 15 Biotite ± hornblende coarse grained to megacrystic pink granite to quartz monzonite
  - 15a Hornblende-granite gneiss
  - 15b Leucocratic megacrystic pink granite
  - 15c Fine grained quartz monzonite
  - 16 Magnetite-biotite-hornblende quartz monzonite
  - 17 Granite pegmatite
  - 18 Diabase
- Pyrite, chalcopyrite, galena, sphalerite, Iron Formation  
 Geological boundary (approximate, assumed, gradational)  
 Drift covered
- \* A four character mnemonic name recorded rock type as part of the 1984 field observations

Geological Survey of Canada  
 Resource Geophysics and Geochemistry Division  
 and  
 Manitoba Department of Energy and Mines  
 Mineral Resources Division

CONTRACTORS

Sample collection by Marshall Macklin Monaghan Ltd., Toronto  
 Sample preparation by Golder Associates, Ottawa

Sediment chemical analysis by Barringer Magenta Ltd., Rexdale, Ontario  
 Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation  
 880 Wellington St.  
 Bay 238  
 Ottawa, Ontario  
 K1R 6K7

The data are also available in digital form. For further information please contact:

The Director  
 Computer Science Centre  
 Department of Energy, Mines and Resources  
 Ottawa, Ontario  
 K1A 0E4

PROGLACIAL AND GLACIAL ENVIRONMENT

- GLACIOLACUSTRINE DEPOSITS: beach and nearshore deposits: sand and gravel 1-4 m thick, forming distinct ridges
- GLACIOLACUSTRINE DEPOSITS: deep basin deposits: silt, clay and sand, 1-30 m thick
- GLACIOFLUVIAL DEPOSITS: gravel, sand and silt, 1-100 m thick


GLACIAL ENVIRONMENT

- GLACIAL DEPOSITS: till: 1-5 m thick, derived primarily from Precambrian bedrock

NONGLACIAL ENVIRONMENT

- BEDROCK
- ORGANIC DEPOSITS: marsh, fen, swamp and bog deposits up to 6 m thick, characterized by seasonal flooding

Striations  
 Flutings, drumlins, and drumlinoid ridges, oriented parallel to ice flow direction  
 Esker (flow direction known or inferred)



Elevation in feet above mean sea level


Mean magnetic declination 1985, 9°06' East, decreasing 23.1' annually. Readings vary from 7°35' in the NE corner to 10°20' in the SW corner of the map area

URANIUM (ppm)  
 GSC OPEN FILE 1105  
 REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 70-1984  
 CANADA - MANITOBA  
 MINERAL DEVELOPMENT AGREEMENT (1984-89)  
 LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY  
 NORTH-WEST MANITOBA, 1984

Scale 1:250 000

Universal Transverse Mercator Projection  
 © Crown Copyrights reserved

Base map at the same scale published by the Surveys and Mapping Branch in 1963



This map forms one of a series of maps released by the Geological Survey of Canada, Open File 1103 to 1105. Each Open File consists of maps of various geochemical variables: 16 for lake sediment, 3 for lake water and 1 sample site location

This map has been reprinted from a scanned version of the original map. Reproduction per numérisation d'une carte sur papier.