




	GLACIOLACUSTRINE DEPOSITS: beach and nearshore deposits: sand and gravel 1-4 m thick, forming distinct ridges
	GLACIOLACUSTRINE DEPOSITS: deep basin deposits: silt, clay and sand, 1-30 m thick
	GLACIOFLUVIAL DEPOSITS: gravel, sand and silt, 1-100 m thick


 **GLACIAL DEPOSITS:** till: 1-5 m thick, derived primarily from Precambrian bedrock


 BEDROCK

 ORGANIC DEPOSITS: marsh, fen, swamp and bog deposits up to 6 m thick, characterized by seasonal flooding

Striations 

Flutings, drumlins, and drumlinoid ridges, oriented
parallel to ice flow direction 


Esker (flow direction known or inferred) 



Mean magnetic declination 1985, 9°06' East,
decreasing 23.1' annually. Readings vary from
7°35' in the NE corner to 10°28' in the SW
corner of the map area


Manitoba
Department of Energy and Mines

Canada
Department of Energy, Mines and Resources
Geological Survey of Canada



GSC OPEN FILE 1105
REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 70-1984
CANADA - MANITOBA
MINERAL DEVELOPMENT AGREEMENT (1984-89)
LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY
NORTH-WEST MANITOBA, 1984

Base map at the same scale published by
the Surveys and Mapping Branch in 1963



This map has been reprinted from a scanned version of the original map
Reproduction par numérisation d'une carte sur papier

URANIUM in water (ppb)
GSC OPEN FILE 1105
NORTH-WEST MANITOBA, 1984

LEGEND

Note: This legend is common for Regional Geochemical Reconnaissance Map 70-1984, Open File 1105

- | | |
|-----|--|
| A* | Metadiorite, hornblendeite of possible Archean age |
| 1 | Amphibolite, volcanic derived with locally preserved pillows |
| 2a | Biotite-feldspar-quartz-paragneiss + garnet + granite \pm muscovite |
| 2b | Biotite metatextite + garnet + granite (25-75% white granitic lit) |
| 2c | Biotite metatextite + garnet + cordierite |
| 3a | Light grey biotite (5-10%) quartz-feldspar-gneiss + magnetite + garnet with discontinuous diorite gneiss lenses |
| 3b | Light grey to dark grey biotite (5-15%) quartz-feldspar-gneiss interlayered with thin layers of amphibolite and/or hornblende-biotite bearing layers |
| 4 | Calc-silicate rock |
| 5 | Amphibolite, metagabbro, locally agmatitic |
| 6a | Metaconglomerate |
| 6b | Thin interlayered amphibolite and hornblende biotite-bearing layers |
| 6c | Arkosic gneiss |
| 6d | Metavolcanic rocks |
| 6e | Metagreywacke |
| 7 | Gneissic diorite and leucodiorite |
| 7a | Biotite \pm hornblende granodiorite gneiss with white granitic lit |
| 7b | Gabbro |
| 8 | Grey, medium to coarse grained biotite (5%) + magnetite-tonalite to quartz monzonite |
| 8a | Hybrid gneiss of grey biotite-quartz monzonite and gneissic diorite |
| 9 | Foliated quartz diorite + magnetite |
| 10 | Biotite (15-20%) - tonalite \pm garnet |
| 11a | Megacrystic biotite-granodiorite |
| 11b | Megacrystic biotite-hornblende \pm pyroxene-granodiorite |
| 11c | Coarse grained leucocratic granodiorite |
| 12 | White leucocratic medium grained to pegmatitic monzogranite \pm garnet |
| 13 | Coarse grained to megacrystic-pyroxene-hornblende-monzonite to monzogranite with olive-brown feldspar |
| 13a | Anorthositic gabbro |
| 13b | Hornblende-biotite-monzonite to quartz monzonite with variegated olive-brown and pink feldspar |
| 14 | Megacrystic-biotite-magnetite quartz monzonite |
| 15 | Biotite \pm hornblende coarse grained to megacrystic pink granite to quartz monzonite |
| 15a | Biotite-hornblende granite gneiss |
| 15b | Leucocratic megacrystic pink granite |
| 15c | Fine grained quartz monzonite |
| 16 | Magnetite-biotite-hornblende quartz monzonite |
| 17 | Granite pegmatite |
| 18 | Dabase |

Pyrite, chalcopyrite, galena, sphalerite,
Iron Formation ▲Py: ▲Cp: ▲Pb: ▲Sp: ▲I.F.
Geological boundary (approximate, assumed, gradational)
Drift covered

* A four character mnemonic name recorded rock type as part of the 1984 field observations

Provisional Compilation map by H.V. Zwanzig,
Manitoba Department of Energy and Mines

This map forms one of a series of maps released by the Geological Survey of Canada, Open File 1103 to 1105. Each Open File consists of maps of various geochemical variables: 16 for lake sediment, 3 for lake water and 1 sample site location

URANIUM in water (ppb)
GSC OPEN FILE 1105
NORTH-WEST MANITOBA, 1984