Canada
Department of Energy, Mines and Resources
Geological Survey of Canada

Province of Saskatchewan
Saskatchewan Energy and Mines
Saskatchewan Geological Survey

64 E, and parts of 74 A, H





Complexes: where two or more classes of terrain are interspersed in a mosaic or repeating pattern the proportion of each component in the combination is given in a three-position designation set off by slashes denoting arbitrary percentage limits. For example, "Mv/0/R" means that at least 60% of the area is underlain by thin till, with up to 40% boggy areas, and less than 15% scattered rock outcrops. "Rc/R" indicates more than 60% bedrock concealed by vegetation and less than 15% outcrop.

SAMPLE LOCATION
GSC OPEN FILE 1106
REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 71-1984
CANADA - SASKATCHEWAN
MINERAL DEVELOPMENT AGREEMENT (1984-89)
LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY
NORTHEASTERN SASKATCHEWAN, 1984

Scale 1:250 000
Kilometres 5 0 5 10 15 20 Kilometres
Universal Transverse Mercator Projection
© Crown Copyrights reserved

Mean magnetic declination 1985, 15°06' East,
decreasing 23.1' annually. Readings
vary from 13°05' in the SE corner to 17°05' in the NW
corner of the map area

Geological Survey of Canada
Resource Geophysics and Geochemistry Division
and
Saskatchewan Energy and Mines
Saskatchewan Geological Survey

CONTRACTORS

Sample collection by Marshall Macklin Monaghan Ltd., Toronto
Sample preparation by Golder Associates, Ottawa

Sediment chemical analysis by Barringer Magenta ltd., Rexdale, Ontario
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
880 Wellington St.
Bay 238
Ottawa, Ontario
K1R 6K7

The data are also available in digital form. For further information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4

SAMPLE LOCATION
GSC OPEN FILE 1106
NORTHEASTERN SASKATCHEWAN, 1984

LEGEND

Note: This legend is common for Regional Geochemical Reconnaissance Map 71-1984, Open File 1106

NEOHELIKIAN-HADRYNIAN

dd — **Diabase-gabbro**: fine to coarse grained, massive to weakly foliated ± olivine ± biotite ± hypersthene

PALEOHELIKIAN

ATHABASCA GROUP

MF — **Manitou Falls Formation**: sandstone and conglomerate
MFc: fluvial sandstone member, locally pebbly
MFb:conglomerate member

LATE APHEBIAN (HUDSONIAN)

X — **Cataclasite**: mylonite and sheared rocks of the Needle Falls Shear Zone; derived from rocks of the Wollaston and Peter Lake Domains

WOLLASTON DOMAIN

LATE APHEBIAN (HUDSONIAN)

Wpeg — **Granitic pegmatite**: variable grain size, generally massive ± biotite ± muscovite ± garnet; contacts with migmatitic supracrustal gneisses commonly gradational

Wg — **Granite and aplasite**: fine to coarse grained, massive to weakly foliated ± biotite ± hornblende ± magnetite ± microcline megacrysts; contacts with migmatitic supracrustal gneisses commonly gradational

Wgdb — **Biotite granodiorite-tonalite**: medium to coarse grained, massive to foliated, grey to buff; local xenoliths and rafts of amphibolite, mica schist and psammitic gneiss

EARLY TO MIDDLE APHEBIAN

Wollaston Group

Wsh — **Quartz-muscovite schist**: pelitic to psammopelitic, fine to medium grained, ± biotite; locally intensely sheared

Wcm — **Calc-silicate gneiss, marble and amphibolite**: medium grained to pegmatitic, variable composition and texture, ± diopside ± albite ± hornblende ± biotite ± actinolite/tremolite ± calcite ± scapolite ± grossularite; occurring as conformable bands and lenses in supracrustal sequence

Wmi — **Meta-arkosic gneiss**: fine to medium grained, massive to foliated to gneissic, locally layered, ± biotite ± hornblende ± diopside ± muscovite ± sillimanite ± garnet ± cordierite ± myrmekite; commonly interlayered with metaquartzite, pelite and calc-silicate rock; commonly anatectic with more than 50 percent leucogranitic neosome

Wmg impure metaquartzite

Wpsm — **Pelitic to psammopelitic gneiss**: fine to medium grained, foliated to gneissic, ± biotite ± quartz ± feldspar ± graphite ± sillimanite ± muscovite ± cordierite ± garnet ± pyrite/pyrrhotite; interlayered metaquartzite, meta-arkosic calc-silicate rock and marble with psammitic types locally abundant; commonly anatectic with more than 50 percent leucogranitic to tonalitic neosome

Wpl graphite-bearing pelitic gneiss

Ws — **Mixed metasediment**: variable unit both along and across strike, generally fine grained and foliated, interlayered laminated slate, meta-argillite, metaquartzite, metachert, meta-arkose, calc-silicate rock, marble and rare banded iron formation

Wq — **Metaquartzite**: fine grained, massive to foliated, locally layered, ± muscovite ± biotite ± garnet ± feldspar ± hornblende ± sulphide; local conglomerate and interlayered graphitic meta-argillite ± andalusite ± garnet ± sulphide

PROBABLY EARLY APHEBIAN (LATE ARCHEAN?)

Wr — **Meta-arkose**: fine to medium grained, massive to foliated, locally layered and cross stratified, ± muscovite ± magnetite ± lithic fragments; local conglomerate with granite, granitic gneiss, metavolcanic and metasedimentary clasts; local interbedded pelitic schist ± andalusite ± staurolite ± garnet

Wv — **Amphibolite**: fine grained, massive to poorly foliated, locally pillowed; local intercalated pelitic gneiss

UNCONFORMITY

ARCHEAN, DEFORMED AND METAMORPHOSED WITH APHEBIAN SUPRACRUSTAL ROCKS DURING THE HUDSONIAN OROGENY

Wfn — **Felsic granitoid gneiss**: fine to coarse grained, foliated to gneissic, generally homogeneous rocks of syenogranitic to monzogranitic composition, ± biotite ± hornblende ± cordierite, with rare hypersthene (of granulite facies metamorphic origin); local inclusions of amphibolite and/or pelitic schist; locally partially anatectic with leucogranite neosome

wfb: biotite-hornblende bearing

RATTENSTONE DOMAIN

APHEBIAN (HUDSONIAN) WITH POSSIBLE ARCHEAN ELEMENTS

WATHAMAN BATHOLITH: ≈ 1880 Ma

Rgp — **Migmatite granitoid**: medium to coarse grained, massive to foliated, with local igneous flow banding; microcline megacrysts ± biotite ± hornblende-quartz-microcline, monzogranite and granodiorite; local pelitic, amphibolite and/or metabasite xenoliths; locally abundant pegmatite, aplite and non-megacrystic granitoid-forward margins

Rgm: sheared variants, with strongly foliated granulitmass and flattened microcline megacrysts, locally subsymoidic

Rgm — **Monzogranite-granodiorite**: containing abundant xenoliths and rafts of amphibolite and amphibolite gneiss derived from unit Pn

Rbd — **Quartz monzodiorite, diorite and gabbro**: fine to coarse grained, massive to foliated, ± hornblende ± plagioclase ± potassium feldspar ± quartz ± biotite; local paragneiss inclusions; contact zones commonly injected with granitic material

Rpeg — **Granitic pegmatite**: massive to foliated, ± biotite ± muscovite ± hornblende ± garnet

Rgt — **Tonalite, granodiorite and trondhjemite**: compositionally variable, generally potassium feldspar-poor, fine to coarse grained, locally pegmatitic, massive to foliated; ± biotite ± hornblende ± garnet ± muscovite; local amphibolite and/or pelite inclusions

Rgtc: metasomatized leucotonalite, pink, massive, porphyritic/aplitic, with introduced potassium feldspar and quartz in aureole of Wathaman Batholith

ROTTENSTONE MIGMATITE COMPLEX

Rmg — **Amphibolite and derived migmatite**: paleosome fine to medium grained, foliated, locally layered, ± hornblende ± feldspar ± biotite ± muscovite ± garnet ± quartz; local pelitic to psammopelitic gneiss; leucosome is commonly tonalitic, granodioritic or trondhjemitic; 00 to 90 percent of unit, medium to coarse grained to pegmatitic, massive to foliated; ± biotite ± muscovite ± garnet ± hornblende

Rng — **Pelitic to psammitic gneiss and derived migmatite**: paleosome fine to medium grained, well foliated to gneissic, locally layered, ± biotite ± quartz ± feldspar ± muscovite ± garnet ± sillimanite ± graphite; leucosome tonalitic, granodioritic or trondhjemitic; 00 to 90 percent of unit, medium to coarse grained to pegmatitic, massive to foliated; ± biotite ± muscovite ± garnet ± hornblende

PETER LAKE DOMAIN

APHEBIAN AND ARCHEAN ROCKS, STRONGLY REWORKED PROBABLY LATE IN THE HUDSONIAN

Ps — **Wylievine**: fine grained to aphanitic, streaky to finely laminated; interlayered mafic and felsic gneiss

Pgn — **Felsic gneiss**: medium grained, strongly foliated to gneissic; locally subsymoidic granitoid rock; ± biotite ± hornblende ± flattened microcline megacrysts

PAINTER LAKE GNEISSES

Pn — **Hornblende-biotite gneiss**: tonalitic to gabbroic composition, medium to coarse grained, with minor metagabbro and amphibolite

Pbn — **Mafic gneiss**: fine to medium grained, foliated to gneissic, locally subsymoidic; ± hornblende - plagioclase ± biotite ± quartz ± potassium feldspar; local pods of weakly foliated to massive mafic rock

Pmg - area cut by numerous granitic dykes and sills likely derived from the Wathaman Batholith

COURTENAY-CAVING LAKE BELT

HUDSONIAN WITH POSSIBLE ARCHEAN ELEMENTS

PETER LAKE COMPLEX

Pg — **Felsic granitoid**: generally medium to coarse grained, locally megacrystic, multiply intrusive, granodioritic to syenitic to monzonitic, ± biotite; local relict igneous textures, generally only weakly foliated except in discrete shear zones; ± biotite ± amphibole; local amphibolite and/or pelitic inclusions

Pgl: late granodioritic to alaskitic and aplitic intrusive sheets

Pgp — **Megacrystic granitoid** (microtexgneissoide in places from unit Pgp)

Pgpv: sheared variants

Pbg — **Mafic plutonic rocks**: fine to coarse grained, massive to weakly foliated; metabasite to metadiorite to metagabbro; local relict ophitic to subophitic texture; ± hornblende ± plagioclase ± biotite ± actinolite/tremolite, with local metaadamellitary xenoliths;

includes amphibolitic rocks of possible metavolcanic origin

Pba: layered metagabbro, minor anorthosite, ultramafic rock and granodiorite

Pbp: megacrystic diorite, massive with potassium feldspar megacrysts

Psf — **Quartzofeldspathic gneiss**: fine to medium grained, equigranular, generally well foliated to gneissic; locally compositionally banded; granitic to granodioritic ± biotite ± hornblende; local biotite-hornblende gneiss lenses

'Campbell River Group'

Psl — **Slate, phyllite and biotite schist**: very fine to fine grained, well foliated, locally layered; ± biotite ± muscovite ± pyrite/pyrrhotite ± carbonaceous material; local hornblende-biotite schist, psammite, amphibolite and metagabbro

SYMBOLS

Single bedrock exposure; approximate area of abundant bedrock exposure

Geological contact: defined to approximate; inferred

Structural lineament; possible to probable fault, as interpreted from geological, geophysical and/or airphoto evidence

Major fold axial trace: antiform; synform

Trend and approximate dip of dominant foliation surface: dip shallow (0-20°); moderate (20-59°); steep (60-84°); subvertical (85-90°)

Mineral prospect
1 George Lake: Zn-Pb (Kerr-Addison Mines Ltd., 1975)
2 Gemini: Au-Ag-Cu (Kerr-Addison Mines Ltd., 1975)
3 Ruth: Mo (Kerr-Addison Mines Ltd., 1975)
4 Gemini: Au-Ag-Cu (Kerr-Addison Mines Ltd., 1975)
5 Gemini: Au-Ag-Cu (Kerr-Addison Mines Ltd., 1975)
6 Gemini: Au-Ag-Cu (Kerr-Addison Mines Ltd., 1975)
7 Gemini: Au-Ag-Cu (Kerr-Addison Mines Ltd., 1975)
8 Gemini: Au-Ag-Cu (Kerr-Addison Mines Ltd., 1975)