HISTOGRAM

Complexes: where two or more classes of terrain are interspersed in a mosaic or repeating pattern the proportion of each component in the combination is given in a three-position designation set off by slashes denoting arbitrary percentage limits. For example, "Mv/0/0" means that at least 60% of the area is underlain by thin till, with up to 40% boggy areas, and less than 15% scattered rock outcrops. "Rc/R" indicates more than 60% bedrock concealed by vegetation and less than 15% outcrop.

Canada
Department of Energy, Mines and Resources
Geological Survey of Canada

Province of Saskatchewan
Saskatchewan Energy and Mines
Saskatchewan Geological Survey

64 E, and parts of 74 A, H

64 E and parts of 74 A, H

MANGANESE (ppm)
GSC OPEN FILE 1106

REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 71-1984
CANADA - SASKATCHEWAN
MINERAL DEVELOPMENT AGREEMENT (1984-89)
LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY
NORTHEASTERN SASKATCHEWAN, 1984

Scale 1:250 000

Kilometres 5 0 5 10 15 20 Kilometres

Universal Transverse Mercator Projection

© Crown Copyrights reserved

Mean magnetic declination 1985, 15°06' East,
decreasing 23.1' annually. Readings
vary from 13°05' in the SE corner to 17°06' in the NW
corner of the map area

Geological Survey of Canada
Resource Geophysics and Geochemistry Division
and
Saskatchewan Energy and Mines
Saskatchewan Geological Survey

CONTRACTORS

Sample collection by Marshall Macklin Monaghan Ltd., Toronto
Sample preparation by Golder Associates, Ottawa

Sediment chemical analysis by Barringer Magenta ltd., Rexdale, Ontario
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
880 Wellington St.
Bay 238
Ottawa, Ontario
K1R 6K7

The data are also available in digital form. For further information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4

MANGANESE (ppm)
GSC OPEN FILE 1106
NORTHEASTERN SASKATCHEWAN, 1984

LEGEND

Note: This legend is common for Regional Geochemical Reconnaissance Map 71-1984, Open File 1106

NECHELIKIAN/HADRYNIAN

- dd — **Diabase-gabbro**: fine to coarse grained, massive to weakly foliated; ± olivine; ± biotite; ± hypersthene

PALEOHELIKIAN

ATHABASCA GROUP

- MF — **Manitou Falls Formation**: sandstone and conglomerate
  MFu: fluvial sandstone member, locally pebbly
  MFb conglomerate member

LATE APHEBIAN (HUDSONIAN)

- X — **Cataclasite**: mylonite and sheared rocks of the Needle Falls Shear Zone; derived from rocks of the Wollaston and Peter Lake Domains

ROTTENSTONE DOMAIN

APHEBIAN (HUDSONIAN) WITH POSSIBLE ARCHEAN ELEMENTS

WATHAMAN BATHOLITH (c.1980 Ma)

- Pgp — **Megacrystic granitoid**: medium to coarse grained, massive to foliated, with local gneiss flow banding; microcline megacrysts ± biotite ± hornblende quartz monzonite, granodiorite composition; ± biotite ± hornblende xenoliths and/or metasedimentary xenoliths; locally abundant pegmatite, aplite and non-megacrystic granitoid toward margins
  Pgpa: sheared variants, with strongly foliated granulitoid and fissured microcline megacrysts, locally subsymplectic
- Pgm — **Monzogranite-granodiorite**: containing abundant xenoliths and rafts of amphibolite and amphibole-gneiss derived from unit Pm
- Pqd — **Quartz monzodiorite, diorite and gabbro**: fine to coarse grained, massive to foliated; ± hornblende ± pigeonites ± potassium feldspar ± quartz ± biotite; local paragneiss inclusions; contact zones commonly injected with granitic material
- Pgeg — **Granitic pegmatite**: massive to foliated; ± biotite ± muscovite ± hornblende ± garnet

ROTTENSTONE MIGMATITE COMPLEX

- Pgt — **Tonalite, granodiorite and trondhjemite**: compositionally variable, generally potassium feldspar-poor, fine to coarse grained, locally pegmatitic, massive to foliated; ± biotite ± hornblende ± garnet; local hornblende, locus amphibolite and/or pinite inclusions
  Pgtz: metasomatized leucotonalite, pink, massive, porphyritoidastic, with introduced potassium feldspar and quartz in aureole of Wathaman Batholith
- Pmg — **Amphibolite and derived migmatite**: paleosome fine to medium grained, well foliated to gneissic, locally layered; ± biotite ± muscovite ± garnet; ± hornblende
  Pmgt: migmatite; locally layered; quartz-rich; biotite to psammitic gneiss and calc-silicate gneiss; neosome tonalitic, granodioritic or trondhjemitic (30 to 90 percent of unit); medium to coarse grained to pegmatitic, massive to foliated; ± biotite ± muscovite ± garnet ± hornblende
- Pmg — **Pelitic to psammitic gneiss and derived migmatite**: paleosome fine to medium grained, well foliated to gneissic, locally layered; ± biotite ± muscovite ± sillimanite ± cordierite ± garnet ± sillimanite ± graphite; neosome tonalitic, granodioritic or trondhjemitic (30 to 90 percent of unit), medium to coarse grained to pegmatitic, massive to foliated; ± biotite ± muscovite ± garnet ± hornblende

PETER LAKE DOMAIN

APHEBIAN AND ARCHEAN ROCKS, STRONGLY REWORKED PROBABLY LATE IN THE HUDSONIAN

- Pv — **Mylonite**: fine grained to aphanitic, streaky to finely laminated; interlayered mafic and felsic gneiss
- Pgn — **Felsic gneiss**: medium grained, strongly foliated to gneissic, locally subsymplectic granitoid rock ± biotite ± hornblende ± flaserod microcline megacrysts
- Pn — **Hornblende-biotite gneiss**: tonalitic to gabbroic composition, medium to coarse grained, with minor metagabbro and amphibolite
- Pbm — **Mafic gneiss**: fine to medium grained, foliated to gneissic, locally subsymplectic; ± hornblende ± plagioclase ± biotite ± quartz ± potassium feldspar; local pods of weakly foliated to massive mafic rock
- Pmg: area cut by numerous granitic dykes and sills likely derived from the Wathaman Batholith

PARKER LAKE GNEISSES

COURTENAY-GAVINS LAKE BELT

PROBABLY EARLY APHEBIAN (LATE ARCHEAN?)

- Wr — **Meta-arkose**: fine to medium grained, massive to foliated, locally layered; ± biotite ± muscovite ± sillimanite ± cordierite ± magnetite ± relict fragmental, local conglomerate with granite, granite gneiss, metavolcanic and metasedimentary clasts; local interbedded pelitic schist ± andalusite ± staurolite ± garnet
- Ww — **Amphibolite**: fine grained, massive to poorly foliated, locally pillowed; local interlaminated pelitic gneiss

UNCONFORMITY

ARCHEAN, DEFORMED AND METAMORPHOSED WITH APHEBIAN SUPRACRUSTAL ROCKS DURING THE HUDSONIAN OROGENY

- Wfm — **Felsic granuloid gneiss**: fine to coarse grained, foliated to gneissic, generally homogenous rocks of pyroxene-to granodiorite composition ± biotite ± amphibole ± hornblende ± garnet; locally derived from rocks of metamorphic origin; local inclusions of amphibolite and/or pelitic schist; locally partially anatectic with leucogranitic neosome

  Wfb biotite-hornblende bearing

WOLLASTON DOMAIN

LATE APHEBIAN (HUDSONIAN)

- Wpeg — **Granitic pegmatite**: variable grain size, generally massive, ± biotite ± muscovite ± garnet; contacts with migmatitic supracrustal gneisses commonly gradational
- Wg — **Granite and aplasite**: fine to coarse grained, massive to weakly foliated; ± biotite ± hornblende ± magnetite ± muscovite ± garnet; contacts with migmatitic supracrustal gneisses commonly gradational
- Wgdb — **Biotite granodiorite-tonalite**: medium to coarse grained, massive to foliated; grey to buff; local xenoliths and rafts of amphibolite, mica schist and psammitic gneiss

EARLY TO MIDDLE APHEBIAN

Wollaston Group

- Wwn — **Quartz-muscovite schist**: pelitic to psammopelitic, fine to medium grained, ± biotite; locally intensely sheared
- Wcm — **Calc-silicate gneiss, marble and amphibolite**: medium grained to pegmatitic, variable composition and texture, ± diopside ± hornblende ± scapolite ± garnet ± phlogopite ± calcite ± scapolite ± grossularite; occurring as conformable bands and lenses in supracrustal sequence
- Wmx — **Meta-arkosic gneiss**: fine to medium grained, massive to foliated to gneissic, locally layered; ± biotite ± hornblende ± diopside ± muscovite ± sillimanite ± garnet ± cordierite ± magnetite ± pyrite; locally interlayered with metapsammite, pelite and calc-silicate rock; commonly anatectic with more than 50 percent leucogranitic neosome

  Wmq: impure metaquartzite
- Wpsm — **Pelitic to psammopelitic gneiss**: fine to medium grained, foliated to gneissic; ± biotite ± quartz ± feldspar ± graphite ± sillimanite ± muscovite ± cordierite ± garnet ± pyrrhotite/pyrite; local interlayered metaquartzite, meta-arkose, calc-silicate rock and marble with psammitic types locally anatectic with more than 50 percent leucogranitic to tonalitic neosome

  Wpt: graphite-bearing pelitic gneiss
- Wis — **Mixed metasediment**: variable unit both along and across strike, generally fine grained and foliated; interlayered metapsammite, meta-argillite, metaquartzite, metachert, meta-arkose, calc-silicate rock, marble and rare banded iron formation
- Wq — **Metaquartzite**: fine grained, massive to foliated, locally layered; ± muscovite ± biotite ± garnet ± feldspar ± hornblende ± sulphide; local conglomerate and interlayered graphitic meta-argillite ± andalusite ± garnet ± sulphide

PROBABLY EARLY APHEBIAN (LATE ARCHEAN?)

- Wr — **Meta-arkose**: fine to medium grained, massive to foliated, locally layered; ± biotite ± muscovite ± sillimanite ± magnetite ± relict fragmental, local conglomerate with granite, granite gneiss, metavolcanic and metasedimentary clasts; local interbedded pelitic schist ± andalusite ± staurolite ± garnet
- Ww — **Amphibolite**: fine grained, massive to poorly foliated, locally pillowed; local interlaminated pelitic gneiss

UNCONFORMITY

ARCHEAN, DEFORMED AND METAMORPHOSED WITH APHEBIAN SUPRACRUSTAL ROCKS DURING THE HUDSONIAN OROGENY

- Wfm — **Felsic granuloid gneiss**: fine to coarse grained, foliated to gneissic, generally homogenous rocks of pyroxene-to granodiorite composition ± biotite ± amphibole ± hornblende ± garnet; locally derived from rocks of metamorphic origin; local inclusions of amphibolite and/or pelitic schist; locally partially anatectic with leucogranitic neosome

  Wfb biotite-hornblende bearing

LA RONGE DOMAIN

PROBABLY HUDSONIAN s.s.? (c.1740 Ma)

- Lgd — **Granodiorite and quartz monzonite**: medium grained, well foliated to gneissic; biotite ± hornblende; abundant xenoliths of paragneiss and amLLgm
- Lgm — **Quartz monzodiorite**: fine to coarse grained, foliated to gneissic; ± hornblende ± biotite; abundant xenoliths of paragneiss and amphibolitic resiosmes of granodioritic to quartz monzonitic material in migmatitic contact zone

HUDSONIAN WITH POSSIBLE ARCHEAN ELEMENTS

- Pg — **Felsic granitoid**: generally medium to coarse grained, locally megacrystic; multiply intrusive; granodioritic to syenitic to alaskitic and aplitic; local relict igneous textures, generally only weakly foliated except in discrete shear zones; ± biotite ± amphibole; local amphibolite and/or pelite inclusions
  Pgt: late granodioritic to alaskitic and aplitic intrusive sheets
- Pgp — **Megacrystic granitoid** (melting/quishable in places from unit Pgp)
  Pgpa: sheared variants

PETER LAKE COMPLEX

PROBABLY MAINLY ARCHEAN

- Pbd — **Mafic plutonic rocks**: fine to coarse grained, massive to weakly foliated; metatonalite to metadiorite to metagabbro; local relict ophitic to subophitic texture; ± hornblende ± plagioclase ± biotite ± actinolite/tremolite, with local metasedimentary xenoliths;
  includes amphibolitic rocks of possible metavolcanic origin
  Pba: layered metagabbro, minor anorthosite, ultramafic rock and granodiorite
  Pbp: megacrystic diorite, massive with potassium feldspar megacrysts
- Pgf — **Quartzofeldspathic gneiss**: fine to medium grained, foliated to gneissic, locally layered; locally migmatitic; compositionally banded; granitic to granodioritic ± biotite ± hornblende; local biotite-hornblende gneiss lenses

'Campbell River Group'

- Pgf — **Slate, phyllite and biotite schist**: very fine to fine grained, well foliated, locally layered; ± biotite ± muscovite ± pyrite/pyrrhotite ± carbonaceous material; local hornblende-biotite schist, psammite, amphibolite and metagabbro

SYMBOLS

- Single bedrock exposure; approximate area of abundant bedrock exposure
- Geological contact: defined to approximate; inferred
- Structural lineament; possible to probable fault, as interpreted from geological, geophysical and/or airphoto evidence

  Major fold axial trace: antiform; synform

  Trend and approximate dip of dominant foliation surface: dip shallow (0-29°); moderate (30-59°); steep (60-84°); subvertical (85-90°)
- Mineral prospect:
  1. George Lake: Zn-Pb (Kaiup-Maller and Brummer, 1970; Coomber, 1971)
- Sample location (geochronology):
  1. 1625±10 Ma, K-Ar muscovite (Wanless et al., 1970)
  2. 1760±30 Ma, Rb-Sr whole rock isochron (Bell and Macdonald, 1982); map location is approximate centre of collecting area
- * No analytical result

* A mnemonic name recorded as rock types as part of field observations

This legend was modified and the geology derived for these geochemical maps from Compilation Bedrock Geology Series 228A, 229A and 232A, Saskatchewan Energy and Mines, Saskatchewan Geological Survey

This map forms one of a series of maps released by the Geological Survey of Canada, Open File 1106. The Open File consists of maps of various geochemical variables: 16 for lake sediment, 3 for lake water and 1 sample site location

MANGANESE (ppm)
GSC OPEN FILE 1106
NORTHEASTERN SASKATCHEWAN, 1984

This map has been reprinted from a scanned version of the original map
Reproduction par numérisation d'une carte sur papier