


The regional geochemical trend map displayed above utilized a moving weighted average using an inverse distance function (1/d²) to filter out minor irregularities and emphasize broad-scale regional features. Single point anomalies may be suppressed or eliminated, however, geological units which are chemically enriched, or large metallic deposits undergoing weathering would be expected to produce identifiable anomalies.



NONGLACIAL ENVIRONMENT

ORGANIC DEPOSITS: marsh, fen, bog and swamp deposits up to 6 m thick, seasonally flooded; commonly overlying fine textured proglacial lake and glacial till deposits

PROGLACIAL AND GLACIAL ENVIRONMENT

GLACIOLACUSTRINE DEPOSITS: clay silt, sand and minor gravel, 1 - 30 m thick, deposited in proglacial lakes

Beach and nearshore deposits: sand and gravel 1 - 4 m; sand blanket 0 - 1 m thick; includes areas of wave washed till and exposed bedrock

Deep basin deposits: silt, clay and sand, 1 - 30 m thick, forming extensive lake plains and discontinuous veneer reflecting underlying topography; may include areas of iceberg scouring, thin alluvium, wave washed till and exposed bedrock

GLACIOFLUVIAL DEPOSITS: gravel, sand and silt 1 - 100 m thick, deposited in an ice-marginal environment; includes subaqueous outwash deposited in glacial lakes, and some late glacial valley fill

GLACIAL ENVIRONMENT

GLACIAL DEPOSITS: unsorted glacial debris, 1 - 10 m thick, reflecting composition of underlying bedrock; predominantly lodgment till; also includes extensive areas of hummocky stagnation moraine, ribbed moraine and water deposited till

Till: 1 - 10 m thick, highly calcareous, derived primarily from Paleozoic carbonate rock; typical composition: silt - 44%, sand - 21%, clay - 19%, gravel - 16%; includes bedrock exposures, evidence of wave-wearing

Till: 1 - 5 m thick, derived primarily from Precambrian bedrock; typical composition: sand - 46%, gravel - 29%, silt - 24%, clay - 4%; includes extensive areas of bedrock outcrop, surface reflects form of underlying bedrock surface

NONGLACIAL ENVIRONMENT

BEDROCK: Precambrian, Paleozoic and Mesozoic bedrock of various lithologies. Detailed geological legend at right

SYMBOLS

Surficial geological boundary

Striae

Flutings, drumlins and drumlinoid ridges

. Moraines (including end, interlobate and recessional)

Beach ridges, bars and strandlines

Esker

Surficial geology derived from:
Nelson, L. & al. (1981) Surficial Geological Map of Manitoba, Aggregate Resources Section, Manitoba Mineral Resources Division, Map 01-1 (1:1,000,000 scale)

CENOZOIC

10 OVBD 44* Overburden; mainly glacial till and glaciolacustrine deposits

PALEOZOIC

ORDOVICIAN

9 DML 14 RED RIVER FORMATION: Mottled dolomitic limestone to dolomite, in part cherty and calcareous

PROTEROZOIC

8 ACIV 04 Felsic to intermediate plutonic rocks

7 IMIV 04 Intermediate plutonic rocks

6 BCIV 04 Mafic to intermediate plutonic rocks. Includes ultramafic rocks

5 AMPB 04 Amphibolite. Includes chert, marble

4 MARK 04 Meta-arkose and quartz-feldspathic gneiss

3 MGCK 04 Meta-greywacke and quartz-biotite gneiss


2 IEVX 04 Intermediate to felsic volcanic rocks

1 BEVX 04 Mafic to intermediate volcanic rocks

*A mnemonic code assigned to rock types and recorded as part of field observations

Geological boundary
Surficial deposit boundary
No analytical results

Provisional Synoptic Geological Compilation at 1:250,000 scale, by S. Parker, Geological Services, Manitoba Energy and Mines, 1985



Mean magnetic declination 1985, 10°30' East, decreasing 20.5' annually. Readings vary from 8°31'E in the SE corner to 12°32'E in the NW corner of the map

Geological Survey of Canada
Resource Geophysics and Geochemistry Division
and
Manitoba Department of Energy and Mines
Mineral Resources Division

CONTRACTORS

Sample collection by MPH Consulting Ltd., Toronto
Sample preparation by Golder Associates

Sediment chemical analyses by Barringer Magenta Ltd., Rexdale, Ontario
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary
Gold analysis by Chemex Labs Limited, Vancouver, B.C.

This map forms one of a series of maps released by the Geological Survey of Canada, Open File 1212. The Open File consists of maps of various geochemical variables: 18 for lake sediment, 3 for lake water and 1 sample site location

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
880 Wellington St.
Box 230
Ottawa, Ontario
K1R 6K7

The data are also available in digital form. For further information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4

