


The regional geochemical trend map displayed above utilized a moving weighted average using an inverse distance function ($1/d^2$) to filter out minor irregularities and emphasize broad-scale regional features. Single point anomalies may be suppressed or eliminated, however, geological units which are chemically enriched, or large metallic deposits undergoing weathering would be expected to produce identifiable anomalies.



NONGLACIAL ENVIRONMENT

ORGANIC DEPOSITS: marsh, fen, bog and swamp deposits up to 6 m thick, seasonally flooded; commonly overlying fine textured proglacial lake and glacial till deposits

PROGLACIAL AND GLACIAL ENVIRONMENT

GLACIOGLACIESTRINE DEPOSITS: clay silt, sand and minor gravel, 1 - 30 m thick, deposited in proglacial lakes

Beach and nearshore deposits: sand and gravel 1 - 4 m; sand blanket 0 - 1 m thick; includes areas of wave washed till and exposed bedrock

Deep basin deposits: silt, clay and sand, 1 - 30 m thick, forming extensive lake plains and discontinuous veneer reflecting underlying topography; may include areas of iceberg scouring, thin alluvium, wave washed till and exposed bedrock

GLACIOFLUVIAL DEPOSITS: gravel, sand and silt 1 - 100 m thick, deposited in an ice-marginal environment; includes subaqueous outwash deposited in glacial lakes, and some late glacial valley fill

GLACIAL ENVIRONMENT

GLACIAL DEPOSITS: unsorted glacial debris, 1 - 10 m thick, reflecting composition of underlying bedrock; predominantly lodgment till; also includes extensive areas of hummocky stagnation moraine, ribbed moraine and water deposited till

Till: 1 - 10 m thick, highly calcareous, derived primarily from Paleozoic carbonate rock; typical composition: silt - 44%, sand - 21%, clay - 19%, gravel - 16%; includes bedrock exposures, evidence of wave-washing

Till: 1 - 5 m thick, derived primarily from Precambrian bedrock; typical composition: sand - 46%, gravel - 28%, silt - 22%, clay - 8%; includes extensive areas of bedrock outcrop, surface reflects form of underlying bedrock surface


NONGLACIAL ENVIRONMENT

BEDROCK: Precambrian, Paleozoic and Mesozoic bedrock of various lithologies. Detailed geological legend at right

SYMBOLS

Surficial geological boundary
Striae
Flutings, drumlins and drumlinoid ridges
Moraines (including end, interlobate and recessional)
Beach ridges, bars and strandlines
Esker

Surficial geology derived from:
Nelson, E. et al. (1983) Surficial Geological Map of Manitoba, Aggregate Resources Section, Manitoba Mineral Resources Division, Map 81-1 (1:1,000,000 scale)



LEGEND

- CENOZOIC
10 QVB9 44* Overburden; mainly glacial till and glaciolacustrine deposits
- PALEOZOIC
- ORDOVICIAN
9 QMLM 14 RED RIVER FORMATION: Mottled dolomitic limestone to dolomite, in part cherty and calcareous
- PROTEROZOIC
- 8 ACIV 04 Felsic to intermediate plutonic rocks
7 IMIV 04 Intermediate plutonic rocks
6 BCIV 04 Mafic to intermediate plutonic rocks. Includes ultramafic rocks
5 AMPB 04 Amphibolite. Includes chert, marble
4 MARK 04 Meta-arkose and quartz-feldspathic gneiss
3 MGCK 04 Meta-gneiss and quartz-biotite gneiss
2 IEKV 04 Intermediate to felsic volcanic rocks
1 BEKV 04 Mafic to intermediate volcanic rocks

*A mnemonic code assigned to rock types and recorded as part of field observations


Geological boundary
Surficial deposit boundary
No analytical results

Au value (ppb) denotes an analysis performed on a sample weight <10 g. () identifies Au values corresponding to repeat analyses. <n denotes a result less than detection level n (ppb). consult text for actual sample weight when Au values denoted by * or < detection level

Examples:
21 Au value of 21 ppb determined on sample weight <10 g.
38(27) Au value of 38 ppb on first analysis, Au value of 27 ppb on repeat analysis for sample weighing <10 g.
*4 Au value less than detection limit of 4 ppb.


Provisional Synoptic Geological Compilation at 1:250,000 scale, by S. Parker, Geological Services, Manitoba Energy and Mines, 1985

Please refer to Open File text for discussion of gold presentation format and geochemical interpretation.



GOLD (ppb)

GSC OPEN FILE 1212
REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 77-1985
CANADA-MANITOBA
MINERAL DEVELOPMENT AGREEMENT (1984-89)
LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY
WEST-CENTRAL MANITOBA, 1985
Scale 1:250,000



GOLD (ppb)
GSC OPEN FILE 1212
WEST-CENTRAL MANITOBA, 1985

Contribution to Canada-Manitoba Mineral Development Agreement 1984-89, a subsidiary agreement under the Economic and Regional Development Agreement. Project funded by the Geological Survey of Canada.

Manitoba Energy and Mines
Energy, Mines and Resources Canada
Energie, Mines et Ressources Canada

Canada

Universal Transverse Mercator Projection
© Crown Copyright reserved