


ENVIRONMENT CATEGORY	GEOMORPHIC MODIFIER
Organic	c concealed
Glaciolacustrine	w weathered
Glaciolacustrine	e eroded
Glaciolacustrine	g gullied
Glaciolacustrine	k collapsed
Glaciolacustrine	pl plain
Moraine	v veneer
Rock	r ridged
Rock	h hummocky
Rock	dr drumlinoid
Rock	t terrace
Soil	

Complexes: where two or more classes of terrain are interspersed in a mosaic pattern the proportion of each component in the combination is given in a three-position designation set off by slashes denoting arbitrary percentage limits. For example, 40/30/30 means that at least 40% of the area is underlain by this soil, with up to 40% boggy areas, and less than 30% scattered rock outcrops. 10/90 indicates more than 90% bedrock concealed by veneer and up to 40% bedrock exposures.

GLACIAL FEATURE SYMBOLS
 Boundary of overburden unit
 Drumlin, drumlinoid ridge, fluting
 Striation, groove (ice direction inferred)
 End moraine
 Esker, crevasse filling

Surficial geology modified from:
 Schertzer, J.T. (1984) Quaternary Geology of the Precambrian Shield, Map 221A (1:1,000,000 scale), to accompany Report 221, Saskatchewan Energy and Mines.



The regional geochemical trend map displayed above utilized a moving weighted average using an inverse distance function (1/r²) to filter out minor irregularities and emphasize broad-scale regional features. Single point anomalies may be suppressed or eliminated, however, geological units which are chemically enriched, or large metallic deposits undergoing weathering would be expected to produce identifiable anomalies.

Geological Survey of Canada
 Resource Geophysics and Geochemistry Division
 Department of Mineral Resources
 Saskatchewan Geological Survey

CONTRACTORS
 Sample collection by NW Consulting Ltd., Toronto
 Sample preparation by Gelder Associates

Sediment chemical analyses by Barringer Regenta Ltd., Redlake, Ontario
 Water chemical analyses by Barringer Regenta Laboratories (Alberta) Ltd., Calgary


This map forms one of a series of maps released by the Geological Survey of Canada, Open File 1213. The Open File consists of maps of various geochemical variables: 16 for lake sediments, 3 for lake water and sample site location

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
 880 Wellington St.
 Box 230
 Ottawa, Ontario
 K1R 8K7

The data are also available in digital form. For further information please contact:
 The Director
 Computer Science Centre
 Department of Energy, Mines and Resources
 Ottawa, Ontario
 K1A 0G8

Contribution to Canada-Saskatchewan Mineral Development Agreement 1984-89, a subsidiary agreement under the Economic and Regional Development Agreement. Project funded by the Geological Survey of Canada.



LEGEND

CEANOZOIC

QUATERNARY

21 (RMR 44) Recessional moraine: unconsolidated sand and gravel of the Ore Lake moraine.

MESOZOIC

LOWER CRETACEOUS

20 (SWS 36) MANVILLE GROUP: Sandstone, variably argillaceous and carbonaceous quartzose sands, local mudstone and lignitic interbeds.

PALEOZOIC

MIDDLE DEVONIAN

19 (DLMT 18) WINNIPEGOSIS FORMATION: Dolomite, dolomitized limestone, calcareous shale.

18 (LGN 18) MEADOW LAKE FORMATION (Upper member): Limestone, dolomite, mudstone, argillaceous dolomite, dolomitic limestone and dolomitic argillite mudstone.

17 (AGC 18) MEADOW LAKE FORMATION (Lower member): Argillaceous dolomite with local interbeds of mudstone, sandstone and limestone, minor gypsum bands.

CAMBRIAN

16 (SWS 12) DEADWOOD FORMATION: Quartz sandstone, unstratified polymictic conglomerate near base, minor sandy dolomite. Overlies severely weathered, saproplitic Precambrian basement.

PRECAMBRIAN

MAINLY APHEBIAN (HUDSONIAN)

15 (UMFC 04) Ultramafic and mafic rocks, includes gabbro, diorite, pyroxenitic quartz diorite and basalt, locally sericitized.

14 (IREM 04) Banded iron formation, alternating bands of magnetite and hematite.

13 (LLOC 04) Calc-silicate and marble, local interbedded meta-argillite and dolomitic marble.

12 (PRGS 04) Pelitic gneiss and meta-arkosic gneiss, with interbedded calc-silicate rock and pelitic gneiss.

11 (PCSC 04) Pelitic to psammopelitic gneiss and schist, generally contain more than 10 percent mafic minerals, biotite ± garnet ± cordierite ± sillimanite ± graphite ± tourmaline.

10 (MGR 04) Metapsiltstone and minor orthoquartzite with a persistent basal polymictic conglomerate.

9 (BGAS 04) Biotite gneiss of possible volcano-sedimentary origin, conformable granitoid sheets may comprise up to 50 percent of unit.

8 (MPRK 04) Mixed metasedimentary and metavolcanic rocks, probably derived from mudstone, sandstone, arkose, clastic rocks.

7 (BGR 04) MICHAMAN BATHOLITH: Biotite-hornblende granodiorite.

6 (BNGT 04) MICHAMAN BATHOLITH: Biotite monzogranite-granodiorite.

5 (SGAT 04) Syenogranite and monzogranite, generally leucocratic.

4 (MGRM 04) Migmatite and mylonite zones; complexes of mixed metasediments and granitic rocks, and strongly flattened or augered Michaman Batholith rocks.

MAINLY ARCHEAN, DEFORMED WITH APHEBIAN ROCKS DURING HUDSONIAN OROGENY

3 (APRH 03) Amphibolite and hornblende bearing gneisses, meta-gabbro and metadiorite.


2 (GRNG 03) Granitoid gneiss, syenogranite to granodioritic in composition, may include albite to amphibolite inclusions.

1 (DORT 03) Diorite


*A mnemonic code assigned to rock types and recorded as part of field observations.

Geological boundary; approximate, assumed
 Fault
 No analytical result

Geological base and legend are derived from: Thomas, W.L. and Stille, W.L. (1983): Compilation Bedrock Geology, 1:500,000 scale map with marginal notes; Saskatchewan Energy and Mines, Report 249 (1:500,000 scale map with marginal notes).
 Thomas, W.L. and Stille, W.L. (1985): Compilation Bedrock Geology, 1:500,000 scale map with marginal notes; Saskatchewan Energy and Mines, Report 255 (1:500,000 scale map with marginal notes).
 Thomas, W.L. and Stille, W.L. (1987): Compilation Bedrock Geology, 1:500,000 scale map with marginal notes; Saskatchewan Energy and Mines, Report 258 (1:500,000 scale map with marginal notes).
 Thomas, W.L. (1984): Preliminary Compilation Bedrock Geology, 1:500,000 scale map with marginal notes; Saskatchewan Energy and Mines, Report 228 (1:500,000 scale map with marginal notes).
 Thomas, W.L. (1986): Preliminary Compilation Bedrock Geology, 1:500,000 scale map with marginal notes; Saskatchewan Energy and Mines, Report 252 (1:500,000 scale map with marginal notes).
 Thomas, W.L. and Stille, W.L. (1980): Geological Map of Northern Saskatchewan, 1:500,000 scale map with marginal notes.



pH in water
 GSC OPEN FILE 1213
 REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 78-1985
 CANADA - SASKATCHEWAN
 MINERAL DEVELOPMENT AGREEMENT (1984-89)
 LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY
 NORTH-CENTRAL SASKATCHEWAN, 1985
 Scale 1:250 000
 Universal Transverse Mercator Projection
 © Crown Copyright reserved



Elevation in feet above mean sea level
 Mean magnetic declination 1985, 17°20' East, decreasing 20.4' annually. Readings vary from 15°30' East in the SE corner to 20°00' East in the NW corner of the map area

Base map assembled by the Geological Survey of Canada from maps published at the same scale by Mapping and Charting Establishment, Department of National Defence and the Survey and Mapping Branch, Department of Energy, Mines and Resources in 1974, 1977, 1982

This map has been prepared from a published version of the original map. Reproduction for dissemination of data with no alterations.
 5 of 20
 pH in water
 GSC OPEN FILE 1213
 NORTH-CENTRAL SASKATCHEWAN, 1985