

The regional geochemical trend map displayed above utilized a moving weighted average using an inverse distance function (1/d<sup>2</sup>) to filter out minor irregularities and emphasize broad-scale regional features. Single point anomalies may be suppressed or eliminated, however, geological units which are chemically enriched, or large metallic deposits undergoing weathering would be expected to produce identifiable anomalies.

#### CONTRACTORS

Sample collection by McIlhenny Surveying and Engineering Ltd., Vancouver

Sample preparation by Golder Associates, Ottawa (1984) and  
Kamloops Research and Assay Laboratory Ltd., Kamloops B.C. (1985)

Sediment chemical analyses by Barringer Magenta Ltd., Rexdale, Ontario (1984) and  
Bondar-Clegg and Company Ltd., North Vancouver, B.C. (1985)

Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary (1984) and  
Chemex Labs Ltd., North Vancouver, B.C. (1985)

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation  
880 Wellington St.  
Bay 238  
Ottawa, Ontario  
K1R 6K7

The data are also available in digital form. For further information please contact:

The Director  
Computer Science Centre  
Department of Energy, Mines and Resources  
Ottawa, Ontario  
K1A 0E4

Contribution to Canada-British Columbia Mineral Development Agreement 1985-89, a subsidiary agreement under the Economic and Regional Development Agreement. West half of the project jointly funded under a letter of understanding by the Geological Survey of Canada and British Columbia. East half of the project funded by British Columbia




Province of British Columbia  
Ministry of Energy, Mines and Petroleum Resources



Energy, Mines and Resources Canada  
Energie, Mines et Ressources Canada

Canada

This document was produced by scanning the original publication. Ce document est le produit d'une numérisation par balayage de la publication originale.



Undivided Glaciofluvial, Glaciolacustrine and morainal deposits: sand, silt, gravel and clay with local accumulations up to 70 m thick along valley bottoms: may include meltwater or outwash channel deposits, alluvium, varved silt, clay and sand, glacial till and ground moraine

Predominantly Rock: below 1,700 m elevation, discontinuous veneers of glacial till and ground moraine, and minor colluvium; valley bottoms may contain drumlinoid features, fluting, kame and esker complexes

Glaciers, permanent snowfields

Ice flow direction inferred from various data

Fluting, glacial striae

Sources of information:  
Prest, V.K., Grant, D.R. and Rampton, V.N. (1967) Glacial Map of Canada, Geological Survey of Canada, Map 1253A (Scale 1:5,000,000).  
Campbell, R.B. and Mountjoy, E.W. (1967) Geology - McBride, British Columbia, Geological Survey of Canada, Map 1356A (Scale 1:250 000)

