

The regional geochemical trend map displayed above utilized a moving weighted average using an inverse distance function ($1/d^3$) to filter out minor irregularities and emphasize broad-scale regional features. Single point anomalies may be suppressed or eliminated; however, geological units which are chemically enriched, or large metallic deposits undergoing weathering would be expected to produce identifiable anomalies.

Geological Survey of Canada
Resource Geophysics and Geochemistry Division
CONTRACTORS

Sample collection by Rogers Exploration Services Ltd., Whitehorse
Sample preparation by Golder Associates, Ottawa
Gold analysis by Chemex Labs Limited, Vancouver, B.C.

Sediment chemical analyses by Barringer Magenta Ltd., Rexdale, Ontario
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary




This map forms one of a series of maps released by the Geological Survey of Canada, Open Files 1217 to 1220. Each Open File consists of maps of various geochemical variables: 21 for stream sediment, 3 for stream water and 1 sample site location.

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
880 Wellington St.
Bay 238
Ottawa, Ontario
K1A 6K7

The data are also available in digital form. For further information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4



- [Symbol] Undivided surficial deposits; alluvium, glacial till and moraine, outwash and ice contact deposits, volcanic ash, loess, colluvium
- [Symbol] Bedrock exposures; includes discontinuous veneer of undivided glacial drift

SYMBOLS


- Surficial deposit boundary
- Limit of Pre-Reid ice advance
- Limit of McConnell (Ruby) ice advance
- Meltwater channels, outwash deposits, indicating direction of flow
- Glaciation lineation parallel to ice flow direction, includes fluting, crag and tail, roches moutonnées and drumlinoid forms, direction of flow indicated
- Drumlinoid form, direction of movement inferred, not inferred

- Sources of information:
Hughes, O.L., Campbell, R.B., Muller, J.E., and Wheeler, J.O. (1968) Glacial Map of Yukon Territory, Geological Survey of Canada, Map 6-1968, (1:1 000 000 scale) to accompany GSC Paper 68-34
Prest, V.K., Grant, D.R., and Rampton, V.N. (1967) Glacial Map of Canada, Geological Survey of Canada (1:5 000 000 scale)
Templeman-Kluit, D.J. (1973) Geology - AISIHIK LAKE, Yukon Territory, Geological Survey of Canada, Map 17-1973, (1:250 000 scale) to accompany Paper 73-41



Contribution to the Canada/Yukon Subsidiary
Agreement on Mineral Resources 1985-1989
under the Canada/Yukon Economic Development
Agreement

Yukon
Government




INDEX MAP

TIN (ppm)
GSC OPEN FILE 1219
REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 84-1985
CANADA-YUKON
MINERAL DEVELOPMENT AGREEMENT (1984-89)
STREAM SEDIMENT AND WATER GEOCHEMICAL SURVEY
SOUTHERN YUKON TERRITORY, 1985

Elevation in feet above mean sea level
Mean magnetic declination 1986, 20°39' East,
decreasing 13.4' annually. Readings vary
from 29°29' E in the SE corner to 29°48' E in
the NW corner of the map area

Scale 1:250 000
Universal Transverse Mercator Projection
© Crown Copyrights reserved



NATIONAL TOPOGRAPHIC SYSTEM REFERENCE INDEX
TO ADDITIONAL GEOLOGICAL SURVEY OF CANADA MAPS

LEGEND	
TERTIARY	
21	LTG 62* Rhyolite porphyry, granite, granodiorite OLIGOCENE AND MIOCENE CARMACKS GROUP
20	OMCV 60 Andesite, basalt, breccia EOCENE MOUNT NANSEN GROUP
19	EMN 59 Acid to intermediate tuff, breccia LOWER(?) TERTIARY
18	TFP 58 Feldspar porphyry dykes and flows
17	TVA 58 Acid tuff
16	TVB 58 Andesite, porphyritic basalt flows and dykes EARLY TERTIARY
15	ETGA 57 Alaskite, granite, quartz monzonite
14	ETQM 57 Granite, quartz monzonite
13	FPPP 57 Feldspar porphyry dykes
JURASSIC AND CRETACEOUS	
12	JKT 51 TANTALUS: Conglomerate, siltstone, arkose, coal
11	JKK 51 KLUANE: Sericitic to biotitic schist, gneiss, amphibolite JURASSIC LABERGE GROUP
10	JL 47 Greywacke, arkose, conglomerate
TRIASSIC	
9	TV 42 Basaltic greenstone
8	TQM 42 Leucocratic, porphyritic quartz monzonite
7	TGD 42 RUBY RANGE: Granodiorite
6	TDGN 42 Foliated hornblende granodiorite, quartz
MESOZOIC UNDIVIDED	
5	MQM 41 Porphyritic quartz monzonite
4	MDI 41 Diorite
PALEOZOIC UNDIVIDED	
3	PM 09 Amphibolite, schist, gneiss
2	HCSN 08 Schist, gneiss, quartzite HADRYNIAN AND CAMBRIAN
PALEOZOIC HADRYNIAN	
1	HC 07 Crystalline limestone

*A mnemonic code assigned to rock types and recorded as part of field observations

Geological boundary
Fault
No analytical result

Geological base and legend are derived from: Map 1398A, MACMILLAN RIVER, YUKON - DISTRICT OF MACKENZIE - ALASKA, NTS SHEET 105, 115. Compiled by H. Gabriele, D.J. Templeman-Kluit, S.L. Blusson and R.B. Campbell, Geological Survey of Canada, Energy, Mines and Resources Canada, 1980. 1:1 000 000 scale