

SURFICIAL GEOLOGY

PROGLACIAL AND GLACIAL ENVIRONMENTS

- Glaciolacustrine Deposits:

 - Varved or massive clay and silt
 - Fine Sands, sands, and deltaic sand

- Glaciofluvial and Ice Contact Deposits:

 - Outwash sands and gravel
 - End moraine, interlobate moraine; sand, gravel and boulders
 - Esker or kame complex; sand, gravel, boulders

- Glacial Deposits:

 - Predominantly clayey till
 - Predominantly silty to sandy till

NONGLACIAL ENVIRONMENT

Bedrock

Complexes: when two or more types of glacial or non-glacial environment are interposed in a mosaic or repeating pattern, the relative dominant/subordinate amount of each type is indicated by sequential order. For example, 2/1 indicates predominantly silty to sandy till with lesser clayey till.

SYMBOLS

Surficial geological boundary

Striae

Fluting, drumlin or drumlinoid ridge

Esker, kame or kame complex

Surficial geological derived from:
Boissonneau, A.N. (1965), Map S465, Ontario Department of Lands, Forests.
Prest, V.K., Grant, D.R., and Rampton, V.N. (1969), Glacial Map of Canada, Geological Survey of Canada, Map 1253A (Scale: 1:5 000 000).

Geological Survey of Canada
Mineral Resources Division
Exploration Geochemistry Subdivision

CONTRACTORS

Sample collection by SIAL Geophysique Inc., Montreal
Sample preparation by Golder Associates, Ottawa

Sediment chemical analyses by Barringer Magenta Ltd., Rexdale, Ontario


Au analyses by Chemex Labs Limited, Vancouver
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

Contribution to Canada - Ontario Mineral Development Agreement 1986 - 1990, a subsidiary agreement under the Economic and Regional Development Agreement Project funded by the Geological Survey of Canada.

Ministry of Northern Development and Mines

Energy, Mines and Resources Canada Energie, Mines et Ressources Canada

Canada



HG

PPB

21000

MAX

255

98

220

95

190

90

136

70

11



MIN

1333 SAMPLES


REGIONAL TREND MAP

KILOMETERS - SCALE 1:1000000

The regional geochemical trend map displayed above utilized a moving weighted average using an inverse distance function ($1/d^3$) to filter out minor irregularities and emphasize large-scale regional features. Single point anomalies may be superimposed on estimates; however, geological units which are chemically enriched, or large metallic deposits undergoing weathering would be expected to produce identifiable anomalies.



CONCENTRATION	FREQUENCY
256 to 21000	N= 27(2.0%)
221 to 255	N= 39(2.9%)
191 to 220	N= 66(5.0%)
137 to 190	N= 256(19.2%)
11 to 136	N= 945(70.9%)



MERCURY (ppb) LAKE SEDIMENTS

GSC OPEN FILE 1357

REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 93-1986

CANADA - ONTARIO

MINERAL DEVELOPMENT AGREEMENT (1986-1990)

LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY

CENTRAL ONTARIO, 1986

Scale 1:250 000 - Échelle 1/250 000

Kilometres 0 5 10 15 20 Kilometres


Universal Transverse Mercator Projection

Projection transversale de Mercator

Crown Copyright reserved

Droits de la Couronne réservés

Base map at the same scale published by the Surveys and Mapping Branch in 1979



41-O

LEGEND

PRECAMBRIAN

LATE PRECAMBRIAN

- 11 LPAC* Carbonatite - alkalic complexes; alkalic syenite, pulaskite, brecciated and fenitized rocks, nepheline syenite, syenite, magnetite-patite rock, urtite, ijolite, metteigite
- 10 LPAD Diabase dykes
- 9 ASUB Shanner Anorthosite Complex; anorthosite to gabbro, gneissic to faser-textured tonalite and monzonite
- 8 AKN Kapuskasing structural zone rocks; meta-igneous rocks, melanocratic granulite, pelitic and psammitic granulites, metasedimentary gneiss and arkosic metasediments
- 7 AGM Massive felsic to intermediate plutonic rocks; granite, granodiorite, tonalite, quartz monzonite, monzodiorite, pegmatite
- 6 AGN Foliated to gneissic felsic to intermediate plutonic rocks; granite, granodiorite, tonalite, quartz monzonite, diorite, migmatite
- 5 AUB Mafic and ultramafic intrusive rocks, including gabbro, diorite and serpentinized ultramafics
- 4 ASGN Paragneiss, orthogneiss and migmatite
- 3 ACSP Metasediments; greywacke, arkose, quartzite, conglomerate, argillaceous and migmatized metasediments biotite-quartz-feldspar schist and gneiss
- 2 AMVF Felsic to intermediate metavolcanics; ryholite to dacite flows and fragments, tuff, lapilli-tuff, agglomerate, breccia, porphyry flows
- 1 AMVB Mafic to intermediate metavolcanics; basalt to andesite flows, porphyritic flows, and pillow lavas, mafic pyroclastics, layered amphibolite, diorite, gabbro, migmatized mafic metavolcanics

IF Iron formation

*A mnemonic code assigned to rock types and recorded as part of field observations.

Geological boundary:

Fault:

No analytical results

The geology base and legend for these geochemical maps were derived from: Geology - Chapleau - Foleyet, Map 2221 Geological Compilation Series, Ontario Department of Mines, 1:253 400. McFrank, J.F.U., and Brown, P.A. (1979), Geology - Plutonic Rocks in Ontario, Geological Survey of Canada, Map 1253A, to accompany GSC Paper 80-23.

Copies of map material and listings of field observations, analytical data and methods, from which the open file was prepared, are available from:

K.G. Campbell Corporation
880 Wellington St.
Bay 238
Ottawa, Ontario
K1R 6K7

Digital data are available on IBM-PC compatible diskette from:

Geological Survey of Canada
Publications Distribution
601 Booth St.
Ottawa, Ontario K1A 0E8
Tel: (613) 995-4342