

0 20 40 60
KILOMETERS - SCALE 1:1000000

SURFICIAL GEOLOGY

This legend is common to Open Files 1356 and 1357

PROGLACIAL AND GLACIAL ENVIRONMENTS

Glaciolacustrine Deposits:

- Varved or massive clay and silt
- Fine Sands, sands, and deltaic sand

Glactofluvial and Ice Contact Deposits:

- Outwash sands and gravel
- End moraine, interlobe moraine; sand, gravel and boulders
- Esker or kame complex; sand, gravel, boulders

Glacial Deposits:

- Predominantly clayey till
- Predominantly silty to sandy till

NONGLACIAL ENVIRONMENT

- Bedrock

Complexes: when two or more types of glacial or non-glacial environment are interspersed in a mosaic or repeating pattern, the relative dominant/subordinate amount of each type is indicated by sequential order. For example, 2/1 indicates predominantly silty to sandy till with lesser clayey till.

SYMBOLS

- Surficial geological boundary
- Striae
- Fluting, drumlin or drumlinoid ridge
- Esker, kame or kame complex

Surficial geology derived from: Boissonneau, A.N. (1965), Map S465, Ontario Department of Lands, Forests, and Parks; Press, V.K., Grant, D.R., and Rampton, V.N. (1969), Glacial Map of Canada, Geological Survey of Canada, Map 1253A (Scale: 1:5 000 000).

Geological Survey of Canada
Mineral Resources Division
Exploration Geochemistry Subdivision

CONTRACTORS

Sample collection by SIAL Geophysique Inc., Montreal
Sample preparation by Golder Associates, Ottawa



Sediment chemical analyses by Barringer Magenta Ltd., Rexdale, Ontario

Au analyses by Chemex Labs Limited, Vancouver
Water chemical analyses by Barringer Magenta Laboratories (Alberta) Ltd., Calgary

Contribution to Canada - Ontario Mineral Development Agreement 1986 - 1990, a subsidiary agreement under the Economic and Regional Development Agreement. Project funded by the Geological Survey of Canada.



Energy, Mines and Resources Canada / Énergie, Mines et Ressources Canada



Elevation in feet above mean sea level
Mean magnetic declination 1987, 8°20' West, increasing 10.9' annually. Readings vary from 9°07'W in the SE corner to 7°30'W in the NW corner of the map area

SAMPLE LOCATION
GSC OPEN FILE 1357
REGIONAL GEOCHEMICAL RECONNAISSANCE MAP 93-1986
CANADA - ONTARIO
MINERAL DEVELOPMENT AGREEMENT (1986-1990)
LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY
CENTRAL ONTARIO, 1986

Scale 1:250 000 - Échelle 1:250 000
Kilometres 0 10 20
Universal Transverse Mercator Projection / Projection transverse universelle de Mercator
© Crown Copyrights reserved / © Droits de la Couronne réservés

Base map at the same scale published by the Surveys and Mapping Branch in 1979

41-O

SAMPLE LOCATION
GSC OPEN FILE 1357
CENTRAL ONTARIO, 1986

LEGEND

- PRECAMBRIAN**
- LATE PRECAMBRIAN**
- 11 LPAC* Carbonatite - alkalic complexes; alkalic syenite, pulaskite, brecciated and fenitized rocks, nepheline syenite, sovite, magnetite-apatite rock, urtite, tsilite, melteigite
- EARLY TO MIDDLE PRECAMBRIAN**
- 10 LPAD Diabase dykes
- EARLY PRECAMBRIAN (ARCHEAN)**
- 9 ASUB Shawmere Anorthosite Complex; anorthosite to gabbro, gneissic to flaser-textured tonalite and monzonite
- 8 AKN Kapuskasing structural zone rocks; meta-igneous rocks, melanocratic granulite, pelitic and psammitic granulites, metasedimentary gneiss and arkosic metasediments
- 7 AGM Massive felsic to intermediate plutonic rocks; granite, granodiorite, tonalite, quartz monzonite, monzodiorite, pegmatite
- 6 AGN Foliated to gneissic felsic to intermediate plutonic rocks; granite, granodiorite, tonalite, quartz monzonite, diorite, migmatite
- 5 AUB Mafic and ultramafic intrusive rocks, including gabbro, diorite and serpentinized ultramafics
- 4 ASGN Paragneiss, orthogneiss and migmatite
- 3 ACPM Metasediments; greywacke, arkose, quartzite, conglomerate, argillaceous and migmatized metasediments, biotite-quartz-feldspar schist and gneiss
- 2 AWF Felsic to intermediate metavolcanics; rhyolite to dacite flows and fragmentals, tuff, lapilli-tuff, agglomerate, breccia, porphyritic flows
- 1 AMWB Mafic to intermediate metavolcanics; basalt to andesite flows, porphyritic flows, and pillow lavas, mafic pyroclastics, layered amphibolite, diorite, gabbro, migmatized mafic metavolcanics

IF Iron formation
*A mnemonic code assigned to rock types and recorded as part of field observations.

Geological boundary:
Fault:
No analytical results:

The geology base and legend for these geochemical maps were derived from: Geology - Chapleau - Foleyet, Map 2221 Geological Compilation Series, Ontario Department of Mines, 1:253 400. McCrank, G.F.D., Mistura, J.D., and Brown, Geology - Plutonic Rocks in Ontario, Geological Survey of Canada Map 1533A, to accompany GSC Paper 80-23.

Copies of map material and listings of field observations, analytical data and methods, from which the open file was prepared, are available from:

K.G. Campbell Corporation
880 Wellington St.
Bay 238
Ottawa, Ontario K1R 6K7

Digital data are available on IBM-PC compatible diskette from:

Geological Survey of Canada
Publications Distribution
601 Booth St.
Ottawa, Ontario K1A 0E8
Tel: (613)995-4342

This map has been reprinted from a scanned version of the original map. Reproduction par numérisation d'une carte sur papier.

SAMPLE LOCATION
GSC OPEN FILE 1357
CENTRAL ONTARIO, 1986
21 of 25