NEW GLASGOW - TONEY RIVER MAP: DESCRIPTIVE NOTES

INTRODUCTION:

The New Glasgow - Toney River map area lies in central northern mainland Nova Scotia. It includes parts of two basement blocks, the Antigonish and Cobequid Highlands, separated by the Cumberland Basin, and the Carboniferous Basin to the north (Fig. 1). Most faults in the area are high angle. Major features include the Cobequid and Hollow faults, which mark the northern and southern boundary of the Cobequid Highlands and the northwestern boundary of the Antigonish Highlands. Stellarton Graben, site of the Pictou Coalfield, is bounded by subsidiary, overlapping ends of these faults. Rocks in the area range in age from middle(?) Proterozoic to late Carboniferous.

This area was first mapped on a regional scale by Logan and others (1863, 1864) and Dawson (1868). Parts of the area have been remapped by various workers (Poole, 1954; Bell, 1946; Halter, 1956; Mawhi, 1961; Gillis, 1964; Donohoe and Wallace, 1978; Murphy, 1985; Murphy, 1986). This map is a compilation of previous mapping, with additional observations. All of the area is till-covered. The best exposures are found along streams, shorelines, and roads.

ANTIGONISH HIGHLANDS:

The oldest rocks in the Antigonish Highlands are assigned to the Georgeville Group, a late Proterozoic assemblage of mafic to felsic volcanic rocks and associated sediments. In the northern Antigonish Highlands, greenish grey, turbiditic, feldspathic wacke and slates of the James River Formation predominate (Murphy, 1985). These rocks are intruded by the Merigomish pluton. In the northern Antigonish Highlands the Georgeville Group is overlain by Cambro-Ordovician strata. An appinite dyke intruding the lower Georgeville Group has yielded a K-Ar hornblende age of 530 +/- 33 Ma (Wanless et al. 1967, corrected to 1978 IUGS standard).

The upper Ordovician to lower Devonian Arisaig Group (Boucot et al., 1974; Mawhi, 1961; Murphy, 1986) unconformably overlies the Georgeville Group. The basal unit is the Beart Brook Formation, consisting of subaerial, mafic to felsic flows, mafic to felsic volcaniclastics, and arkose. A whole rock Rb/Sr isochron of 430 +/- 12 Ma has been obtained from Beart Brook rhyolites (Donohoe and Wallace, 1978; recalculated using IUGS standards by Keppie et al, 1987). The Beaubush Cove Formation, comprising reddish arkose and greenish, fossiliferous shale and sandstone, overlies the Beart Brook volcanics disconformably, and is overlain conformably by dark grey to black fossiliferous shales and limestones of the Ross Brook Formation. These rocks, in which strata were deposited during Llandovelian time, Greenish grey and dark grey, micaceous, shallow marine siltstone and sandstone of the lower Ludlovian McAdam Formation overlie the Ross Brook Formation disconformably. Wenlockian strata are missing from the Antigonish Highlands sequence. Pridolian to lower Devonian Handsomans and siltstones, interbedded red and green siltstone and fluvialite, red, calcareous siltstone and marl of the Pleasant Bay Formation overlie the McAdam conformably. Blue-grey to green-grey, micaceous, cross-laminated, shallow marine siltstone and massive sandstone of the Pridolian to lower Bonnehouse Formation and overlie McAdam and Ross Brook strata.

The Arisaig Group is conformably overlain by red, fluvialite siltstones, sandstones, and minor conglomerate of the Devonian to lower Siegenian Knoydart Formation. The Knoydart may be correlated with the Old Red Sandstone of western Europe (Boucot et al, 1974).

Green, subaerial basalts, reddish felsic tuffs, and oligomictic, alluvial fan conglomerates of the Devono-Carboniferous Fountain Lake Group (Donohoe and Wallace, 1982; Donal et al, 1983) overlie pre-Devonian strata in the Beaubush Mountain - McLellan Brook area. These rocks are correlative with the Fountain Lake Group in the Cobequid Highlands. Red knoydart clasts predominate in the conglomerates.

COBEQUID HIGHLANDS:

The oldest rocks in the Cobequid Highlands are 934 +/- 82 Ma (whole rock Rb/Sr) granitic orthogneisses of the Mt. Thom Complex (Donohoe, 1976; Wallace and Donohoe, 1977; Gaudette et al. 1984). Relationships between the Mt. Thom Complex and younger rocks are masked by several generations of felsic plutons described below.

Rhyolites, dacites, and minor andesites, intruded by late Paleozoic plutons, are believed to pass westward into shallow marine rocks of the Georgeville Group in the Antigonish Highlands (Wallace and Donohoe, 1977), and are thus correlative with the Arisaig Group. Recent investigations, however, suggest that these volcanics are Precambrian (R. Murphy, personal communication, 1987).

Felsic and mafic terrestrial volcanics, and polymictic orthoconglomerates, wackes, and siltstones of the Diamond Brook Formation (Donohoe and Wallace, 1982) overlie Mt. Thom rocks unconformably on Pictou strata west of the map area. The conglomerates may be distinguished from the overlying Falls and Millsville conglomerates by the scarcity of igneous clasts (Fralick, 1985). Spore assemblages (Donohoe and Wallace, 1982), and the presence of Diamond Brook clasts in the basal conglomerate of the Fountain Lake Group (Cormier, 1983), indicate deposition in Emsian-Tournaisian time.

The late Devonian(?) to early Carboniferous Falls Formation, a thick wedge of polymictic orthoconglomerate with minor sandstone and paraconglomerate, overlies the Fountain Lake Group with angular unconformity (Wallace and Donohoe, 1977). The Falls Formation is the oldest unit preserved in the Stellarton Graben. Felsic fluvial deposits of the Tournaisian Nuttby Formation, both formations can be assigned to the Horton Group.

A plutonic complex forms the core of the western Cobequid Highlands. The largest intrusion is the Salmon River Suzolite-hornblende granite. A Visean age is probable since it is nonconformably overlain by Riversdale strata and is intrusive into Canso Group strata (Donohoe and Wallace, 1978). Several other plutons, age relationships of the latter, which include the Glen Road granite, the Eight Mile Brook syenite, and the Brookland and Mount Thom diorites, are uncertain.

STELLARTON GAP:

The Windsor Group is exposed at the eastern end of the Antigonish Highlands, where it lies unconformably on older strata (Giles, 1982), and in fault blocks along the eastern part of the Cobequid fault system (Belt, 1968; Giles, 1979). Gypsum, anhydrite, and carbonates of the early middle Visean Bridgeville Formation (Bell, 1946) overlie the Windsor Group. Fluvial, deltaic, evaporitic and local conglomerates of the Forbes Lake Formation, and sandstone, siltstone, and carbonates of the mid-Visean to earliest Namurian Churchville Formation. Volcanic clasts predominate in the Forbes Lake conglomerate.

Lower Carboniferous strata are overlain by four diachronous cycles of middle to late Carboniferous alluvial fan and fluvial sediments: the Canso, Riversdale, Cumberland, and Pictou Groups.

The late Visean to early Namurian Canso Group rests conformably on the upper Windsor Group in the southern part of the map area. It comprises red and grey fluvial sandstones and conglomerates. Canso Group strata are folded in the southern part of the map-area, and along fault zones. Canso sandstones are phyllitic. The Canso Group also includes sandstones of the Lismore Formation and polymictic orthoconglomerates of the Hollow Formation deposited north of the Hollow fault (Fralick and Schenk, 1981). Lismore strata may be distinguished from Canso strata by the abundance of greenish quartzite clasts in the former (Fralick and Schenk, 1981). Late Visean foraminifera from a limestone bed in the Lismore Formation (Mamet, 1970) suggest a transitional relationship between the Windsor and Canso Groups (Fralick and Schenk, 1981).

The early middle Carboniferous Riversdale Group is in fault contact with older Carboniferous strata along the eastern and western end of the Cobequid Highlands. A lower unit, the Millsville Conglomerate, contains predominantly reddish-brown, polymictic orthoconglomerate with minor sandstone and siltstone. The presence of white quartzite and felsic plutonic clasts (Fralick, 1985) indicates a source in the Cobequid Highlands to the south. This unit is transitional to the overlying Boss Point Formation, a sequence of grey and reddish-brown lithic sandstone and conglomerate with minor siltstone and coal (Fralick, 1981). Red and green siltstone of the eastern Cobequid Highlands and the Boss Point Formation are assigned to the Riversdale Group (Donohoe, 1982). The Riversdale Group was shed eastward into Stellarton Gap from the Cobequid Highlands.

The early Upper Carboniferous Cumberland Group includes two formations in Stellarton Gap. Predominantly red, fluvial sandstone, mudstone, and minor conglomerate of the middle Westphalian B Middle River Formation rests disconformably on Riversdale strata in the eastern part of the gap, and lies conformably or disconformably on Boss Point strata to the north. The Westphalian B to early Westphalian C New Glasgow Conglomerate is a fluvial sandstone and conglomerate with minor sandstone and siltstone. It conformably overlies downwards with the Middle River Formation. Contacts with older strata (e.g.: south of Fraser's Mountain) are not exposed. The New Glasgow Conglomerate can be distinguished from older conglomerates by the presence of Canso and Windsor clasts, especially carbonates.

The upper Carboniferous Pictou Group overlies the New Glasgow Conglomerate disconformably or conformably. Fluvialite redbeds are typical of the Pictou Group throughout the Maritimes, but in Stellarton Gap greybeds predominate. These strata, which comprise the Coal Measures, consist of cyclic, fining upward channel sandstone beds alternating with thick grey or red mudstones, and minor thin coals and oil shales. A thin stromatolitic limestone lies at the base of the Formation. To the west, the Merigomish Formation is overlain by predominantly red sandstone of the Pictou Landing Formation. The Merigomish Formation was deposited in late Westphalian C to Westphalian D time.

Partly coeval with the Merigomish Formation, but separated from it by an easterly-trending ridge of Cumberland and Canso Group rocks, is the Stellarton Formation. This comprises oil shale- and coal-bearing, grey and red, lacustrine, and alluvial fan sediments. Palynological data from the Stellarton Formation indicate an early Westphalian C to early Westphalian D age. The Stellarton Formation is entirely restricted to the Stellarton Graben.

SURFICIAL GEOLOGY

Till blankets most of the map area. Till thickness is variable, up to 25 m or more locally. Three tills have been recognized: a red clay till derived from the north; a hybrid till produced by reworking and mixing of the lower till with material from the south, and a local stony till produced by limited ice movement in the Cobequid Highlands during the last glaciation (N. Rea, personal communication, 1987). Most of the ice contact features such as kames and eskers are restricted to the Cobequid and Antigonish Highlands.

STRUCTURAL GEOLOGY

The Mt. Thom Complex has undergone at least three major deformations, thought to be early Hadrynian or older (Donohoe and Wallace, 1985). Early isoclinal folds with axial plane foliation are refolded by open, upright folds, on locally steep dipping incipient, axial planar, crenulation cleavage.

Georgeville Group strata in the western Cobequid Highlands have undergone folding about variable axes. Steep dips are due to rotation of fault blocks. Folding predates cleavage development (Murphy, 1986).

Major regional deformation took place during the middle Devonian. Arisaig Group and Carloont Formation strata show open folding about NE axes (Boucot et al, 1974; Donohoe and Wallace, 1977; Murphy, 1986). A regional cleavage, best developed in fine-grained strata, was imposed on early Devonian and older rocks at this time. In the Cobequid Highlands slaty cleavage is only developed locally. In the western Antigonish Highlands cleavage dips steeply southeast. It is cut by the Hollow and Coronation Faults, but dips westerly to the west of it (Murphy, 1986). Reverse movement on the Hollow fault (Boucot et al, 1974) and the Coronation fault (Murphy, 1986) in the western Antigonish Highlands (Murphy, 1986) indicate regional SE-NW compression. The end of this deformation is marked by normal movement on the Hollow Fault (Boucot et al, 1974).

Renewed SE-NW compression in the late Carboniferous resulted in gentle to open folding of older Carboniferous strata about NE axes, and dextral movement on the Hollow and Cobequid Faults. This led to formation of the Stellarton Graben (Yeo and Gao, 1986 Gao). The length of the graben suggests that late Westphalian dextral movement on the Cobequid fault was significant; minor open folding about NE axes and second-order faulting developed. Near the major faults penetrative fabrics are developed in middle Carboniferous strata.

Northerly-trending Mesozoic normal faults cross-cut all earlier structures.

GEOLOGICAL HISTORY

The Mt. Thom Complex records a middle Proterozoic intrusive event, as well as the Grenville orogeny, when Cobequid rocks were metamorphosed to upper greenschist facies. Andesite-dacite-rhyolite (late Proterozoic, cratonic rifting is recorded by the volcanics and marine sediments of the Georgeville Group (Murphy, 1985).

The absence of lower Paleozoic strata in the map area suggests that this was an area of emergence at that time. Middle Paleozoic strata of the Arisaig Group lap onto the Antigonish Highlands from the southeast. Strata of the Arisaig Group were deposited under terrestrial conditions. Volcanism was followed by lower Llandoverian marine transgression (Boucot et al, 1974). Lower Llandovery Ross Brook shallow marine clastics were derived from eroded Middle Arisaig strata to the southeast. The absence of Wenlockian strata within the map area, a facies change from west to the east, suggests local uplift. Shallow marine conditions returned again until late Ludlovian time, when deposition of the McAdam Formation took place. Deposition of strata in the Pridolian - Emsian Bonnehouse Formation (Boucot et al, 1974), derived from a new source to the southeast, possibly the Meguma Terrane, was followed by regression and deposition of coastal floodplain redbeds in the Siegenian (Boucot et al, 1974). Middle Devonian deformation is probably related to the collision of the Meguma Terrane with eastern North America and the final closure of the Iapetus Ocean.

Major regional deformation took place in the middle Devonian, as noted above. Terrestrial volcanism, accompanied by shedding of coarse-grained molasse from the Cobequid and Antigonish Highlands is recorded in the Fountain Lake Group and its correlatives (Donohoe and Wallace, 1982). Volcanism and sedimentation continued intermittently at least into the late Devonian. Cobequid and Hollow Faults were active by this time. Alluvial fan sedimentation persisted into the early Carboniferous.

A series of extensive, marine transgression-regression cycles are recorded by deposition of carbonates, evaporites, and clastics of the Windsor Group in Visean to early Namurian time. The end of Windsor deposition is marked by fine-grained terrestrial clastics of the Canso Group. Renewed uplift of the Cobequid Highlands is signalled by development of alluvial fan and fluvial plains of the Riversdale Group, shed eastward into Stellarton Gap during late Namurian - early Westphalian time.

Renewed activity on the Cobequid-Hollow fault system in the middle Westphalian is indicated by alluvial fan deposits of the Cumberland Group, which prograded northerly into Stellarton Gap over fine-grained fluvial deposits shed from the highlands.

Dextral movement on the faults in late Westphalian time led to formation of the Stellarton Graben. Lacustrine and fluvio-deltaic deposits of the Stellarton Formation, including coals and oil shales, accumulated. Deposition of the Pictou Group began at about the same time from northerly flowing fluvial systems. Fine-grained, lacustrine and overbank sedimentation persisted regionally until the Permian, long after subsidence and deposition within Stellarton Graben had ceased.

At least four phases of Wisconsinan glaciation may be recognized in the area (Grant, 1977; Rampton et al, 1984; Rea and Finck, 1984; Rivers and Stea, 1988). About 38,000 years ago (Grant, 1977), following the Sangamonian (interglacial) stage, a major ice lobe moved over the area, moving towards the east and southeast. During a period of retreat, the ice advanced again, this time moving towards the west and southwest. In another interstadial period, ice again moved into the area, but this time from the south. In the final phase, ice flowed northwards from the Cobequid and Antigonish Highlands, across Stellarton Gap and around the Cobequid Highlands. Glaciation ended about 14,000 years ago (Grant, 1977).

ECONOMIC GEOLOGY

The Pictou Coalfield in Stellarton Graben is one of the major coal fields in Nova Scotia (Dickie, 1988; Hacquebard and Donaldson, 1969). Production from 1918 to 1984 was 40 million tonnes. Small-scale production still continues near Westville. Estimates of reserves remaining in the Stellarton Formation range from 64 (Chesal, 1982) to 69 million tonnes (Hacquebard, 1979) of which about 30 million tonnes could be recovered (Dickie, 1988). Preliminary mining feasibility studies by Suncor Inc. indicate that about 27 million tonnes could be recovered from the Foord seam east of the old workings, with further potential. Ash content is relatively high, but sulphur is low, making these coals suitable for use in power plants (Hacquebard, 1979). Extensive oil shales are also found in the Stellarton Formation (Macauley and Ball, 1984; Naylor and Smith, 1986; Kalkreuth and Macauley, in press). Hydrocarbon yields are generally too low for them to be of economic interest, however (Naylor and Smith, 1986).

A number of small copper occurences and one zinc showing are found in the Cobequid Highlands. Copper mineralization is locally associated with coaly zones, presumably redox traps (Dunsmore, 1977), possibly localized above faults, while sphalerite is found in a small pyritic coal seam on the Toney River. Galena is found locally on fracture surfaces in the Stellarton Formation. None of the occurences have known economic potential.

The Windsor Group appears to have the most potential of Carboniferous strata to host massive sulphides (Kirkham, 1985). Disseminated chalcopyrite is found in Windsor gypsum, near Lismore (Giles, 1979; Kirkham, 1985). Chalcopyrite and sphalerite also occur in Windsor limestone near Churchville (Kirkham, 1985).

Several small iron deposits are found in the northwestern Cobequid Highlands. The only one of note within the map area is an oolitic hematite bed in Beechhill Cove sandstones near Falls Brook (Wright, 1975).

Willemite and native silver are found in calcite veins in Georgeville Group metasedimentsnear Kirshmount. These veins are thought to be related to faults (Rangsiter, 1986).

Crushed and dimension stone has been taken from several quarries in the Merigomish and Boss Point formations and from Eight Mile Brook syenite and Salmon River granite. Quarries in Metalarna Mountain are currently operating. Attractive building stone can be found at Eight Mile Brook and at Sawmill Brook (Dickie, 1988; in preparation). Occurences of fine-grained Windsor Group limestone, potentially suitable for quarrying, have been found at Lismore, Fox River, Churchville, Brookfield, Stony Brook, and Limerock. Shale from the Stellarton Formation is quarried south of New Glasgow for tile and brick-making.

Gravel for road construction is extracted from several deposits in the western Cobequid Highlands (Fowler and Dickie, 1977).

REFERENCES

Bell, W.A., 1946. The Pictou Coalfield, Nova Scotia. Geological Survey of Canada Memoir 235, 161p.

Boucot, A.J., Dewey, J.F., Dineley, D.L., Fletcher, L., Fyson, W.K., Griffin, J.G., Hickox, C.F., McKerrow, W.S., and Ziegler, W., 1974. Geology and Paleontology of the Arisaig Area, Nova Scotia. Geological Society of America Special Paper 139, 135 p.

Chandler, F.W., Sullivan, R.W., and Currie, K.L., 1987. The age of the Springdale Group, western Newfoundland, and correlative rocks in Cape Breton Island: implications for stratigraphy of the Canadian Appalachians. Transactions of the Royal Society of Edinburgh: Earth Sciences (in press).

Chesal, S., 1982. Pictou Coalfield Project. An Evaluation of Coal Resources in the Westville, Stellarton, and Thorburn Districts of the Pictou Coalfield. Nova Scotia Department of Mines and Energy internal report, 92p.

Cormier, R.F., 1982. Rb/Sr age data for the Fountain Lake Group volcanics, Nova Scotia Department of Mines and Energy Report 82-1, p. 314-315.

Dickie, G.R., 1988. Building stone project. Nova Scotia Department of Mines and Energy Report 88-1, p. 45-49.

Dickie, G., (in preparation). Sandstones in Nova Scotia. Nova Scotia Department of Mines and Energy Bulletin.

Donohoe, H.V., 1976. The Cobequid Mountains Project. Nova Scotia Department of Mines and Energy Report 76-2, p. 113-126.

Donohoe, H.V. and Wallace, P.I., 1982. Debert-Mt. Thom region, Nova Scotia Department of Mines and Energy Map 82-9 (1:50,000 scale).

Donohoe, H.V. and Wallace, P.I., 1985. Repeated orogeny, faulting, and stratigraphy in the Cobequid Highlands, Avalon Terrain of Northern Nova Scotia. GAC Annual Meeting (Fredericton), Guidebook Excursion 7, 77p.

Dostal, J., Keppie, J.D., and Dupuy, C., 1981. Petrology and geochemistry of Devono-Carboniferous volcanic rocks in Nova Scotia. Maritime Sediments, v. 19, p. 59-71.

Dunsmore, H.E., 1977. A new genetic model for uranium-copper mineralization, Permo-Carboniferous Basin, northern Nova Scotia. Geological Survey of Canada Paper 77-1B, p. 247-251.

Fletcher, W., 1902. Toney River map sheet. Geological Survey of Canada Map 668 (1:63,360).

Fowler, J.H. and Dickie, G.R., 1977. Sand and Gravel Resources of Pictou County, Nova Scotia Department of Mines and Energy Paper 77-6, 42p.

Fralick, P.W., 1980. Tectonic and Sedimentological Development of the Late Paleozoic Eastern Cumberland Basin, Maritimes Basin, Maritime Canada. MSc thesis, Dalhousie University, 166p.

Fralick, P.W. and Schenk, P.E., 1981. Molasse deposition and basin evolution in a strike-slip setting: the late Paleozoic eastern Cumberland Basin, Maritime Canada. In Sedimentation in Tectonics in Alluvial Basins. Edited by A.D. Miall. Geological Association of Canada Special Paper 23, p. 77-98.

Gao, M., 1987. Deformation Characteristics of the Eastern Cobequid and Hollow Fault Zones and Stellarton Basin, Nova Scotia. MSc thesis, University of New Brunswick, 227p.

Gaudette, W.J., Olszewski, W.J., and Donohoe, H.V., 1984. Rb/Sr isochrons of Precambrian age from plutonic rocks in the Cobequid Highlands, Nova Scotia. Nova Scotia Department of Mines and Energy Report 84-1, p. 295-302.

Giles, P.S., 1979. Upper Windsor limestone at Limerock, Pictou County, Nova Scotia Department of Mines and Energy Report 79-1, p. 73-80.

Giles, P.S., 1981. Major transgressive-regressive cycles in middle to late Visean rocks of Nova Scotia. Nova Scotia Department of Mines and Energy Report 81-3, 43p.

Giles, P.S., 1982. Geological map of the Eureka area, central Nova Scotia. Nova Scotia Department of Mines and Energy (1:50,000 scale).

Gillis, J.W., 1964. Geology of the northwestern Pictou County, Nova Scotia, Canada. Unpublished PhD thesis, Pennsylvania State University, 133 p.

Grant, D.R., 1977. Glacial style and ice limits, the Quaternary stratigraphic record, and changes of land and ocean level in the Atlantic Provinces, Canada. Geographie physique et Quaternaire, v. 31, p. 247-260.

Hacquebard, P.A., 1979. A geological appraisal of the coal resources of Nova Scotia. Canadian Institute of Mining and Metallurgy Bulletin, v. 72, p. 76-87.

Hacquebard, P.A. and Donaldson, J.R., 1969. Carboniferous coal deposits associated with flood plain and limic environments in Nova Scotia. Geological Society of America, Special Paper 114, P, 143-191.

Halter, T.R., 1956. Some geological aspects of the Pictou Coalfield and their economic influence on mining operations. 3rd Third Conference on the Origin and Constitution of Coal, Crystal Cliffs, Nova Scotia, 1956. Nova Scotia Department of Mines and Nova Scotia Research Foundation, Halifax, p. 39-63.

Kalkreuth, W. and Macauley, G., in press. Organic petrology and geochemical (Rock-Eval) studies on Oil shales and coals from the Pictou and Antigonish areas, Nova Scotia, Canada. Canadian Society of Petroleum Geologists Bulletin.

Kirkham, R.V., 1985. Base metals in upper Windsor (Codroy) Group pollitic and stromatolitic limestones in the Atlantic Provinces. Geological Survey of Canada Paper 85-1A, p. 573-585.

Logan, W., Harvey, E., Fletcher, H., Fairbank, E.R., and Robert, 1902. Stellarton, New Glasgow, Pictou, and Westville map sheets, Geological Survey of Canada Maps 598, 600, 609, and 615 (1:63360).

Macauley, G. and Ball, F.D., 1984. Oil Shales of the Big Marsh and Pictou areas, Nova Scotia. Geological Survey of Canada Open File Report 1037, 57p.

Mamet, B.R., 1961. The Older Paleozoic of Pictou County, Nova Scotia. Nova Scotia Department of Mines and Energy Memoir 4, 112p.

Mamet, B.L., 1970. Carbonate microfacies of the Windsor Group (Carboniferous), Nova Scotia and New Brunswick. Geological Survey of Canada Paper 70-21, 121p.

Murphy, J.B., 1986. Geology of the western Antigonish Highlands, Nova Scotia. Geological Survey of Canada Paper 85-1A, p. 445-453.

Murphy, J.B., 1985. Geology of the southwestern Antigonish Highlands, Nova Scotia. Geological Survey of Canada Paper 85-1B, p. 519-530.

Myers, B.A. and Stea, R.R., 1986. Surficial mapping results: Pictou, Guysborough, and Antigonish Counties, Nova Scotia. Nova Scotia Department of Mines and Energy Report 86-1, p. 189-193.

Naylor R.D. and Smith, W.D., 1986. Stratigraphy, Lithology, and Composition of the Stellarton Group Oil Shales, Pictou County, Nova Scotia. Nova Scotia Department of Mines and Energy Open File Report 86-01.

Poole, H.S., 1954. Geological map of the Pictou coalfield. Geological Survey of Canada map 833 (1:30,375).

Stea, R.R., 1984. The sequence of glacier movements in northern mainland Nova Scotia determined through mapping and till provenance studies. In Correlation of Quaternary Chronologies. Edited by W.C. Mahaney, Geo Books, p. 279-291.

Stea, R.R. and Fowler, J.H., 1986. Patterns of glacier movement in Cumberland, Colchester, Hants and Pictou Counties, Northern Nova Scotia. Geological Survey of Canada Paper 84-1A, p. 677-684.

Wallace, P.I. and Donohoe, H.V., 1977. The Cobequid Highlands Survey. Nova Scotia Department of Mines and Energy Report 77-1, p. 187-188.

Wanless, R.K., Stevens, R.D., Lachance, G.R., and Edmonds, C.M., 1967. Age Determinations and Geological Studies, K-Ar Isotopic Ages. Report 8. Geological Survey of Canada Paper 67-2, p. 126.

Wright, J.D., 1975. Iron Deposits of Nova Scotia. Nova Scotia Department of Mines and Energy Economic Geology Series 75-1, p. 108.

Yeo, G.M. and Gao, R., 1986. Late Carboniferous dextral movement on the Cobequid - Hollow fault system, Nova Scotia: evidence and implications. Geological Survey of Canada Paper 86-1A, p. 399-410.

Contribution to Canada-Nova Scotia Mineral Development Agreement 1984-89, a subsidiary agreement under the Economic and Regional Development Agreement. Project funded by the Geological Survey of Canada.

Department of Mines and Energy

Energy, Mines and Resources Canada | Énergie, Mines et Ressources Canada

Canada


PICTOU MUN - PICTOU COUNTY
NOVA SCOTIA
CARIBOU CHANNEL
CARIBOU ISLAND
CARIBOU HARBOUR
NORTHUMBERLAND STRAIT
(GULF OF ST LAWRENCE)
PICTOU ISLAND
PICTOU HARBOUR
PICTOU ROAD
NEW GLASGOW
COBEQUID MOUNTAINS
Toney River
Caribou Harbour
Pictou
Boat Harbour
Little Harbour
Scotsburn
Westville
Stellarton
Trenton
Abercrombie
Millsville
Hardwood Hill
Rogers Hill
Durham
Glengarry
Glen Road
Mount William
Mount Thom
Brookland
Sutherlands River
Green Hill
Lismore
Fox River
Churchville
Brookfield
Stony Brook
Limerock
Salmon River
Eight Mile Brook
Sawmill Brook
Falls Brook
Beaubush Mountain
McLellan Brook
Metalarna Mountain
Fraser's Mountain
Dalhousie Mountain
Caribou Mountain
Mount Thom
Mount William