

Note: This legend is common for National Geochemical Reconnaissance Map 37-1978, Open File 557; Map 38-1978, Open File 558; Map 39-1978, Open File 559; Map 40-1978, Open File 560.

SEDIMENTARY, VOLCANIC AND METAMORPHIC ROCKS

TRIASSIC
[26] (VCRK) Andesitic volcanics and shallow intrusives of the Mistastin Formation

HADRYDIAN AND/OR NEOHELIXIAN

[25] (ARKS) Red conglomerate, arkose and siltstone

GRENVILLE PROVINCE

APHEBIAN AND EARLIER (?)

[24] (GRNG) Metasedimentary granitoid gneisses, minor amphibolite, sillinitic gneiss, metaquartzite, marble

SUPERIOR PROVINCE

ARCHEAN

[23] (PXGL) Pyroxene granulite, unseparated acidic intrusives

CHURCHILL PROVINCE

HELIKIAN

NEOHELIXIAN

[22] (SMRX) Quartzite, conglomerate, arkose, shale-Shipiskan Formation

PALEOHELIXIAN

[21] (QRTZ) Quartzite, grit and conglomerate of Sims Formation

APHEBIAN

[20] (BSLT) Basaltic flows and pyroclastics, quartzite, greywacke, slate, argillites, conglomerate, minor iron formation

[19] (SMRX) Grit, arkose, conglomerate, quartzite, greywacke, slate, acidic to basic volcanics, dolomite, limestone, chert breccia

[18] (SLTE) Ferruginous slate and iron formation

APHEBIAN AND EARLIER (?)

[17] (GRNL) Granulite, pyroxene gneiss, charnockite, minor granitic gneiss, mylonitic gneiss, amphibolite, ultrabasic intrusions

[16] (GRSS) Garnet-quartz-feldspar gneiss, chiefly mylonitized, locally graphitic

[15] (GRNG) Granitic gneiss, granodioritic gneiss, migmatite, agmatite, amphibolite

[14] (PRGS) Paragneiss; includes biotite-quartz-feldspar gneiss, garnet-biotite-quartz-feldspar gneiss, hornblende-gneiss, augen and graphitic gneiss

[13] (AMPB) Amphibolite, pyroxene amphibolite, chlorite schist, garnet- and biotite-rich gneisses

[12] (MSOM) Metasedimentary rocks, mainly quartzite and marble

NAIN PROVINCE

APHEBIAN

[11] (SLTE) Slate, argillite, siltstone, quartzite, greywacke, dolomite and basalt of LOWER CROTEAU GROUP

ARCHEAN

[10] (SCST) Mafic schistose rocks, greenstone, metasedimentary rocks, amphibolite, minor ultrabasic intrusions

[9] (GRGG) Granitic and granodiorite gneiss, migmatite, granulite and amphibolite

INTRUSIVE ROCKS

HELIKIAN

PALEOHELIXIAN

[8] (GRNT) Granite, quartz monzonite, granodiorite, quartz diorite, syenite

[7] (QZMZ) Adamellite suite: adamellite, monzonite, syenite, granodiorite, granite and their hypersthene-bearing equivalents forsunidite, mangerite, opdalite and charnockite

[6] (ANRS) Anorthosite suite: anorthosite, anorthositic gabbro, leucotroctolite

[5] (UMFC) Gabbro, norite, anorthositic gabbro, troctolite, diorite, derived basic gneiss and amphibolite

PALEOHELIXIAN AND EARLIER (?)

[4] (GRGR) Granitic to granodiorite, massive to poorly foliated, with inclusions of granitic gneiss

[3] (GRNT) Granite, quartz monzonite, granodiorite, quartz diorite

[2] (GBBR) Gabbro, metagabbro, glomerophytic gabbro and diorite

ARCHEAN

[1] (PXGD) Massive to poorly foliated pyroxene-bearing granodiorite and syenodiorite.

*A four letter mnemonic name recorded as rock type as part of field observations.

Geological boundary.....

Fault.....

Mainly acidic volcanic rocks.....

Mainly basic volcanic rocks.....

No analytical result.....


This legend was modified and the geology derived for this geochemical map from Geology Map of Labrador, Mineral Resources Division, Department of Mines, Agriculture and Resources, Province of Newfoundland and Labrador.

Geological Survey of Canada
Resource Geophysics and Geochemistry Division
and
Newfoundland Department of Mines and Energy

CONTRACTORS
Sample collection by Marshall Macklin Monaghan Ltd.
Sample preparation by Golder Associates.
Uranium in sediment chemical analyses by Atomic Energy of Canada Ltd.
Other sediment chemical analyses by Chemex Labs Ltd.
Water chemical analyses by Barringer Magenta Ltd.

This map forms one of a series of 68 maps released by the Geological Survey of Canada, Open Files 557, 558, 559 and 560. Each Open File consists of maps for 12 elements for lake sediments, 2 elements for lake water, and 1 each for sample site location, sediment loss on ignition and water pH.

pH in water
OPEN FILE 559
CENTRAL LABRADOR 1978



Elevation in feet above mean sea level

Mean magnetic declination 1978, 32°53.7' West, decreasing 10.2' annually. Readings vary from 32°28.8' in the SE corner to 33°21.6' in the NW corner of the map

pH in water
OPEN FILE 559


NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 39-1978

URANIUM RECONNAISSANCE PROGRAM


LAKE SEDIMENT AND WATER GEOCHEMICAL SURVEY

CENTRAL LABRADOR 1978

Scale 1:250,000



Base-map at the same scale published by the Mapping and Charting Establishment, Department of National Defence, 1968



This map has been reprinted from a scanned version of the original map. Reproduction par numérisation d'une carte sur papier.


pH in water
OPEN FILE 559
CENTRAL LABRADOR 1978

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
880 Wellington St.,
Bay 238
Ottawa, Ontario
K1R 6K7

The data is also available in digital form. For further information please contact:

The Director
Computer Science Centre
Department of Energy, Mines and Resources
Ottawa, Ontario
K1A 0E4



INDEX MAP