



Note: This legend is common for National Geochemical Reconnaissance Map 54-1982, Open File 901; Map 55-1982, Open File 902; Map 56-1982, Open File 903; Map 57-1982, Open File 904.

- SEDIMENTARY, VOLCANIC AND METAMORPHIC ROCKS
- HADRYANIAN AND/OR NEOHELIXIAN
- 34 [HDHL]*, (ARKS)** Red conglomerate, arkose, and siltstone.
- GRENVILLE PROVINCE
- HELIXIAN AND EARLIER (?)
- 33 [HUGP] Paragneisses, granitoid gneisses of probable sedimentary origin, minor quartzite and marble.
- APHEBIAN
- 32 [APHG] Dolomite, marble, quartzite, iron formation, parashist and paragneiss.
 - 31 [HUGN] Sillimanite gneiss, commonly migmatitic. Minor amphibolite.
 - 30 [HUGG] Granitic gneiss, mainly pink quartzo-feldspathic gneisses, commonly banded and migmatitic.
 - 29 [AGWF] Iron formation: WABUSH FORMATION
 - 28 [HUGB] Intermediate to basic gneiss, amphibolite.
 - 27 [HUGA] Garnetiferous gabbroic gneiss.
- APHEBIAN AND EARLIER (?)
- 26 [AUGP], (GRNG) Metasedimentary granitoid gneisses, minor amphibolite, sillimanite gneiss, metaquartzite, marble.
 - 25 [AUGN] Sillimanite gneiss, with abundant intrusive pegmatitic material, minor amphibolite.
 - 24 [AUGR] Granulitic, dioritic to granodioritic gneiss.
 - 23 [AUGB] Amphibolite, pyroxene amphibolite, chlorite schist, garnet- and biotite-rich gneisses.
 - 22 [AUGS] Marble.
- ARCHEAN
- 21 [ARCG] Granitic gneiss, amphibolite, unseparated massive acidic intrusives.
- SUPERIOR PROVINCE
- ARCHEAN
- 20 [ARCS], (PXGL) Pyroxene granulite, unseparated acidic intrusives: ASHUANIPI COMPLEX.
- CHURCHILL PROVINCE
- HELIXIAN
- NEOHELIXIAN
- 19 [NHWS] Quartzite, conglomerate, arkose, shale.
- PALEOHELIXIAN
- 18 [PHWS], (QRTZ) Quartzite, grit and conglomerate of SIMS FORMATION.
- PALEOHELIXIAN AND/OR APHEBIAN
- 17 [PHAW] Greywacke, quartzite, arkose, slate, phyllite, basic to intermediate volcanic rocks, derived schists and gneisses.
- APHEBIAN
- 16 [VAM2], (BSLT) Basaltic flows and pyroclastics, quartzite, greywacke, slate. (basic volcanic rocks).
 - 15 [APW1], [VAPW], [UAPW], (SMRK) Grit, arkose, conglomerate, quartzite, greywacke, slate, acidic to basic volcanics.
 - 14 [AW1S], (SLTE) Ferruginous slate and iron formation RUTH and COKOMAN FORMATIONS OF KNOB LAKE GROUP.
- APHEBIAN AND EARLIER (?)
- 13 [AUMR], (GRNL) Granulite, pyroxene gneiss, charnockite; minor granitic gneiss, mylonitic gneiss.
 - 12 [AUNG], (GRNG) Granitic gneiss, granodioritic gneiss, migmatite, agmatite, amphibolite.
 - 11 [AUMB], (AMPB) Amphibolite, pyroxene amphibolite, chlorite schist, garnet- and biotite-rich gneiss.

- INTRUSIVE ROCKS
- CAMBRIAN AND EARLIER
- 10 [CM18] Diabase dykes, radiometric ages range from Cambrian to Archean.
- HELIXIAN
- NEOHELIXIAN
- 9 [NH17] Diabasic olivine gabbro, intermediate and ultramafic intrusive rocks.
- NEOHELIXIAN AND EARLIER (?)
- 8 [NH15] Granite to granodiorite, massive to poorly foliated, porphyritic in part.
- PALEOHELIXIAN
- 7 [PH14], (GRNT) Granite, quartz monzonite, granodiorite, quartz diorite, syenite.
 - 6 [PH13], (QZMZ) Adamellite suite: adamellite, monzonite, agenite, granodiorite, granite and their hypersthene.
 - 5 [PH11], (ANRS) Anorthosite suite: anorthosite, anorthositic gabbro, leucotroctolite.
 - 4 [PH10], (UMFC) Gabbro, norite, anorthositic gabbro, troctolite, diorite, derived basic gneisses and amphibolite.
- APHEBIAN
- 3 [APW6], (GBBR) Gabbro-WAKUACH GABBRO-Gabbro, metagabbro, glomeroporphyritic gabbro and diorite.
- ARCHEAN
- 2 [ARC2] Massive granite and quartz monzonite.
 - 1 [ARC1], (PXGD) Massive to poorly foliated pyroxene-bearing granodiorite and syenodiorite.

* A four letter mnemonic name recorded as rock type as part of 1982 field observations.
 ** A four letter mnemonic name recorded as rock type as part of 1978 field observations.

Geological boundary
 Fault
 Mainly acidic volcanic rocks
 Mainly basic volcanic rocks
 No analytical result

This legend was modified and the geology derived for this geochemical map from Geology Map of Labrador, Mineral Resources Division, Department of Mines, Agriculture and Resources, Province of Newfoundland and Labrador.

Copies of map material and listings of field observations and analytical data, from which the material was prepared, may be available at users expense by application to:

K.G. Campbell Corporation
 880 Wellington St.
 Bay Z38
 Ottawa, Ontario
 K1R 6K7

The data is also available in digital form. For further information please contact:

The Director
 Computer Science Center
 Department of Energy, Mines and Resources
 Ottawa, Ontario
 K1A 0E4



Elevation in feet above mean sea level

Magnetic declination 1983 varies from 27°55.8' westerly at centre of west edge to 28°39.6' westerly at centre of east edge. Mean annual change -12.25' westerly

SILVER (ppm)
 OPEN FILE 901
 NATIONAL GEOCHEMICAL RECONNAISSANCE MAP 54-1982

CANADA-NEWFOUNDLAND
 co-operative mineral program 1982-1984

WEST LABRADOR 1982

Scale 1:250 000 Échelle

Universal Transverse Mercator Projection
 Projection transversale universelle de Mercator
 © Crown Copyrights reserved
 © Droits de la Couronne réservés

Base-map assembled by the Geological Cartography Unit from maps published at the same scale by the Surveys and Mapping Branch in 1965

