


BIOSTRATIGRAPHIC AND MATURATION STUDIES  
OF THE SCOTIAN SHELF  
PART 2  
HUSKY-BOW VALLEY et al.  
CHEBUCTO G-90



This document was produced  
by scanning the original publication.

Ce document est le produit d'une  
numérisation par balayage  
de la publication originale.

**BIOSTRATIGRAPHIC AND MATURATION  
STUDIES OF THE  
SCOTIAN SHELF**

**- PART 2 -**

HUSKY-BOW VALLEY et al. CHEBU<sup>K</sup>CTO G-90

**BY**

**LENTIN INTERNATIONAL  
BIOSTRATIGRAPHIC, LTD.**

Suite 2110 London House  
505 4th Ave. S.W.  
Calgary, Alberta, T2P 0J8  
(403) 264-0173

OPEN FILE  
DOSSIER PUBLIC  
18 53  
GEOLOGICAL SURVEY  
COMMISSION GEOLOGIQUE  
OTTAWA

7

## C O N T E N T S

INTRODUCTION.....	1
Chapter 1. PALYNOLOGY.....	2
Chapter 2. MICROPALAEONTOLOGY.....	7
Chapter 3. KEROGEN ANALYSIS.....	13
APPENDIX - A. Photographic record.....	15
APPENDIX - B. Foram and ostracod locations.....	31
APPENDIX - C. Kerogen analysis sheets.....	38

### MAP POCKETS:

#### PALYNOLOGY DISTRIBUTION CHART CHEBUCTO G-90

Part 1

Part 2

#### MICROPALAEO. DISTRIBUTION CHART CHEBUCTO G-90

Part 1

Part 2

1

BIOSTRATIGRAPHIC AND MATURATION STUDIES  
OF THE SCOTIAN SHELF

- PART 2

HUSKY-BOW VALLEY et al. CHEBUCTO G-90

GSC locality: D242

Location: 43 39'25.45"N; 50 42'43.93"W

K.B.elevation: 22m      Water depth: 122m

Casing set at: 396.2m, 922.3m, 3407m, 3713.4m, 4807.3m.

Total depth: 5234m      Interval studied: 440-5234m

Palynology by: S. de Gasparis.

Micropaleontology by: S.E. Cameron

INTRODUCTION

This is the second in a series of five reports detailing the palynology, micropaleontology kerogen and fluorescence analysis from four wells on the Scotian Shelf. Each specialty will be presented in separate chapters in each of the well reports. Rangecharts, which integrate graphically the three parts of the studies, are included in a pocket at the back of each report. The fifth and final report will compare and correlate the four wells.

Species cards with photographs of all palynological taxa will be presented with the final report. Appendix A of each well report contains a complete list of photographs of palynomorphs from that well. All foraminifera taxa are represented by at least one specimen, stained green and attached to a designated square in the assemblage slides. The depth and square number is given following the name of each taxon in Appendix B.

## CHAPTER 1

### PALYNOLOGY

One hundred sixty one cutting samples were studied from this well. All of the cuttings samples contained palynomorphs, with the exception of samples 3410-3420m and 4900-4910m, which were barren. The following biostratigraphic zonations and age determinations have been made:

0440-0550m Miocene

0570-0970m Oligocene

0990-1120m Early Oligocene

-Unconformity-

1140-1360m *A. reticulense* Peak Zone (Middle Eocene)

1380-1630m *A. senonensis* s. Gocht Peak Zone (Early Eocene)

1650-1720m Late Paleocene

1740-1775m *P. pyrophorum* Zone (Early Paleocene)

-Unconformity-

1795-1895m *O. operculata* Zone (Campanian)

1915-1925m *P. truncigerum* Zone (Santonian)

-Unconformity-

1945-1985m *K. williamsii* Zone (Cenomanian)

2005-2650m *C. cf. vestitum* Zone (Albian)

2670-3420m undiagnostic

3430-5180m *A. anaphrissa* Peak Zone (Barremian)

5200-5234m *C. elegantulum* Zone? (Hauterivian)

A questionably identified specimen of *Occisucysta* sp. A of Bujak

& Williams, 1978 is present at 5200-5210m, giving a possible age of Hauterivian to the lowest 34m of the well. Sedimentation is probably essentially uninterrupted in the Lower Cretaceous, resulting in a thickness of over thirty two hundred meters of Lower Cretaceous deposits. An unconformity is present in the Upper Cretaceous, as attested by the absence of part of the Cenomanian, the Turonian, the Coniacian and most of the Santonian record. The Upper Cretaceous sediments are two hundred and eighty meters thick. In the Tertiary, no record of Late Eocene sediments is present. The total thickness of the Tertiary sediments is in excess of thirteen hundred meters. Extensive reworking was observed in this well, most notably of *Adnatosphaeridium multispinosum* and *Areoligera senonensis* s. Gocht into Oligocene sediments, and of *Subtilisphaera perlucida* into Albian sediments.

The depositional environment appears to have been mostly marine, with the exception of level 3730-3740m, which contains only terrestrial fossils. The possible masking effect of caving at other levels cannot, however, be discounted.

#### SELECTED PALYNOMORPHS:

##### **0440-0550m: Miocene.**

*Pentadinium laticinctum*, *Operculodinium centrocarpum*, *Spiniferites pseudofurcatus*, *Tuberculodinium vancampoe*, *Tsugapollenites igniculus*, *Achomosphaera alvicornu*, *Achomophsaera ramulifera*, *Cordosphaeridium inodes*, *Glaphyrocysta ordinata*, *Hemicystodinium zoharii*, *Homotryblium tenuispinosum* (reworked), *Reticulatosphaera stellata*, *Operculodinium placitum*, *Spiniferites ramosus*, *Impagidinium paradoxum*.

##### **0570-0970m: Oligocene.**

*Cyclopsiella vieta*, *Tricolporites* sp. C of Williams & Brideaux 1975, *Glaphyrocysta microfenestrata*, *Lejeunecysta hyalina*, *Caryapollenites simplex*, *Ovoidites arcticus*, *Spiniferites granulatus*, *Spiniferites speciosus*, *Tricolporopollenites* sp. N of Williams & Brideaux 1975 (reworked), *Lingulodinium macherophorum*, *Cordosphaeridium cantharellum*, *Hystrichokolpoma unispinum* (reworked), *Hystrichosphaeropsis quasricibrata* (reworked).

This interval contains assemblages with dominant *Cyclopsiella vieta* alternating with dominant *Cyclopsiella conata*. *Cyclopsiella* is considered to be indicative of estuarine conditions.

0990-1120m: Early Oligocene.

*Wetzeliella ovalis*, *Wetzeliella symmetrica*, *Fibrocysta axialis*, *Hystriochokolpoma salacium*, *Distatodinium craterum*, *Thalassiphora delicata* s. Williams & Brideaux 1975, *Chiropteridium lobospinosum*, *Exochosphaeridium bifidum* (reworked), *Heteraulacacysta campanula*, *Thalassiphora pelagica*.

1140-1360m: *A. reticulense* Peak Zone (Middle Eocene).

*Dinopterygium fehmannense*, *Hystriochokolpoma cinctum*, *Homotryblium abbreviatum*, *Areoligera senonensis* s. Gocht, *Adnatosphaeridium multispinosum*, *Delandrea phosphoritica*, *Bipolaribucina paradoxum*, *Lejeunecysta granosa*, *Glaphyrocysta exuberans*, *Litosphaeridium siphoniphorum* (reworked), *Homotryblium pallidum*.

1380-1630m: *A. senonensis* Peak Zone (Early Eocene).

*Deflandrea oebisfeldensis*, *Apectodinium homomorphum*, *Heteraulacacysta porosa*, *Ceratiopsis speciosa glabra*, *Deflandrea andromienensis*, *Deflandrea heterophlycta*, *Membranophoridium aspinatum*, *Rhom-bodinium draco*, *Cordosphaeridium funiculatum*, *Areosphaeridium arcuatum*, *Areosphaeridium dictyoplokus*, *Diphyes colligerum*, *Distatodinium ellipticum*, *Gochtodinium spinulum*, *Rottnestia borussica*, *Cordosphaeridium gracile*, *Melitasphaeridium pseudorecurvatum*, *Spiniferites cingulatus*, *Spiniferites cornutus*, *Wetzeliella articulata conopia*, *Wetzeliella echinulata*, *Wilsonidium lineidentatum*, *Wilsonidium tabulatum*, *Cribroperidinium fetchamense* (reworked), *Leptodinium maculatum*, *Membranilarnacia ursulae*, *Turbiosphaera filosa*.

1650-1670m: Late Paleocene.

*Cordosphaeridium fibrospinosum*, *Palaeoperidinium pyrophorum*, *Ceratiopsis pannucea*, *Lejeunecysta hyalina*, *Oligosphaeridium complex*, *Turbiosphaera galatea*.

1740-1775m: *P. pyrophorum* Zone (Early Paleocene).

*Danea californica*, *Isabelidium bakeri*, *Ceratiopsis diebeli*, *Hystriochosphaeridium bowerbankii* (reworked).

1795-1825m: *O. operculata* Zone (Campanian).

*Hystriosphæridium bowerbankii*, *Odontochitina costata*, *Odontochitina porifera*, *Xenascus ceratioides*, *Xenascus gochtii*.

1915-1925m: *P. truncigerum* Zone (Santonian).

*Pervosphæridium truncigerum*, *Codoniella campanulata*.

1945-1985m: *K. williamsii* Zone (Cenomanian).

*Epelidosphaeridia spinosa*, *Florentinia cooksonae*, *Ariadnaesporites spinocaperatus*, *Oligosphaeridium pulcherrimum*, *Cyclonephelium vannophorum*, *Surculosphaeridium longifurcatum*, *Cyclonephelium distinctum*, *Cyclonephelium paucispinum*, *Litosphaeridium siphoniphorum*, *Oligosphaeridium asterigerum* (reworked), *Rugubivesciculites reductus*.

2005-2650m: *C. cf. vestitum* Zone (Albian).

*Chichaouadinium cf. vestitum*, *Oligosphaeridium poculum*, *Costatoperforosporites foveolatus*, *Vesperopsis mayi*, *Kiokansium williamsii*, *Palaeoperidinium cretaceum*, *Cribroperidinium edwardsii*, *Subtilisphaera perlucida* (reworked), *Aequitriradites ornatus*, *Florentinia verdierii*, *Odontochitina ancala*, *Senoniasphaera rotundata*, *Cyclonephelium chabaca*, *Palaeoperidinium sp. A* of Bujak & Williams, 1978, *Acanthotriletes reductus*, *Rugubivesciculites rugosus*, *Polysphaeridium lamina-spinosum*, *Appendicisporites bifurcatus*, *Ariadnaesporites fustiformis*, *Oligosphaeridium totum*, *Appendicisporites auritus*, *Appendicisporites problematicus*, *Cymososphaeridium validum*, *Florentinia mantellii*, *Palaeohytrichophora infusorioides*, *Rouseisporites reticulatus*, *Ascodinium scabrosum hostium*, *Gonyaulacysta cassidata*, *Taurocusporites segmentatus*, *Appendicisporites cristatus*, *Microreticulatisporites diatretus*, *Chichaouadinium vestitum*, *Callaiosphaeridium asymmetricum*, *Aequitriradites spinulosum*, *Canningia attadalica*, *Trilobosporites marylandensis*.

2670-3420m Undiagnostic.

No diagnostic fossils were found in this section of the well. Due to extensive reworking of Aptian fossils, notably *Subtilisphaera perlucida*, the Aptian cannot be located with assurance in this well.



3430-5180m *A. anaphrissa* Peak Zone (Barremian).

*Cicatricosisporites australiensis*, *Contignisporites cooksonae*, *Pseudoceratium pelliferum*, *Coronifera albertii*, *Endoceratium ludbrookiae*, *Callialasporites trilobatus*, *Occisucysta duxburii*, *Oligosphaeridium anthophorum*, *Hystrichosphaerina schindewolfii*, *Muderongia simplex*, *Occisucysta tentoria*, *Cerbia tabulata*, *Cleistosphaeridium polypes*, *Callaiosphaeridium trycherium*, *Stiphrosphaeridium dictyophorum*, *Trichodinium speetonense*, *Cribroridinium cooksonae*, *Aptea eisenackii*, *Tuberositriletes grosse-tuberculatus*, *Taleisphaera hydra*, *Aptea polymorpha*, *Oligosphaeridium diluculum*, *Heterosphaeridium heteracanthum*, *Subtilisphaera rotundata*, *Cicatricosisporites pseudotripartitus*, *Cicatricosisporites sprumontii*, *Veryhachium reductum*, *Muderongia parjata*, *Aptea anaphrissa*, *Pseudoceratium expolitum*, *Pseudoceratium parvum*.

5200-5234m *C. elegantulum* Zone? (Hauterivian).

*Occisucysta* sp. A of Bujak & Williams, 1978 was tentatively identified from this interval.

## CHAPTER 2

### MICROPALAEONTOLOGY

One hundred and sixty-two cuttings samples were studied for micropaleontology for Chebucto G-90. The following zonations have been made:

0415-0940m	Oligocene
0960-1450m	Upper Eocene
1470-1570m	Middle Eocene
1590-1630m	Lower Eocene
1650-1750m	Paleocene
-Unconformity-	
1765-1865m	Campanian
1885-1895m	Santonian
-Unconformity-	
1915-2015m	Cenomanian
2035-2255m	Albian
2275-3300m	Aptian
3320-4790m	Barremian
4810-5234m	Barremian or Hauterivian

The environment of deposition of the Barremian or Hauterivian section of the well was marginal marine to fresh or brackish swamp. The Barremian section was mostly marginal marine with a few inner neritic incursions in its lower part and some evidence of swamp conditions in the upper part. During the Aptian, conditions fluctuated between marginal and inner neritic at first and then became marginal. Albian deposition took place in a marginal marine environment. During the Cenomanian, conditions varied

from inner to middle neritic at first and then became outer neritic. During the Santonian and Campanian, the environment was outer neritic. it became bathyal at the base of the Paleocene, and did not change until late in the Upper Eocene. A shallowing episode caused conditions to become outer neritic in the late Upper Eocene, inner to outer neritic in the lower part of the Oligocene and inner neritic to marginal marine in the upper part of the Oligocene.

#### SELECTED FOSSILS:

**O415-O940m: Oligocene.**

**FORAMINIFERA:** *Turborotalia opima opima*, *Globigerina ouachitaensis*, *Orbulina universa*, *Globigerina officinalis*, *Nonionella auris*, *Cibicides floridana*, *Planulina depressa*, *Marginulina superba*, *Pullenia bulloides*, *Plectofrondicularia vauhani*, *Marginulinopsis tuberculata*, *Melonis planatus*, *Nodosaria stainforthi*, *Heterolepa pappen*, *Bulimina inflata*, *Tritaxia alazanensis*, *Oridorsalis umbonatus*, *Spiroplectammina mississippiensis*, *Globocassidulina subglobosa*, *Stilostomella curvatura*, *Melonis pompilioides*.

**ENVIRONMENT OF DEPOSITION:** Top of interval - low diversity benthonic and planktonic foraminifera, common echinoid and bivalve debris, suggesting an inner neritic to marginal marine environment. The presence of glauconite and bone fragments suggests a relatively low sedimentation rate, while frosted sand grains and carbonized plant rootlets indicate a depositional site close to shore in a relatively high energy environment. Abundant Miocene cavings are present in the upper part of the interval. More numerous planktonic individuals and higher benthonic diversity in the lower part of the interval suggest deeper depositional conditions.

**O960-1450m: Upper Eocene.**

**FORAMINIFERA:** *Globorotalia centralis*, *Globigerina venezuelana*, *Globigerina eocena*, *Siphonina danvillensis*, *Hoeglundina eocenica*, *Nodosaria lamellata*, *Stilostomella cookei*, *Heterolepa pseudoungerianus*, *Marginulinopsis texasensis*, *Planulina cooperensis*, *Guttulina elegans*, *Guttulina spicaeformis*, *Massilina decorata*, *Globocassidulina subglobosa*, *Marginulina hantkeni*, *Heterolepa pappen*, *Gyroidinoides planata*, *Stilostomella*

*subspinosa*, *Bulimina cooperensis*, *Uvigerina cocoaensis*, *Plectofrondicularia cookei*, *Ammodiscus glabratus*, *Ceratobulimina contraria*, *Pullenia quinqueloba*, *Melonis affine*, *Glomospira diffundens*, *Bulimina alazaensis*, *Siphonina jacksonensis*.

ENVIRONMENT OF DEPOSITION: in the upper and middle part of the interval, a well oxygenated outer neritic environment is indicated by the high diversity of benthonic and planktonic foraminifera. In the lower part, bathyal conditions are indicated by the presence of abundant deep water agglutinated and planktonic species. Much Miocene and Oligocene caving throughout.

1470-1570m : Middle Eocene.

FORAMINIFERA: *Pseudohastigerina micra*, *Globorotalia cerroazulensis cerroazulensis*, *Globorotalia cerroazulensis cocoaensis*, *Globorotalia cerroazulensis pommeroli*, *Globorotalia broedermanni*, *Globigerinathea mexicana*, *Globigerinathea index*, *Globorotalia bullbrooki*, *Globorotalia lehneri*, *Bulimina tuxpamensis*, *Pleurostomella cubensis*, *Bulimina impendens*.

ENVIRONMENT OF DEPOSITION: Very rich planktonic facies with highly diversified benthonic fauna suggests an upper slope bathyal marine environment. Moderate amount of Oligocene and Upper Eocene caving.

1590-1630m : Lower Eocene.

FORAMINIFERA: *Globorotalia acuta*, *Globorotalia aragonensis*, *Globorotalia pentacamerata*, *Pseudohastigerina wilcoxensis*, *Globorotalia formosa*, *Gavelinella cushmani*, *Bulimina trinitatensis*, *Tritaxia globulifera*, *Marssonella trinitatensis*, *Cibicidoides constrictus*, *Tritaxilina cubensis*, *Allomorphina subtriangularis*.

ENVIRONMENT OF DEPOSITION: Bathyal. Rich planktonic and diverse benthonic species including such deeper water indicators as *Cibicidoides*, *Tritaxia* and ornate *Bulimina* spp.. Very chalky facies with fairly common Oligocene and Eocene caving.

1765-1865m: Campanian.

FORAMINIFERA: *Planoglobulina glabrata*, *Pseudotextularia difformis*, *Rosita fornicata*, *Rosita contusa*, *Abathomphalus*

mayaroensis, *Heterohelix punctulata*, *Globotruncanella petaloidea*, *Planoglobulina brazoensis*, *Pseudotextularia elegans*, *Globotruncanita elevata*, *Globotruncanita stuartiformis*, *Marginotruncana marginata*, *Globotruncana bulloides*, *Globotruncana linneiana*, *Bolivinoidea draco miliaris*, *Neoflabellina rugosa leptodisca*, *Stensioina exsculpta exsculpta*, *Bolivina incrassata gigantea*, *Arenobulimina americana*, *Stensioina excolata*, *Stensioina pommerana*, *Gaudryina rudita*, *Dorothia bulletta*, *Heterostomella americana*, *Heterostomella austinana*, *Gaudryina austinana*.

ENVIRONMENT OF DEPOSITION: Very rich and diversified planktonic and benthonic faunas suggest an open marine, outer shelf environment. Abundant Maastrichtian indicators in the upper portion suggest that a thin Maastrichtian interval occurs in the top few meters or immediately above the interval studied. The Maastrichtian forms are well preserved, while the Campanian forms tend to be chalky.

1885-1895m : Santonian.

FORAMINIFERA: *Marginotruncana marginata*, *Marginotruncana coronata*, *Marginotruncana pseudolinneiana*, *Marginotruncana schneegansi*, *Whiteinella baltica*, *Whiteinella brittonensis*, *Dicarinella asymmetrica*, *Lenticulina munsteri*.

ENVIRONMENT OF DEPOSITION: Very chalky, planktonic rich environment. An open marine, outer shelf depositional site is suggested.

1915-2015m: Cenomanian.

FORAMINIFERA: *Praeglobotruncana stephani*, *Heterohelix moremani*, *Hedbergella delrioensis*, *Rotalipora greenhornensis*, *Gavelinella tourainensis*, *Gavelinella cenomanica*, *Ammobaculites comprimatus*, *Ammobaculites subcretaceus*, *Marssonella trochus*, *Quinqueloculina lirellangula*, *Tritaxia pyramidata*, *Quasispiroplectammina nuda*, *Gavelinella ammonoides*.

OSTRACODES: *Protocythere alexanderi*, *Schuleridea jonesiana*.

ENVIRONMENT OF DEPOSITION: Fairly good diversity of planktonics and benthonics suggests an open marine, outer shelf environment. A thin Turonian layer occurs in the uppermost part of this interval or immediately above. The lower part of the interval was deposited in a somewhat shallower, inner neritic environment,

where agglutinating foraminifers and ostracodes were much more frequent. Abundant Upper Cretaceous and Lower Tertiary cavings throughout.

2035-2255m: Albian.

FORAMINIFERA: *Ammobaculites goodlandensis*, *Lenticulina gaultina*, *Ammobaculites subcretaceus*, *Ammobaculites comprimatus*, *Ammobaculites reophacoides*.

OSTRACODES: *Eocytheropteron paenorbiculatum*, *Protocythere speetonensis*, *Cornicythere bonnemai*.

ENVIRONMENT OF DEPOSITION: Microfaunas consist of a dominance of shallow marine ostracodes and agglutinating foraminifers. A marginal marine environment is indicated. Many of the in situ species are limonite stained suggesting a very shallow, well oxygenated, low energy environment.

2275-3300m: Aptian.

FORAMINIFERA: *Favusella washitaensis*, *Hedbergella trochoidea*, *Hedbergella delrioensis*, *Lenticulina nodosa*, *Gavelinella intermedia*, *Gavelinella berthelini*, *Epistomina cretosa*, *Gavelinella cenomanica*, *Gavelinella barremiana*, *Tritaxia singularis*, *Epistomina carpenteri*.

OSTRACODES: *Mandocythere harrisiana*.

ENVIRONMENT OF DEPOSITION: Low diversity benthonic, especially agglutinating, foraminiferal fauna, and fairly common ostracodes indicate marginal marine conditions at the top of the interval. Slightly deeper inner neritic conditions exist in places in the lower part of the interval, where a few planktonic and calcareous benthonic species occur. Marginal marine conditions are also indicated by an abundance of coal, wood and carbonized plant rootlets.

3320-4790m: Barremian.

FORAMINIFERA: *Verneuilioides neocomiensis*, *Ammobaculites reophacoides*, *Gavelinella barremiana*, *Epistomina hechti*, *Reophax minuta*, *Gaudryinella tealbyensis*, *Textularia foeda*, *Marssonella kummi*, *Epistomina ornata*, *Epistomina caracolla*, *Marginulinopsis humilis*, *Planularia crepidularis*.

ENVIRONMENT OF DEPOSITION: The upper part of the interval contains low diversity agglutinating fauna and few ostracodes, sug

where agglutinating foraminifers and ostracodes were much more frequent. Abundant Upper Cretaceous and Lower Tertiary cavings throughout.

2035-2255m: Albian.

FORAMINIFERA: *Ammobaculites goodlandensis*, *Lenticulina gaultina*, *Ammobaculites subcretaceus*, *Ammobaculites comprimatus*, *Ammobaculites reophacoides*.

OSTRACODES: *Eocytheropteron paenorbiculatum*, *Protocythere speetonensis*, *Cornicythere bonnemai*.

ENVIRONMENT OF DEPOSITION: Microfaunas consist of a dominance of shallow marine ostracodes and agglutinating foraminifers. A marginal marine environment is indicated. Many of the *in situ* species are limonite stained suggesting a very shallow, well oxygenated, low energy environment.

2275-3300m: Aptian.

FORAMINIFERA: *Favusella washitaensis*, *Hedbergella trochoidea*, *Hedbergella delrioensis*, *Lenticulina nodosa*, *Gavelinella intermedia*, *Gavelinella berthelini*, *Epistomina cretosa*, *Gavelinella cenomanica*, *Gavelinella barremiana*, *Tritaxia singularis*, *Epistomina carpenteri*.

OSTRACODES: *Mandocythere harrisiana*.

ENVIRONMENT OF DEPOSITION: Low diversity benthonic, especially agglutinating, foraminiferal fauna, and fairly common ostracodes indicate marginal marine conditions at the top of the interval. Slightly deeper inner neritic conditions exist in places in the lower part of the interval, where a few planktonic and calcareous benthonic species occur. Marginal marine conditions are also indicated by an abundance of coal, wood and carbonized plant rootlets.

3320-4790m: Barremian.

FORAMINIFERA: *Verneuilinoides neocomiensis*, *Ammobaculites reophacoides*, *Gavelinella barremiana*, *Epistomina hechti*, *Reophax minuta*, *Gaudryinella tealbyensis*, *Textularia foeda*, *Marssonella kummi*, *Epistomina ornata*, *Epistomina caracolla*, *Marginulinopsis humilis*, *Planularia crepidularis*.

ENVIRONMENT OF DEPOSITION: The upper part of the interval contains low diversity agglutinating fauna and few ostracodes, sug

gesting a marginal marine environment for the most part. A complete absence of marine indicators and presence of coal and rootlets suggest brackish or fresh water, swamp conditions. The lower part of the interval contains a few marine incursions, with a few calcareous and agglutinating benthonics.

4810-5234m: Barremian or Hauterivian.

FORAMINIFERA: A few poorly preserved and undiagnostic agglutinating foraminifera.

ENVIRONMENT OF DEPOSITION: Very poor *in situ* foraminiferal fauna. A few, poorly preserved, undiagnostic agglutinating forms may be in place. The abundance of coal, goethite and other terrestrial indicators suggests an environment ranging from marginal marine to fresh or brackish water swamp.



## CHAPTER 3

### INTRODUCTION

Kerogen analysis has been completed on twenty-four samples. These samples, spaced approximately 200 meters apart in the well, have been examined for kerogen type, determination of the thermal alteration index (TAI), and epifluorescence. The data from these analyses are presented on analysis sheets provided by the Geological Survey and are included as Appendix C. A summary of each analysis follows.

### KEROGEN

It is impossible to determine the kerogen distribution in the 820m, 1000m, and 1210m samples with any degree of confidence due to the large amounts of mineral matter in the preparations. Most of the organic residue in this section appears to be finely degraded collinite with very rare spores and pollen. Below this level the samples are dominated by phyrogen and melanogen. The kerogen at 4820m appears to consist of coal fragments which may have been a mud additive.

Structured sapropels (spores, pollen, dinocysts and cuticle) vary from trace amounts to 20% of the total kerogen. High values were recorded at 610m, 1600m, 1805m, and 2015m, all within the immature section. Apart from these samples, the kerogen in the remainder of the well appears to be gas-prone.

### THERMAL ALTERATION INDEX

A series of color readings was taken on spores wherever possible and the results are presented on the chart. Where values fall between two TAI levels, both are plotted.

Some of the spores present down to 2015m suggest a higher maturity factor than other factors would indicate. These may have been reworked or partially oxydized prior to burial. The well penetrates an immature section down to 2015m. Below this level, maturity values equal to TAI 2 were encountered. The maturity level gradually increases to TAI 2+ at 3620m and TAI 3- at 4220m. At 5234m (T.D.), the maturity reaches a value of TAI 3. The oil window was reached at approximately 2225m, but the kerogens indicate a gas-prone source rock. The maturation levels appear to be high enough to generate wet gas at and below 4220m and the oil floor was probably reached at approximately 4700m.

## FLUORESCENCE

The strong fluorescence of the mounting medium strongly affected the study of the epifluorescence of the kerogens throughout the well. This was particularly noticeable below 3620m where the kerogens appear to have ceased fluorescing. The maturity values indicate that the palynomorphs should show some residual, very dull orange/brown colors down to at least 4400m.

Fluorescent material is very rare in the above-mentioned clay-dominated samples down to 1210m. *In situ* colors down to 1660m are generally a strong, bright yellow or yellow-green. There is a distinct reworked or oxydized component down to 1420m, which either does not fluoresce or shows dull orange colors. Cavings contaminants are obvious down to 2225m. The brightness of the *in situ* component diminishes and the colors shift through orange to orange/brown as the maturity increases. The fluorescence "floor" appears to be at 4220m, but this level is probably too high, due to the masking effect of the fluorescent mounting medium.

## APPENDIX A

## PHOTOGRAPHIC RECORD

Chebucto G-90

#	Name	Well #	Depth	Coordinate
S-1	Spiniferites sp. #TA WB/75	D242	500-510m	34.0 x 110.3
S-2	Homotryblum tenuispinosum	D242	540-550m	39.3 x 94.1
S-3	Operculodinium placitum	D242	540-550m	31.1 x 97.4
S-4	Operculodinium israelianum s. WB/75	D242	540-550m	37.5 x 100.4
S-5	Tuberculodinium sp. WB/75	D242	540-550m	34.0 x 103.8
S-6	Achomosphaera ramulifera	D242	500-510m	28.7 x 93.1
S-7	Glaphyrocysta sp. B WB/75	D242	500-510m	35.6 x 95.7
S-8	Canningia sp. #TA	D242	500-510m	39.6 x 95.1
S-9	Operculoidinium centrocarpum	D242	500-510m	35.7 x 98.0
S-10	Wetzeliiella sp. #TA	D242	500-510m	43.1 x 98.3
S-11	Glaphyrocysta ordinata	D242	500-510m	31.7 x 99.8
S-12	Tsugapollenites igniculus	D242	500-510m	41.9 x 106.3
S-13	Lentinia extensa	D242	500-510m	31.8 x 112.5
S-14	Lingulodinium sp. B WB/75	D242	470-480m	49.9 x 92.2
S-15	Pentadinium laticinctum	D242	470-480m	35.8 x 97.0
S-16	Osmundacidites sp.	D242	470-480m	34.6 x 108.3

S-17	<i>Tuberculodinium</i> <i>vancampoe</i>	D242	440-450m	44.4 x 109.5
S-18	<i>Spiniferites ramosus</i>	D242	570-580m	36.2 x 91.4
S-19	<i>Spiniferites speciosus</i>	D242	690-700m	34.2 x 97.3
S-20	<i>Spiniferites</i> <i>pseudofurcatus</i>	D242	570-580m	42.4 x 101.0
S-21	<i>Tricolporopollenites</i> sp. #TA	D242	600-610m	45.8 x 94.4
S-22	<i>Camarozonosporites</i> sp. #TA	D242	600-610m	29.9 x 92.1
S-23	? <i>Platicarya</i> sp. #TA	D242	810-820m	47.5 x 92.0
S-24	? <i>Corylus</i> sp. #TA	D242	780-790m	40.7 x 99.0
S-25	<i>Membranophoridium</i> sp. #TA	D242	720-730m	30.9 x 102.8
S-26	<i>Lingulodinium</i> <i>macherophorum</i>	D242	720-730m	39.7 x 100.8
S-27	<i>Homotryblum</i> <i>abbreviatum</i>	D242	1110-1120m	44.6 x 105.0
S-28	<i>Distatodinium craterum</i>	D242	1110-1120m	36.0 x 111.2
S-29	<i>Ascostomocystis potane</i>	D242	1140-1150m	36.0 x 111.2
S-30	<i>Canningia</i> <i>ringnesiorum</i>	D242	1140-1150m	29.2 x 101.2
S-31	<i>Taxodiaceopollenites</i> <i>hiatus</i>	D242	1140-1150m	05.5 x 103.9
S-32	<i>Fibrocysta axialis</i>	D242	1170-1180m	40.6 x 92.6
S-33	<i>Exochosphaeridium</i> <i>bifidum</i>	D242	1170-1180m	34.3 x 97.7
S-34	<i>Dinopterygium</i> <i>fehmarne</i>	D242	1200-1210m	43.6 x 91.9
S-35	<i>Thalassiphora pelagica</i>	D242	1230-1240m	45.6 x 100.5
S-36	<i>Heteraulacacysta</i> <i>campanula</i>	D242	1260-1270m	43.6 x 101.6
S-37	<i>Homotryblum pallidum</i>	D242	1260-1270m	44.7 x 109.8

S-38	Glaphyrocysta exhuberans	D242	1260-1270m	36.0 x 111.4
S-39	Cordosphaeridium sp. A WB/75	D242	810-820m	43.5 x 93.0
S-40	Hystrichokolpoma rigaudiae	D242	810-820m	38.5 x 97.0
S-41	Hystrichosphaeropsis quasicribrata	D242	720-730m	42.8 x 94.8
S-42	Impagidinium paradoxum	D242	570-580m	40.2 x 109.0
S-43	Impletosphaeridium sp. I s. M/76	D242	500-510m	31.5 x 110.7
S-44	Caryapollenites simplex	D242	600-610m	30.3 x 95.2
S-45	Cyclopsiella vieta	D242	600-610m	34.7 x 97.4
S-46	Lentinia serrata	D242	600-610m	49.0 x 97.7
S-47	Tricolpate grain	D242	600-610m	43.5 x 97.3
S-48	Onagraceous pollen	D242	600-610m	38.8 x 103.8
S-49	Kallosphaeridium cf. capulatum	D242	600-610m	45.4 x 104.8
S-50	Triporate grain	D242	600-610m	32.0 x 108.4
S-51	Faguspollenites sp. #TA	D242	600-610m	40.5 x 109.6
S-52	Achomosphaera #TA	D242	690-700m	35.0 x 99.1
S-53	Appendicisporites sp. #TA	D242	690-700m	38.0 x 98.2
S-54	Ovoidites arcticus	D242	690-700m	40.1 x 97.0
S-55	Spiniferites speciosus	D242	690-700m	43.4 x 92.9
S-56	Triatriopollenites sp. #TA	D242	660-670m	14.2 x 108.9
S-57	Impagidinium sp. #TA	D242	660-670m	39.6 x 107.0
S-58	Operculodinium sp. #TA	D242	660-670m	45.5 x 104.4
S-59	Tricolporites sp. C s. WB/75	D242	630-640m	19.6 x 103.8

S-60	<i>Glaphyrocysta</i> <i>microfenestrata</i>	D242	630-640m	42.4 x 104.0
S-61	<i>Thalassiphora delicata</i>	D242	630-640m	41.9 x 98.3
S-62	<i>Lejeunecysta hayalina</i>	D242	600-610m	35.2 x 112.0
S-63	<i>Adnatosphaeridium</i> <i>multispinosum</i>	D242	630-640m	35.6 x 96.4
S-64	<i>Lingulodinium</i> sp. A s. WB/75	D242	630-640m	30.0 x 93.6
S-65	<i>Apteodinium</i> sp. s. G/69	D242	1350-1650m	42.2 x 111.0
S-66	<i>Pyxidiniopsis</i> sp. #TA	D242	1350-1360m	42.0 x 102.2
S-67	<i>Chiropteridium</i> sp. #TA	D242	1350-1360m	30.0 x 95.5
S-68	? <i>Apectodinium</i> sp. #TC	D242	1350-1360m	29.9 x 92.8
S-69	<i>Kleithriasphaeridium</i> cf. <i>truncatum</i>	D242	1290-1300m	40.3 x 90.9
S-70	<i>Kleithriasphaeridium</i> <i>truncatum</i>	D242	1290-1300m	33.0 x 93.3
S-71	<i>Exochosphaeridium</i> sp. #TA	D242	1290-1300m	49.8 x 101.0
S-72	<i>Thalassiphora</i> <i>delicata</i> s. WB/75	D242	990-1000m	43.8 x 99.8
S-73	<i>Nyssapollenites</i> sp. #TA	D242	960-970m	32.2 x 101.5
S-74	<i>Liquidambar</i> sp. #TA	D242	960-970m	34.9 x 96.9
S-75	<i>Pterocarya</i> sp. #TA	D242	930-940m	22.0 x 108.5
S-76	<i>Cyclopsiella</i> sp. #TA	D242	930-940m	42.5 x 108.3
S-77	<i>Lejeunecysta spatiosa</i>	D242	930-940m	30.0 x 101.2
S-78	<i>Achomosphaera</i> <i>alcicornu</i>	D242	810-820m	41.5 x 104.5
S-79	<i>Spiniferites speciosus</i>	D242	810-820m	46.0 x 101.0
S-80	<i>Adnatosphaeridium</i> <i>caulleryi</i>	D242	810-820m	29.2 x 99.7

S-81	<i>Tuberculodinium</i> sp. s. WB/75	D242	810-820m	36.0 x 97.8
S-82	<i>Hemicystodinium</i> <i>zoharii</i>	D242	810-820m	41.0 x 97.1
S-83	<i>Hystriochokolpoma</i> <i>unispinum</i>	D242	810-820m	40.7 x 96.8
S-84	<i>Cyclopsiella coniata</i>	D242	810-820m	31.0 x 105.6
S-85	<i>Retitricolpites</i> sp. #TA	D242	900-910m	28.4 x 110.6
S-86	<i>Hafniasphaera saeptata</i>	D242	870-880m	42.0 x 93.7
S-87	<i>Apectodinium</i> sp. #TB	D242	840-850m	39.9 x 106.1
S-88	<i>Lejeunecysta</i> sp. #TA	D242	840-850m	34.3 x 93.6
S-89	<i>Palaeocystodinium</i> <i>gozlowense</i>	D242	840-850m	36.0 x 92.4
S-90	<i>Apectodinium</i> sp. #TA	D242	840-850m	45.6 x 91.0
S-91	<i>Selenopemphix</i> cf. <i>selenoides</i>	D242	840-850m	41.0 x 92.0
S-92	<i>Cordosphaeridium</i> <i>cantharellum</i>	D242	810-820m	39.0 x 106.2
S-93	<i>Momipites</i> sp. #TA	D242	990-1000m	06.0 x 103.8
S-94	<i>Eurydinium ingramii</i>	D242	0990-1000m	34.0 x 101.5
S-95	<i>Selenopemphyx</i> <i>nephroides</i>	D242	1020-1030m	43.0 x 96.2
S-96	<i>Wetzeliella</i> cf. <i>ovalis</i>	D242	1380-1390m	41.9 x 92.8
S-97	<i>Pentadinium laticinctum</i> <i>granulatum</i>	D242	1380-1390m	37.0 x 92.1
S-98	<i>Deflandrea</i> <i>phosphoritica</i>	D242	1380-1390m	40.3 x 96.7
S-99	<i>Wetzeliella symmetrica</i>	D242	1380-1390m	44.2 x 96.7
S-100	<i>Deflandrea medcalfii</i>	D242	1380-1390m	41.7 x 100.7
S-101	<i>Wetzeliella symmetrica</i> <i>incisa</i>	D242		32.7 x 102.5

S-102	Deflandrea oebisfeldensis	D242	1380-1390m	31.0 x 107.3
S-103	Ilexpollenites sp. A s. WB/75	D242	1380-1390m	09.0 x 103.8
S-104	Apectodinium homo- morphum quinquelatum	D242	1410-1420m	30.9 x 92.0
S-105	Apectodinium homomorphum	D242	1410-1420m	41.4 x 95.0
S-106	Deflandrea andromiensis	D242	1410-1420m	38.6 x 96.4
S-107	Membranophoridium aspinatum	D242	1410-1420m	45.2 x 97.4
S-109	Rhombodinium draco	D242	1410-1420m	48.3 x 101.0
S-110	Adnatosphaeridium cf. vittatum	D242	1410-1420m	43.8 x 101.3
S-111	Hystriochokolpoma salacium	D242	1410-1420m	32.9 x 101.4
S-112	Achilleodinium sp. #TA	D242	1410-1420m	42.6 x 107.3
S-113	Tricolporate grain 1	D242	780-790m	48.2 x 103.0
S-114	Deflandrea heterophlycta	D242	1410-1420m	35.5 x 112.6
S-115	Hystriochokolpoma eisenackii	D242	1410-1420m	28.3 x 112.5
S-116	Gen. et sp. indet.	D242	1440-1450m	36.6 x 93.0
S-117	Melitasphaeridium sp. #TA	D242	1440-1450m	35.0 x 92.5
S-119	Heteraulacacysta porosa	D242	1440-1450m	46.9 x 94.0
S-120	Cordosphaeridium sp. #TB	D242	1140-1450m	48.6 x 94.6
S-121	Apteodinium sp. B s. WB/75	D242	1440-1450m	44.0 x 94.2
S-122	Bipolaribucina paradoxum	D242	1140-1450m	30.0 x 95.0



S-123	<i>Areosphaeridium dictyoplokus</i>	D242	1470-1480m	46.3 x 91.5
S-124	<i>Glaphyrocysta</i> sp. #TB	D242	1470-1480m	42.2 x 92.7
S-125	<i>Cyclonephelium</i> sp. #TA	D242	1470-1480m	39.6 x 92.0
S-126	<i>Gochtodinium spinulum</i>	D242	1470-1480m	36.4 x 95.1
S-127	<i>Aerosphaeridium arcuatum</i>	D242	1470-1480m	33.8 x 96.0
S-128	<i>Lejeunecysta granosa</i>	D242	1470-1480m	32.8 x 98.0
S-129	<i>Wetzeliiella</i> sp. #TA	D242	1470-1480m	33.3 x 98.1
S-130	<i>Kisselovia</i> cf. <i>coleothrypta</i>	D242	1470-1480m	40.5 x 97.5
S-132	<i>Distatodinium ellipticum</i>	D242	1500-1510m	48.0 x 92.5
S-133	<i>Melitasphaeridium pseudorecurvatum</i>	D242	1500-1510m	36.4 x 98.2
S-134	<i>Letinia</i> sp. #TA	D242	1500-1510m	39.0 x 98.5
S-135	<i>Cleistosphaeridium</i> cf. <i>tenuifilum</i>	D242	1500-1510m	42.0 x 98.7
S-136	<i>Wetzeliiella articulata</i> v. <i>conopia</i>	D242	1560-1570m	45.5 x 92.6
S-137	<i>Diphyes colligerum</i>	D242	1560-1570m	45.5 x 92.6
S-138	<i>Wilsonidium tabulatum</i>	D242	1560-1570m	40.7 x 92.6
S-139	<i>Wilsonidium lineidentatum</i>	D242	1560-1570m	28.8 x 92.5
S-140	<i>Firocysta</i> sp. #TA	D242	1560-1570m	47.7 x 93.3
S-141	<i>Rottnestia borussica</i>	D242	1560-1570m	30.4 x 95.0
S-142	<i>Wetzeliiella echinulata</i>	D242	1560-1570m	47.1 x 98.2
S-143	<i>Hystrihostrogylon</i> sp. #TA	D242	1560-1570m	33.7 x 104.0
S-144	<i>Spiniferites cornutus</i>	D242	1560-1570m	31.3 x 104.4
S-147	<i>Cannosphaeropsis</i> sp. #TA	D242	1590-1600m	36.6 x 96.0

S-148	Gonyaulacysta sp. #TA	D242	1620-1630m	45.6 x 92.2
S-149	Membranilarnacia ursulae	D242	1620-1630m	35.0 x 92.2
S-151	Leptodinium maculatum	D242	1620-1630m	32.7 x 93.6
S-152	Heteraulacacysta sp. #TA	D242	1620 1630m	30.7 x 93.0
S-153	Areoligera senonensis s. G/69	D242	1620-1630m	31.9 x 96.4
S-154	Cordosphaeridium gracilis	D242	1620-1630m	97.0 x 96.0
S-155	Cribroperidinium fetchamense	D242	1620-1630m	47.0 x 97.5
S-156	Turbiosphaera filosa	D242	1620-1630m	39.5 x 92.6
S-157	Ceratiopsis speciosa glabra	D242	1620-1630m	32.9 x 95.2
S-158	Cordosphaeridium fibrospinosum	D242	1650-1660m	41.5 x 104.4
S-159	Oligosphaeridium complex	D242	1680-1690m	47.5 x 93.6
S-159	Palaeoperidinium pyrophorum	D242	1680-1690m	32.3 x 92.6
S-160	Ceratiopsis pannucea	D242	1580-1690m	54.7 x 102.9
S-161	Turbiosphaera galatea	D242	1680-1690m	29.8 x 107.7
S-162	Palaeocystodinium sp. #TA	D242	1680-1690m	38.7 x 112.4
S-163	Cleistosphaeridium sp. #TA	D242	1710-1720m	45.4 x 91.3
S-164	Deflandrea sp. #TA	D242	1710-1720m	29.6 x 93.3
S-165	Isabelidinium bakeri	D242	1710-1720m	44.7 x 95.6
S-166	Oligosphaeridium sp. #TA	D242	1710-1720m	43.2 x 99.1
S-167	Ceratiopsis diebeli	D242	1765-1775m	41.0 x 95.6
S-168	Danea mutabilis	D242	1740-1750m	30.0 x 95.5

S-169	<i>Hystrichosphaeridium bowerbankii</i>	D242	1765-1775m	39.8 x 107.2
S-170	<i>Odontochitina operculata</i>	D242	1795-1805m	40.5 x 94.2
S-171	<i>Criboperidinium</i> sp. #TB	D242	1795-1805m	42.7 x 95.0
S-172	<i>Odontochitina porifera</i>	D242	1795-1805m	34.7 x 97.1
S-173	<i>Turbiosphaera magnifica</i>	D242	1825-1835m	42.8 x 99.4
S-175	<i>Craspedodinium</i> sp. #TA	D242	1855-1865m	46.1 x 97.8
S-176	<i>Xenascus ceratioides</i>	D242	1855-1865m	34.8 x 99.4
S-177	<i>Xenascus gochtii</i>	D242	1855-1865m	39.1 x 102.6
S-178	<i>Odontochitina/Xenascus</i>	D242	1855-1865m	40.6 x 102.6
S-179	? <i>Wallodinium</i> sp. #TA	D242	1855-1865m	36.1 x 106.6
S-180	<i>Hystrichodinium</i> sp. #TA	D242	1855-1865m	32.8 x 108.2
S-181	<i>Hystrichosphaeridium truncigerum</i>	D242	1885-1895m	35.4 x 96.5
S-182	<i>Codoniella campanulata</i>	D242	1885-1895m	34.6 x 96.5
S-183	<i>Cyclonephelium vannophorum</i>	D242	1915-1925m	36.2 x 95.5
S-184	<i>Surculosphaeridium longifurcatum</i>	D242	1915-1925m	34.6 x 95.2
S-185	<i>Ariadnaesporites spinocaperatus</i>	D242	1915-1925m	44.6 x 97.3
S-186	<i>Florentinia cooksoniae</i>	D242	1915-1925m	30.8 x 100.2
S-187	<i>Oligosphaeridium anthophorum</i>	D242	1915-1925m	37.7 x 101.9
S-188	<i>Epelidosphaeridia spinosa</i>	D242	1915-1925m	35.2 x 103.8
S-189	<i>Membranophoridium</i> sp. #TB	D242	1915-1925m	36.0 x 105.0

S-189B	Arcellites sp. #TA	D242	1915-1925m	36.0 x 105.0
S-192	Oligosphaeridium asterigerum	D242	1945-1955m	41.9 x 91.8
S-193	Litosphaeridium siphoniphorum	D242	1945-1955m	48.6 x 94.6
S-194	Cyclonephelium distinctum	D242	1945-1955m	44.9 x 94.8
S-195	Biharisporites sp. #TA	D242	1945-1955m	44.8 x 109.1
S-196	Kiokansium williamsii	D242	1975-1985m	33.7 x 93.6
S-197	Chichaouadinium cf. vestitum	D242	1975-1985m	42.2 x 93.6
S-198	Cribroperidinium edwardsii	D242	2005-2015m	34.0 x 91.8
S-199	Odontochitina cf. rhakodes	D242	2005-2015m	35.6 x 92.5
S-200	Florentinia deanei	D242	2005-2015m	36.2 x 93.0
S-201	Vesperopsis sp. #TA	D242	2005-2015m	42.4 x 95.6
S-202	Vesperopsis mayi	D242	2035-2045m	42.2 x 97.9
S-203	Aequitriradites ornatus	D242	2035-2045m	45.1 x 104.6
S-204	Subtilisphaera perlucida	D242	2035-2045m	23.0 x 101.4
S-204	Cicatricosisporites angustus	D242	2035-2045m	43.0 x 204.5
S-205	Palaeohystrichophora infusorioides	D242	2065-2075m	41.8 x 93.6
S-206	?Appendicisporites	D242	2065-2075m	46.5 x 104.0
S-207	Odontochitina ancala	D242	2065-2075m	47.5 x 105.5
S-208	Senoniasphaera rotundata	D242	2065-2075m	37.0 x 110.8
S-209	Chlamydophorella sp. #TA	D242	2095-2105m	42.3 x 91.8
S-210	Cibotiidites sp. #TA	D242	2095-2104m	35.6 x 94.0

S-212	<i>Rugubivesiculites reductus</i>	D242	2115-2125m	33.8 x 94.6
S-213	<i>Acanthotriletes</i> sp. #TA	D242	2115-2125m	41.0 x 102.8
S-214	<i>Polysphaeridium laminaspinosum</i>	D242	2155-2165m	30.2 x 91.8
S-215	? <i>Exochosphaeridium</i> sp.	D242	2155-2165m	29.2 x 91.4
S-216	<i>Appendicisporites bifurcatus</i>	D242	2155-2165m	46.5 x 98.0
S-217	<i>Rouseisporites</i> sp. #TA	D242	2155-2165m	35.1 x 107.2
S-218	<i>Ariadnaesporites fustiformis</i>	D242	2155-2165m	30.5 x 110.3
S-219	<i>Florentinia verdieri</i>	D242	2185-2195m	35.5 x 104.3
S-220	<i>Oligosphaeridium</i> sp. #TA	D242	2215-2525m	49.6 x 98.0
S-222	<i>Costatoperforosporites foveolatus</i>	D242	2215-2525m	35.5 x 93.2
S-224	<i>Biretisporites</i> sp. #TA	D242	2215-2225m	31.5 x 98.7
S-225	<i>Appendicisporites auritus</i>	D242	2215-2225m	43.8 x 99.5
S-226	<i>Oligosphaeridium totum</i>	D242	2215-2225m	29.5 x 106.0
S-227	<i>Appendicisporites problematicus</i>	D242	2215-2225m	36.7 x 108.0
S-228	<i>Cymosphaeridium validum</i>	D242	2245-2255m	44.5 x 91.2
S-229	<i>Oligosphaeridium poculum</i>	D242	2245-2255m	48.2 x 97.7
S-230	<i>Megaspore</i> sp. #TA	D242	2245-2255m	40.2 x 99.7
S-230	<i>Cibotioidites</i> sp. #TB	D242	2245-2255m	43.2 x 99.7
S-231	<i>Appendicisporites</i> sp. #CB	D242	2275-2285m	29.6 x 99.5
S-232	<i>Florentinia mantellii</i>	D242	2275-2285m	29.6 x 100.6

S-233	<i>Palaeoperidinium</i> <i>cretaceum</i>	D242	2275-2284m	34.9 x 103.7
S-234	<i>Cribroperidinium</i> sp. #TC	D242	2275-2285m	32.6 x 112.3
S-235	<i>Kiokansium</i> sp. #TA	D242	2305-2315m	28.6 x 93.4
S-236	<i>Ascodinium</i> <i>scabrosum</i> <i>ostium</i>	D242	2305-2315m	31.8 x 94.1
S-237	<i>Gonyaulacysta</i> <i>cassidata</i>	D242	2305-2315m	35.0 x 96.6
S-238	<i>Apteodinium</i> sp. #TA	D242	2305-2315m	44.0 x 92.2
S-239	<i>Cyclonephelium</i> <i>chabaca</i>	D242	2305-2315m	28.6 x 107.0
S-240	<i>Pterodinium</i> sp. #TB	D242	2365-2375m	32.3 x 94.3
S-241	<i>Pterodinium</i> sp. #TC	D242	2400-2410m	39.7 x 97.2
S-242	<i>Taurocusporites</i> <i>segmentatus</i>	D242	2400-2410m	35.5 x 97.8
S-243	<i>Aequitriradites</i> sp. #TA	D242	2430-2440m	34.1 x 93.4
S-244	<i>Appendicisporites</i> <i>cristatus</i>	D242	2430-2440m	35.8 x 96.0
S-245	<i>Kiokansium</i> sp. #TB	D242	2430-2440m	36.1 x 111.3
S-246	<i>Cicatricosisporites</i> sp. #TA	D242	2460-2470m	37.3 x 94.6
S-247	? <i>Tubotuberella</i> sp. #TA	D242	2520-2530m	37.2 x 107.8
S-248	<i>Callaiosphaeridium</i> <i>asymmetricum</i>	D242	2550-2560m	40.6 x 111.4
S-249	<i>Microreticulati-</i> <i>sporites diatretus</i>	D242	2580-2590m	37.6 x 94.0
S-250	<i>Canningia</i> <i>attadalica</i>	D242	2610-2620m	30.8 x 111.1
S-251	<i>Protoellipsodinium</i> <i>spinosum</i>	D242	2640-2650m	46.6 x 102.2
S-252	<i>Trilobosporites</i> <i>marylandensis</i>	D242	2670-2680m	46.0 x 92.7

S-253	<i>Concavissimisporites</i> <i>apigranulosus</i>	D242	2670-2680m	35.2 x 103.7
S-254	<i>Nyktericysta</i> <i>cf. vitrea</i>	D242	2850-2860m	30.7 x 104.2
S-255	<i>Neoraistrichia</i> <i>truncata</i>	D242	2940-2950m	33.4 x 100.0
S-256	<i>Nyktericysta davisii</i>	D242	2970-2980m	34.9 x 86.6
S-257	<i>Chichaouadinium</i> <i>vestitum</i>	D242	3090-3100m	40.0 x 99.4
S-258	<i>Maculatisporites</i> <i>microverrucatus</i>	D242	3120-3130m	36.0 x 101.0
S-259	<i>Achomosphaera</i> <i>verdieri</i>	D242	3180-3190m	39.0 x 96.7
S-260	<i>Pilosisorites</i> <i>trichopapillosus</i>	D242	3320-3330m	38.9 x 103.0
S-261	<i>Systematophora</i> <i>cf. silyba</i>	D242	3430-3440m	37.8 x 96.2
S-262	<i>Subtilisphaera terrula</i>	D242	3430-3440m	49.2 x 102.4
S-263	<i>Callialasporites</i> <i>trilobatus</i>	D242	3430-3440m	47.2 x 104.1
S-264	<i>Canningia</i> sp. #TB	D242	3430-3440m	38.6 x 107.6
S-265	<i>Canningia</i> sp. #TE	D242	3430-3440m	49.1 x 107.6
S-266	<i>Pilosisorites</i> sp. #TA	D242	3430-3440m	46.0 x 109.5
S-267	<i>Cicatricosisporites</i> <i>australiensis</i>	D242	3460-3470m	35.7 x 93.9
S-268	<i>Oligosphaeridium</i> <i>anthophorum</i>	D242	3460-3470m	45.8 x 108.0
S-269	<i>Hystrichosphaerina</i> <i>schindewolfii</i>	D242	3520-3530m	42.1 x 93.3
S-270	<i>Occisucysta tentoria</i>	D242	3580-3590m	43.2 x 100.8
S-271	<i>Amphorosphaeridium</i> sp. #TA	D242	3580-3590m	45.0 x 102.7
S-272	<i>Endoscrinium</i> cf. <i>galeritum</i>	D242	3580-3590m	43.8 x 102.1

S-273	<i>Contignisporites cooksonae</i>	D242	3640-3650m	29.8 x 97.0
S-274	<i>Cerbia tabulata</i>	D242	3670-3680m	33.8 x 102.0
S-274	<i>Pseudoceratium pelliiferum</i>	D242	3670-3680m	46.0 x 102.2
S-285	<i>Cleistosphaeridium polypes s.s.</i>	D242	3790-3800m	44.7 x 92.0
S-276	<i>Callaiosphaeridium trycherium</i>	D242	3910-3920m	38.8 x 108.4
S-278	<i>Stiphrosphaeridium dictyophorum</i>	D242	3940-3950m	47.0 x 92.8
S-279	<i>Appendicisporites</i> sp. #TC	D242	3970-3980m	43.1 x 106.7
S-280	<i>Trichodinium speetonensis</i>	D242	4060-4070m	41.6 x 95.9
S-281	<i>Kleithriasphaeridium</i> sp. #TA	D242	4060-4070m	44.9 x 98.1
S-282	<i>Pterodinium</i> sp. #TD	D242	4060-4070m	38.2 x 106.4
S-283	<i>Occisucysta duxburyi</i>	D242	4060-4070m	45.1 x 107.4
S-284	<i>Canningia</i> sp. #TC	D242	4090-4100m	34.7 x 100.3
S-285	<i>Trilobosporites</i> sp. #TA	D242	4120-4130m	43.2 x 96.0
S-286	<i>Tuberositriletes grossetuberculatus</i>	D242	4120-4130m	45.3 x 104.0
S-287	<i>Cribroperidinium cooksonae</i>	D242	4250-4160m	43.4 x 112.2
S-288	<i>Coronifera albertii</i>	D242	4030-4040m	40.7 x 99.1
S-289	<i>Polystephanephorus</i> sp. #TA	D242	4180-4190m	45.8 x 101.2
S-290	<i>Aptea polymorpha</i>	D242	4300-4310m	20.1 x 103.3
S-291	<i>Taleisphaera hydra</i>	D242	4270-4280m	39.0 x 93.3
S-292	<i>Gleicheniidites</i> sp. #TB	D242	4270-4280m	38.3 x 99.9
S-293	<i>Aptea eisenackii</i>	D242	4270-4280m	34.6 x 103.2



S-294	<i>Oligosphaeridium diluculum</i>	D242	4300-4310m	44.3 x 103.5
S-295	<i>Kreuselisporites</i> sp. #TA	D242	4300-4310m	52.1 x 103.5
S-296	<i>Microreticulatisporites</i> sp. #TA	D242	4330-4340m	30.0 x 101.8
S-297	<i>Pseudoceratium pelliiferum</i>	D242	4330-4340m	39.6 x 103.3
S-300	<i>Oligosphaeridium</i> cf. complex	D242	4360-4370m	47.4 x 100.7
S-301	<i>Subtilisphaera</i> rotundata	D242	4420-4430m	47.0 x 95.0
S-302	<i>Canningia</i> sp. #TC	D242	4420-4430m	24.3 x 104.6
S-303	<i>Cicatricosisporites pseudotripartitus</i>	D242	4450-4460m	29.5 x 92.0
S-303A	<i>Cicatricosisporites sprumontii</i>	D242	4450-4460m	31.0 x 94.7
S-304	<i>Contignisporites</i> cf. fornicatus B/80	D242	4450-4460m	44.3 x 96.6
S-305	<i>Cerbia</i> sp. #TA	D242	4450-4460m	46.1 x 99.9
S-306	<i>Callaiosphaeridium</i> sp. #TA	D242	4480-4490m	35.8 x 94.1
S-309	<i>Staplinisporites</i> sp. #TA	D242	4540-4550m	43.3 x 101.3
S-310	<i>Cribroperidinium</i> sp. #TD	D242	4540-4550m	32.0 x 102.4
S-313	<i>Tehamadinium</i> cf. tenuiceras	D242	4570-4580m	38.2 x 96.5
S-314	<i>Lycopodiumsporites</i> sp. #TA	D242	4660-4670m	29.0 x 112.3
S-315	<i>Muderongia parvata</i>	D242	4690-4700m	42.2 x 107.5
S-316	<i>Endoceratium</i> cf. ludbrookiae	D242	4720-4730m	32.6 x 101.7
S-319	<i>Coronifera</i> sp. #TB	D242	4720-4730m	31.1 x 110.0

S-320	<i>Pseudoceratium</i> <i>expolitum</i>	D242	4780-4790m	49.6 x 98.1
S-321	<i>Pseudoceratium</i> <i>parvum</i>	D242	4780-4790m	45.2 x 105.0
S-322	<i>Aequitriradites</i> sp. #TB	D242	5200-5210m	29.5 x 92.7
S-323	? <i>Occisucysta</i> sp. A s. BW/78	D242	5200-5210m	43.0 x 97.9
S-324	<i>Palaeoperidinium</i> <i>cretaceum</i>	D242	2365-2375m	32.4 x 96.1
S-327	<i>Alnipollenites</i> sp. #TA	D242	960-970m	32.7 x 94.4
S-328	<i>Quercoidites</i> sp. #TA	D242	1170-1180m	21.7 x 104.3
S-118	<i>Cordosphaeridium</i> <i>funiculatum</i>	D242	1440-1450m	33.6 x 91.6
S-143	<i>Caryapollenites</i> sp. #TA	D242	990-1000m	43.7 x 105.7
S-144	<i>Cribroperidinium</i> sp. #TA	D242	1590-1600m	34.0 x 103.9
S-145	<i>Pyxidiniopsis</i> sp. #TB	D242	1590-1600m	34.6 x 98.3
S-173	<i>Ulmipollenites</i> sp. #TA	D242	1050-1060m	10.8 x 111.6
S-174	<i>Retitricolpites</i> sp. #TA	D242	780-790m	38.5 x 92.6
S-298	<i>Heterosphaeridium</i>	D242	4330-4340m	34.0 x 104.8
S-299	<i>Contignisporites</i> sp. #TA	D242	4360-4370m	36.3 x 101.6
S-307	<i>Tehamadinium</i> cg. sp. #TA	D242	4510-4530m	29.6 x 106.8
S-308	<i>Veryhachium reductum</i>	D242	4540-4550m	47.8 x 96.1
S-324	<i>Diervilla echinata</i>	D242	1140-1150m	30.1 x 99.3
S-325	<i>Membranophoridium</i> , sp. #TC	D242	1170-1180m	47.3 x 98.7

S-325	<i>Spiniferites</i> cf. <i>cingulatus</i>	D242	1500-1510m	36.4 x 102.6
S-327	<i>Spiniferites</i> <i>granulatus</i>	D242	690-700m	43.3 x 92.3
S-328	<i>Tricolporopollenites</i> sp. N s. WB/75	D242	690-700m	40.7 x 111.3
S-311	<i>Odontochitina</i> sp. #TA	D242	4540-4550m	43.3 x 102.5
S-312	<i>Subtilisphaera</i> <i>perlucida</i>	D242	4570-4580m	45.0 x 97.4
S-325	<i>Tiliaepollenites</i> sp. #TA	D242	780-790m	38.4 x 104.6
S-326	<i>Tripoporopollenites</i> sp. F s. WB/75	D242	780-790m	37.6 x 97.3
S-8	<i>Spiniferites</i> sp. A s. WB/75	D242	540-550m	45.1 x 100.4

## Appendix B

### LOCATION OF FORAMINIFERA AND OSTRACODE SPECIES

0415-0940m: Oligocene.

#### FORAMINIFERA

*Turborotalia opima opima* (440-50, sq. 6)  
*Globigerina cf. praebulloides* (440-50, sq. 18)  
*Globigerina cf. bulloides* (470-80, sq. 8)  
*Globigerina cf. trilobus* (500-10, sq. 20)  
*Globigerina cf. senelis* (540-50, sq. 18)  
*Globigerina ouachitaensis* (570-80, sq. 5)  
*Orbulina universa* (750-60, sq. 21)  
*Globigerina officinalis* (840-50, sq. 19)  
*Globigerina cf. tripartita* (840-50, sq. 32)  
*Marginulina cf. glabra* (415-25, sq. 5)  
*Nonionella auris* (480-90, sq. 31)  
*Textularia cf. badensis* (470-80, sq. 22)  
*Cibicides floridana* (540-50, sq. 7)  
*Planulina depressa* (570-80, sq. 6)  
*Uvigerina cf. danvillensis* (630-40, sq. 18)  
*Marginulina subperba* (630-40, sq. 31)  
*Pullenia bulloides* (630-40, sq. 19)  
*Hoeglundina cf. eocenica* (630-40, sq. 30)  
*Plectofrondiculina vauhan* (720-30, sq. 5)  
*Marginulinopsis tuberculata* (720-30, sq. 7)  
*Melonis planatus* (720-30, sq. 20)  
*Nodosaria stainforthi* (720-30, sq. 19)  
*Sphaeroidina cf. bulloides* (720-30, sq. 18)  
*Heterolepa peppeni* (720-30, sq. 17)  
*Bulimina inflata* (720-30, sq. 31)  
*Tritaxia alazanensis* (750-60, sq. 8)  
*Uvigerina cf. yazooensis* (750-60, sq. 19)  
*Gyroidinoides cf. soldani* (750-60, sq. 9)  
*Oridorsalis umbonatus* (750-60, sq. 22)  
*Spiroplectammina mississippiensis* (780-90, sq. 7)  
*Bolvina cf. gardnerai* (780-90, sq. 31)  
*Globocassidulina subglobosa* (780-90, sq. 8)  
*Stilostomella curvatura* (780-90, sq. 20)  
*Melonis pompiliodes* (810-20, sq. 21)

0960-1450m: Upper Eocene

#### FORAMINIFERA

*Globorotalia cennetralis* (960-70, sq. 8)

*Globigerina venezuelana* (1380-90, sq. 20)  
*Globigerina eocaena* (1380-90, sq. 22)  
*Siphonina danvillensis* (960-70, sq. 7)  
*Hoeflundina eocenica* (960-70, sq. 9)  
*Ceratobulimia* af. *eximia* (960-70, sq. 33)  
*Nodosaria lamellata* (960-70, sq. 32)  
*Stilostomella cookei* (960-70, sq. 22)  
*Uvigerina* cf. *glabrans* (960-70, sq. 10)  
*Heterolepa pseudoungerinaus* (960-70, sq. 46)  
*Marginulinopsis texasensis* (960-70, sq. 20)  
*Planulina cooperensis* (990-1000, sq. 7)  
*Guttulina elegans* (990-1000, sq. 8)  
*Guttulina spicaeformis* (990-1000, sq. 20)  
*Gyroidinoides* cf. *soldanii* (990-1000, sq. 32)  
*Massilina decorata* (1050-60, sq. 21)  
*Globocassidulina subglobosa* (1050-60, sq. 9)  
*Marginulina hantkeni* (1050-60, sq. 20)  
*Heterolepsa pippeni* (1080-90, sq. 21)  
*Cibicides* cf. *lobatus* (1080-90, sq. 9)  
*Gyroidinoides planata* (1110-20, sq. 9)  
*Stilostomella subspinosa*  
*Cassidulinoides* sp. #CA (1140-50, sq. 6)  
*Bulimina cooperensis* (1170-80, sq. 7)  
*Uvigerina cocoaensis* (1200-10, sq. 6)  
*Uvigerina cocoaensis* (1230-40, sq. 7)  
*Plectofrondicularia cookei* (1230-40, sq. 6)  
*Martinotiella* cf. *petrosa* (1350-60, sq. 22)  
*Ammodiscus glabratus* (1350-60, sq. 10)  
*Cyclammina* cf. *cancellata* (1350-60, sq. 8)  
*Ceratobulimina contraria* (1380-90, sq. 7)  
*Hoeglundina* af. *eocenica* (1380-90, sq. 8)  
*Pullenia quinqueloba* (1380-90, sq. 21)  
*Melonis affine* (1380-90, sq. 10)  
*Martinotiella* cf. *cocoaensis* (1380-90, sq. 34)  
*Glomospira diffundens* (1410-20, sq. 9)  
*Bulimina alazanensis* (1440-50, sq. 19)  
*Uvigerina cocoaensis* (1440-50, sq. 7)  
*Siphonina jacksonensis* (1440-50, sq. 32)  
*Valvulina* cf. *jarvisi* (1440-50, sq. 8)

#### OSTRACODES:

*Henryhowella* sp. #CA (990-1000, sq. 44)

#### 1470-1570m: Middle Eocene

*Pseudohastigerina micra* (1470-80, sq. 32)  
*Globorotalia cerroazulensis* s.s (1480-90, sq. 8)

*Globorotalia cerroazulensis cocoaensis* (1470-80, sq. 20)  
*Globorotalia cerroazulensis pomeroli* (1470-80, sq. 9)  
*Globorotalia broedermanni* (1470-80, sq. 21)  
*Globigerinatheca mexicana* (1470-80, sq. 7)  
*Globigerinatheca index* (1470-80, sq. 19)  
*Globorotalia bulbrookii* (1470-80, sq. 22)  
*Globorotalia cf. spinulosa* (1470-80, sq. 34)  
*Globorotalia lehneri* (1500-10, sq. 8)  
*Bulimina tuxpamensis* (1500-10, sq. 7)  
*Bulimina cf. semicostata* (1500-10, sq. 19)  
*Pleurostomella cubensis* (1569-70, sq. 8)  
*Bulimina impendens* (1560-70, sq. 18)

OSTRACODES:

*Krithe* sp. #CA (1470-80, sq. 10)

1590-1630m: Lower Eocene

*Globorotalia acuta* (1590-1600, sq. 30)  
*Globorotalia aragonensis* (1590-1600, sq. 32)  
*Globorotalia pentacamerata* (1590-1600, sq. 31)  
*Pseudohastigerina wilcoxensis* (1620-30, sq. 7)  
*Globorotalis formosa* (1620-30, sq. 32)  
*Chiloguembelina cf. moresii* (1620-30, sq. 18)  
*Gavelinella cushmani* (1590-1600, sq. 20)  
*Bulimina trinitatensis* (1590-1600, sq. 8)  
*Anomalinoidea cf. vulgaris* (1590-1600, sq. 18)  
*Tritaxia cf. jarvisi* (1590-1600, sq. 5)  
*Tritaxia globulifera* (1590-1600, sq. 6)  
*Marssonella trinitatensis* (1590-1600, sq. 17)  
*Cibicidioides constrictus* (1590-1600, sq. 29)  
*Marginuliopsis cf. enbornensis* (1620-30, sq. 19)  
*Tritaxia cubensis* (1620-30, sq. 9)  
*Allomorpha subtriangularis* (1620-30, sq. 31)

OSTRACODES:

*Occultocythereis* (?) sp. #CA (1590-1600, sq. 19)  
*Cytherella* #CA  
*Bairdia* #CA  
*Krithe* #CB

1650-1750m: Paleocene

*Globorotalia pseudomenardii* (1650-60, sq. 33)  
*Chiloguembelina crinita* (1680-90, sq. 8)

*Spiroplectammania spectabilis* (1650-60, sq. 32)  
*Textularia plummerae* (1650-60, sq. 21)  
*Glomospira charoides* (1650-60, sq. 9)

1765-1865m: Campanian

FORAMINIFERA

*Planoglobulina glabrata* (1765-75, sq. 32)  
*Pseudotextularia difformis* (1765-75, sq. 9)  
*Rosita fornicata* (1765-75, sq. 33)  
*Rosita contusa* (1765-75, sq. 45)  
*Abathomphalus mayaroensis* (1765-75, sq. 31)  
*Heterohelis punctulata* (1765-75, sq. 18)  
*Globotruncanella petaloidea* (1765-75, sq. 42)  
*Planoglobulina brazoensis* (1765-75, sq. 20)  
*Pseudotextularia elegans* (1795-1805, sq. 20)  
*Globotruncanella elevata* (1795-1805, sq. 8)  
*Globotruncanella stuartiformis* (1795-1805, sq. 32)  
*Marginotruncana marginata* (1795-1805, sq. 19)  
*Globotruncana bulloides* (1795-1805, sq. 9)  
*Globotruncana linneiana* (1855-65, sq. 32)  
*Bolivinoidea draco miliaris* (1765-75, sq. 7)  
*Neoflabellina rugosa leptodisca* (1765-75, sq. 8)  
*Stensolina exsculpta s.s.* (1765-75, sq. 22)  
*Bolivina incrassata gigantea* (1765-75, sq. 19)  
*Arenobulimina americana* (1765-75, sq. 6)  
*Stensolina excolata* (1765-75, sq. 30)  
*Stensolina pommerana* (1765-75, sq. 21)  
*Gaudryina rudita* (1795-1805, sq. 9)  
*Dorothyia bullata* (1795-1805, sq. 21)  
*Heterostomella americana* (1795-1805, sq. 34)  
*Heterostomella austinana* (1825-35, sq. 20)  
*Gavelinella cf. clementiana* (1855-65, sq. 21)  
*Gaudryina austinana* (1855-65, sq. 9)

1885-1895m: Santonian

FORAMINIFERA:

*Marginotruncana marginata*  
*Marginotruncana coronata* (1885-1895, sq. 44)  
*Marginotruncana pseudolinneiana* (1885-95, sq. 45)  
*Marginotruncana schneegansi* (1885-95, sq. 32)  
*Whiteinella baltica* (1885-95, sq. 21)  
*Dicarinella cf. concavata* (1885-95, sq. 9)  
*Dicarinella asymetrica* (1885-95, sq. 8)

*Lenticulina munsteri*  
*Stensoina exsculpta s.s.*

OSTRACODES:

*Cythereis* (?) sp. #CA (1885-95, sq. 20)

1915-2015: Cenomanian

FORAMINIFERA

*Praiglobotruncana stephani* (1915-25, sq. 8)  
*Heterohelix moremani* (1915-25, sq. 31)  
*Hedbergella delrioensis* (1915-25, sq. 7)  
*Rotalipora greenhornensis* (1945-55, sq. 43)  
*Rotolipora cf. deeckeii* (1945-55, sq. 42)  
*Gavelinella tourainensis* (1915-25, sq. 21)  
*Gavelinella cenomanica* (1915-25, sq. 20)  
*Dorothia af. filiformis* (1915-25, sq. 19)  
*Ammobaculites comprimatus* (1945-55, sq. 18)  
*Ammobaculites subcretaceus* (1945-55, sq. 19)  
*Marssonella trochus* (1945-55, sq. 8)  
*Quinqueloculina lirellangula* (1945-55, sq. 20)  
*Tritaxia pyramidata* (1945-55, sq. 30)  
*Quasispiroplectammina nuda* (1945-55, sq. 32)  
*Gavelinella ammonides* (1975-85, sq. 18)

OSTRACODES:

*Rhacythereis* sp #CA (1915-25, sq. 32)  
*Protocythere alexanderi* (1945-55, sq. 44)  
*Schuleridea jonesiana* (2005-2015, sq. 20)  
*Rhacythereis af. dentonensis* (2005-15, sq. 19)  
*Cythereis cf. eaglefordensis* (2005-15, sq. 7)

2035-2255m: Albian

FORAMINIFERA

*Ammobaculites goodlandensis*  
*Lenticulina gaultina*  
*Ammobaculites subcretaceus*  
*Ammobaculites comprimatus*  
*Ammobaculites reophacoides* (2245-55, sq. 6)



OSTRACODES:

*Cornicythereis* cf. *subgoodlandnesis* (2035-45, sq. 6)  
*Pontocyprrella* (?) sp. #CA (2035-45, sq. 17)  
*Protocythere* sp. #CA 2035-45, sq. 19)  
*Eocytheropteron* *paenorbiculatum* (2035-45, sq. 7)  
*Mandocythere* (?) sp. #CA (2035-45, sq. 18)  
*Protochthere* *speetonensis* (2065-75, sq. 7)  
*Rehacythereis* cf. *dentonensis* (2065-75, sq. 19)  
*Rehacythereis* af. *reticulata* (2125-35, sq. 7)  
*Cornicythereis* *bonnemai* (2125-35, sq. 18)  
*Cytherella* sp. #CC  
*Asciocythere* sp. #CA  
*Rehacythereis* sp. #CB  
*Eocytheropteron* sp. #CB

2275-3300: Aptian

FORAMINIFERA

*Favusella* *washitaensis* (2610-20, sq. 19)  
*Hedbergella* *trochoidea* (2610-20, sq. 8)  
*Hedbergella* *delrioensis* (2610-20, sq. 20)  
*Lenticulina* *nodosa* (2275-85, sq. 29)  
*Gavelinella* *intermedia* (2275-85, sq. 17)  
*Gavelinella* *berthelini* (2305-15, sq. 29)  
*Trochammina* sp. #CA (2335-45, sq. 18)  
*Cribrostomoides* sp. #CA  
*Recurvoides* sp. #CA  
*Bathysiphon* sp. #CB  
*Verneuulinoides* sp. #CA  
*Epistomina* *cretosa* (2550-60, sq. 18)  
*Gavelinella* *cenomanica*  
*Trocholina* af. *infragranulata* (2640-50, sq. 7)  
*Gavelinella* *barremiana* (2640-50, sq. 8)  
*Epistomina* cf. *carpenteri* 2640-50, sq. 6)  
*Tritaxia* *singularis* (2760-70, sq. 8)  
*Epistomina* *carpenteri* (2760-70, sq. 9)  
*Gyroidinoides* sp. #CA  
*Saracenaria* sp. #CA  
*Saracenaria* cf. *frankei* (2850-60, sq. 7)  
*Haplophagmium* (?) sp. #CA  
*Marssonella* cf. *oxsycona* (3150-60, sq. 19)

OSTRACODES:

*Mandocythere* *harrisiana* (2305-15)  
*Neocythere* cf. *mertensi*

Cornicythereis (?) sp. #CA  
Rehacythereis sp #CC  
Eocytheropteron sp. #CB  
Paracypris sp. #CA

3320-4790: Barremian

FORAMINIFERA

*Verneuilinoides neocomiensis* (3320-30, sq. 18)  
*Ammobaculites reophacoides*  
*Gavelinella barremiana*  
*Epistomina hechti* (3550-60, sq. 5)  
*Reophax minuta* (3550-60, sq. 17)  
*Gaudryinella tealbyensis* (3610-20, sq. 4)  
*Textularia foeda* (3700-10, sq. 2)  
*Discorbis* sp. #CA  
*Cancris* sp. #CA  
*Marsonella kummi* (4240-50, sq. 5)  
*Epistomina oronata* (4540-50, sq. 18)  
*Epistomina caracolla* (4540-50-60, sq. 6)  
*Lagena* sp. #CA  
*Marginuliopsis humilus* (4630-40, sq. 4)  
*Lenticulina* cf. *guttata* (4630-40, sq. 5)  
*Planularia crepidularis* (4780-90, sq. 14)

OSTRACODES:

*Asciocythere* sp. #CB

4810-5234m: Barremian or Hauterivian

FORAMINIFERA:

A few poorly preserved and undiagnostic agglutinating foraminifera.

[illegible]

[illegible]

## GEOLOGICAL SURVEY OF CANADA - QUALITATIVE FLUORESCENCE ANALYSIS SHEET

PAGE 1 OF 1

[illegible]