75 O

NOTES

The Artillery Lake map area consists of part of the western margin of the early Proterozoic Thelon Tectonic Zone of the northwestern Churchill Province and its linear contact with the Slave Province to the west. Linear gneisses and schists which have been juxtaposed against the Archean Slave Province with its curvilinear structural trends, relatively massive plutonic bodies and mainly low to intermediate metamorphic grade metasediments and metavolcanic rocks (Fig. 1). The east half of the area was mapped previously by Fraser (1972) and various aspects of the geology have been discussed in brief reports on the current mapping program following successive field seasons (Henderson and Macfie, 1985, 1986; Henderson et al., 1987). Geologically and for purposes of discussion the area can be divided into four structurally bounded domains (Figure 2).

Western Domain

The western domain is typical of much of the Slave Province in that it consists of steeply dipping Archean Yellowknife metasediments and metavolcanics ranging in grade from greenschist to amphibolite that have been intruded by a variety of Archean, mainly massive, equidimensional, bulbous plutons of varied composition.

A major unit of dominantly mafic Yellowknife volcanic rocks occurs along the northeastern side of Clinton-Colden Lake to east of Ptarmigan Lake. It is interrupted by plutonic intrusions between the two lakes. The unit consists mainly of mafic pillowed and massive flows that are locally amygdaloidal with lesser amounts of carbonate cemented mafic breccias and finer volcaniclastic sediments. Small felsic domes and volcaniclastic units are present locally but are not a major component of the unit. The whole unit has a U-Pb zircon age of 2673 +3/-2 Ma (van Breemen and Henderson, 1988). Most of the volcanics are at amphibolite grade although the volcanics on the large island in Clinton-Colden Lake are mainly at greenschist grade.

Yellowknife metasediments underlie a large area south and west of Clinton-Colden Lake. They consist of metamorphosed interbedded greywacke mudstone turbidites that are typical of the Yellowknife elsewhere in the Slave. With the exception of bedding and load grading, primary sedimentary structures are not well preserved. Most of the metasediments are at amphibolite grade and contain assemblages of the following minerals: quartz-plagioclase-biotite-muscovite-andalusite-sillimanite-staurolite-cordierite-garnet. Staurolite is more abundant than cordierite, in contrast to most other areas of the province. Greenschist grade rocks occur in the Lac de Gras area, Clinton-Colden Lake and west of Ptarmigan Lake.

Supracrustal rocks of the Yellowknife Supergroup are intruded by members of several plutonic suite. The Tarantula plutons are part of a widespread gneissic suite that ranges in composition from granodiorite in the northern part of the Healey Lake area to the north (Henderson and Thompson, 1982) to tonalite and diorite in this area. They have U-Pb zircon ages of 2622 + 1.4/-1.2 Ma and 2616 + 2/-6 Ma at localities in the Healey Lake and Artillery Lake areas respectively (van Breemen and Henderson, 1988; van Breemen et al, 1987b). The Charlotin granodiorite pluton, with a U-Pb zircon and sphene age of 2617 + 7/-5 Ma (van Breemen and Henderson, 1988) is similar to the Tarantula in terms of morphologically, compositionally, texturally and chronologically to the Defeat Pluteonic Suite at Yellowknife (Henderson, 1983). Other major plutonic units include the Artillery granite, a coarsely megacrystic unit that occurs mainly west of Artillery Lake but also in the central domain as well as in the Healey Lake area to the north. Ages determined on the Artillery granite include a U-Pb monazite age of 2596 + 3/-6 Ma (van Breemen and Henderson, 1988) and a U-Pb monazite and sphene age of 2596 + 1/0 Ma (van Breemen et al., 1987b). The two-mica Ptarmigan granite, which also occurs to the north, has a U-Pb monazite age of 2656 +/-10 (van Breemen and Henderson, 1988). It is similar in age, composition and texture to granites of the Yellowknife (Henderson, 1985). Several small pyroxenite intrusions less than 100 m in size occur in both this and the central domain to the east. They are similar to the line of somewhat larger intrusions north of Clinton-Colden Lake it the Healey Lake area (Henderson and Thompson, 1982). They are inferred to be more abundant than mapped on the basis of local concentrations of pyroxenite boulders in areas without an obvious source. A 3 km body of hornblende gabbro-anorthosite occurs east of the mafic volcanics at the east end of Clinton-Colden Lake (Macfie, 1987). Associated with it but mapped separately from it is a similarly sized diorite to quartz diorite body immediately to the south. A silli-like body of similar anorthositic gabbro occurs east of the mafic volcanics east of Ptarmigan Lake and a number of smaller unmapped ultramafic and recrystallized gabbroic anorthosite occurs within the central domain to the south.

Rock units of early Proterozoic age include the Artillery Lake formation that unconformably overlies Archean granitoid units at Artillery Lake (Gandhi, 1984, 1985; Rosoce et al., 1987). It is a locally stromatolitic dolomite containing, in places, minor lenses of quartz pebble conglomerate. Rare interbedded grey and red shale and dolomite are locally exposed but on the basis of the abundance of dolomite boulders in the drainage basin the bulk of the lake may well be underlain by shale. In the lake, the chains of islands underlain by dolomite follow the north-northeasterly regional fault pattern of the area and probably represent resistant fault scarps.

Two sets of early Proterozoic diabase dykes are present in the domain: the dominant set has an easterly trend while the second set trends east-northeasterly to northeasterly. Both sets occur in the Healey Lake area to the north where they become progressively metamorphosed towards the Thelon Tectonic Zone (Henderson and Thompson, 1982). The youngest unit in the area is the north-northeasterly trending, middle Proterozoic Mackenzie diabase dyke swarm. Only locally exposed, their mapped distribution is based on their strong magnetic signature.

At the map scale, structural trends are southerly in the central and eastern parts of the domain whereas in the northern part of the domain, trends follow the generally northwesterly structural trend of the eastern Slave. At more detailed scales the trends are commonly more irregular. In the low-grade rocks west of Ptarmigan Lake a pronounced horizontal crenulation is common. The relatively weak foliation becomes more intense with more consistent trends towards the eastern margin of the domain. Several north northeasterly trending faults and narrow shear zones that are parallel to the domain boundary structures and the abundant shear zones of the eastern domain, are present within the western domain.

Central Domain

The central domain differs from that to the west in that a moderate to strong foliation with a more consistent, generally north-northeasterly trend is ubiquitous. As well, parts of the domain have been variably affected by Proterozoic metamorphism. The Archean metamorphic grade is significantly higher than to the west on the basis of migmatite grade in the Yellowknife metasediments. The boundary with the western domain is a series of shear zones that in general become narrower to the north. Moderate to steep, down-dip, easterly plunging lineations occur locally in these zones. The northern half of the domain is also expressed by a background magnetic intensity at least 100 nanoteslas higher than that to the west (Geological Survey of Canada, 1987). A symmetric paired gravity anomaly characterizes the area between the Bathurst and McDonald faults (Fig. 1) and is influenced to some extent by the Archean metamorphic grade is significantly higher than to the west on the basis of migmatite grade in the Yellowknife metasediments. The boundary with the western domain is a series of shear zones that in general become narrower to the north. Moderate to steep, down-dip, easterly plunging lineations occur locally in these zones. The northern half of the domain is also expressed by a background magnetic intensity at least 100 nanoteslas higher than that to the west (Geological Survey of Canada, 1987). A symmetric paired gravity anomaly characterizes the area between the Bathurst and McDonald faults (Fig. 1) and is influenced to some extent by the central domain. The negative component occurs on the Slave Province side while the somewhat smaller positive anomaly is centered over the Thelon Tectonic Zone to the east but to some extent laps over the Slave Province to the north and south (Henderson et al., 1987). Within the map area the anomaly is sensitive to the central domain in that the general trend of the line of inflection between the positive and negative anomalies follows the western boundary of the domain in the southern part of the area before crossing over to the eastern boundary (Observatories Division, 1987). Another expression of the presence of the shift of the gravity anomaly pattern, the relatively higher background magnetic intensity, and the high metamorphic grade of the Yellowknife metasediments within it in contrast to the presence of similar but lower grade metasediments immediately to the north in the Healey Lake area. The domain as defined by these criteria does not continue far beyond the northern boundary of the map area although the bounding shear zone can be traced 40 km north to Healey Lake. This would suggest that if the central domain represents an uplifted crustal block exposing Slave Province rocks at a deeper structural level, the amount of uplift was greater in the south. Many of the Archean rock units recognized in the western domain, as the southern that are not present to the west, occur in the central domain.

The oldest rocks in the domain are believed to occur in the heterogeneous granitoid gneiss complex west and northwest of Sifton Lake. It contains compositionally layered, recrystallized granitic gneiss to mafic amphibolitic gneiss as well as foliated granitoid plutons that could be significantly younger. This complex is considered as a probable correlative with the Healey complex in the Healey Lake area to the north (Henderson and Thompson, 1982). There a meta-quartz diorite has a U-Pb zircon age of 2679 +/-3 Ma which falls between the zircon ages on Yellowknife volcanics in the region (2671 + 2/-4 Ma at Clinton Colden Lake and 2692 +/ 2 Ma 70 km to the northwest on the Back River (van Breemen and Henderson, 1988; van Breemen et al. 1987b). Two ages on the Healey complex granitic phases of the Healey complex to the east suggest ages close to 2.6 Ga (van Breemen et al., 1987b).

The metasediments within the domain are quartz-plagioclase-biotite-(muscovite-sillimanite-garnet) migmatites with varied proportions of leucosome, grading locally into areas of diatexite with only wisps and diffuse patches of relict metasediment. The major zone of metasedimentary rocks near the eastern boundary of the domain was mapped by Fraser (1972) and Gandhi (1988) with no hint of an Archean component. Particularly common within the granitoid gneisses in a zone between Morain Lake and the Hanbury River but occuring elsewhere as well are a series of narrow bodies ranging in composition from black gabbro through anorthositic gabbro to rarely white anorthosite. These bodies have been illustrated on the map as lenses but may well be more extensive along strike than indicated.

Two granitic units differ from the previously discussed major units of the domain in that they form discrete bodies that are significantly younger and do not appear to have undergone granulite grade metamorphism. The Crichchil granite forms a prominent pluton in the northeast part of the area. Foliated pink and black megacrystic granite that than normal for the Archean low-pressure, high-temperature facies-series typical of the Slave Province, its growth outlasted presumed Proterozoic shear structures (James, 1985) and it is close to the very high grade Proterozoic metamorphic rocks of the eastern domain. Within the major shear zone in the eastern belt of metasediment, migmatite layering becomes completely disrupted to form a relatively homogeneous porphyroclastic rock.

A major zone of amphibolite up to 3 km wide occurs within the eastern belt of Yellowknife rocks north of Sifton Lake. The rocks consist of compositionally and texturally layered amphibolites to migmatitic amphibolites with variably-developed, white, biotite-bearing leucosome. Thin metasedimentary units are locally present but rare. Although no primary textures or structures are preserved, this unit, because of its thickness, mafic composition and association with the other metasedimentary migmatites, is thought to be derived from a volcanic unit. West of Sifton Lake the metasedimentary migmatites contain abundant thin, silli-like amphibolite bodies that may represent more or less syn-volcanic mafic intrusives into the sediments as is seen elsewhere in the Slave Province. Other small mafic units along with rare carbonate lenses occur to the south within this supracrustal belt.

Granitoid intrusive rocks make up a major proportion of the domain. All contain a variably developed foliation. The oldest unit is the Sifton granitoid complex in the eastern part of the domain. It is a grey, medium-grained to coarse, megacrystic, biotite granodiorite to tonalite with varied proportions of lenssets veins of pink to white, mafic-poor, generally coarser-grained leucosome that is commonly gradational with the grey phase. The grey tonalitic phase has a U-Pb age of 2661 +/-2 Ma (van Breemen and Henderson, 1988). To the north the Muscliw granite forms a large, homogeneous, foliated, pink and black megacrystic granite that has a U-Pb zircon age of 2605 +3/-2 Ma (van Breemen and Henderson, 1988). Both the Smart and Muscliw granites have close but typically more deformed lithological analogues in the eastern domain although U-Pb geochronological data indicate that they are distinctly younger. More strongly foliated and locally migmatitic phases of the Smart granite and two-mica Ptarmigan granite of the western domain extend across the boundary faults into the central domain. The remaining granitoid rocks have been left in an undifferentiated unit that consists of an heterogeneous assemblage of foliated granite to tonalite and includes dioreitic derived from the Yellowknife metasediment. Sound, where the rocks of the domain are strongly deformed, it is commonly difficult to differentiate between porphyroclastic gneisses derived from Yellowknife migmatites and biotite-rich meta-tonalites.

The same sets of Proterozoic diabase dykes present in the western domain also occur in the central zone. The northeasterly to east-northeasterly set tends to have a more northeasterly trend and dominates in this domain. The trend of the easterly set is somewhat variable but is generally east-northeast to more northeasterly trend of the other set is said due to rotation by internal deformation of the domain but may be more a response to the pre-existing foliation in the domain. Both dyke sets become metamorphosed and foliated to some degree toward the Thelon Tectonic Zone, as was seen to the north in the Healey Lake area (Henderson and Thompson, 1982). Dykes in the southern part of the central zone are less metamorphosed than those to the north suggesting that there is also a north-south Proterozoic metamorphic gradient as well.

Eastern Domain

The eastern domain, representing part of the Thelon Tectonic Zone, is bounded by faults with the western and central domains of the Slave province. Despite this, the boundary with the central domain is the more poorly defined of any of the domain boundaries, being a shear zone between the somewhat more homogeneous, deformed granitoids and gneisses on the Slave side and the somewhat more heterogeneous, deformed granitoid gneisses of the Sifton gneiss of the Thelon Tectonic Zone. It is best defined near the southern border of the map area where James (1985), on the basis of detailed mapping, placed the boundary about 1.5 km east of the zone of Yellowknife supracrustal rocks.

The eastern domain consists of a heterogeneous assemblage of severely deformed granitoid and supracrustal gneisses, most of which are, or have been retrograded from, granulite metamorphic grade. Mylonitic shear zones occur throughout the domain, resulting in a generally much more linear map pattern relative to the domains to the west. The domain has been divided into several subdivisions, that are commonly bounded by pronounced, NNE striking, shear zones. The subdivision may have differing lithological proportions, locally contrasting structural trends and style and, more rarely, contrasting psphysical expression. These shear zones are similar in trend to those that bound the central domain and form the major shear zone on the east side of the central domain (the 'straight zone' (Henderson and Thompson, 1982), and are parallel as well to the Sifton Lake shear zones that occur within the western domain.

The eastern domain is divided into five major units: two gneiss units of presumed supracrustal origin, a heterogeneous layered gneiss and two granitoid gneiss units. In many cases the rock packages are totally gradational such that the unit designation indicates the probable dominant rock type present in the region with all or any of the other rock types also possibly present. Both the supracrustal and granitoid gneisses are commonly migmatitic with highly varied proportions of in-situ leucosome together with introduced granitoid phases. The common occurrence of orthopyroxene indicates that much of the domain is at granulite metamorphic grade. This is particularly true in the area immediately north of Mary Frances Lake where the occurrence of orthopyroxene in rocks of appropriate composition is rather sporadic. Reconnaissance geothermobarometric measurements in the domain suggest a pressure of 7.4 kbar at 780°C for a relatively undeformed sample while several pyroxene/mylonite samples gave significantly lower values between 4 and 5 kbar at 700°C (Haggart, 1985).

Migmatite consisting of biotite-rich paleosome and white quartz-plagioclase leucosome of presumed metasedimentary parentage dominates in the southern part of the domain. An introduced pink granitic phase is commonly present at all scales. Rarely do the migmatites contain sillimanite but where it is present, the migmatites resemble the high-grade migmatites in the western part of the central domain. More commonly however, hornblende and clinopyroxene are present in the biotite-rich rocks, suggesting a more calcic parent. As a result, previous correlations of this unit with the Yellowknife (Fraser, 1972; Henderson and Macfie, 1985) is in some doubt. A possible correlative occurs at lower grade south of the McDonald fault, 150 km WSW of the most westerly occurrence of the biotite migmatite in the Artillery Lake area. There the 'external paragneisses' of Hanmer and Needham (1988) consist of 'micaceous meta-arenite, rhythmically interlayered meta-arenite and metapelite, calc-silicate and amphibolite' of which the calcsilicate forms a major component. Associated with it are 'internal paragneisses' that consist of meta-arenite and extensive units of felsic gneisses, amphibole-rich rocks ranging from amphibolites to amphibolitic gneisses and amphibolitic migmatites that may represent mafic to intermediate volcanics in a supracrustal succession. Like the biotite migmatites, an introduced, concordant granitic component is common. No geochronological data on the supracrustal gneisses are presently available.

The largest unit in the domain is a heterogeneous assemblage of undifferentiated layered gneisses. A large component consists of quartzofeldspathic amphibole-bearing gneisses, possibly of supracrustal origin. Sheets and veins of granitoid gneiss occur throughout the unit. Some into the layered gneiss, south and east of Morain Lake is a series of discontinuous layers and lenses of granitoid gneiss commonly associated with these gneisses. Immediately north of the area at Morraine Lake, gneisses dated at two localities indicate an age of metamorphism of between 1950 and about 1990 Ma (U-Pb on monazite and zircon), with evidence of the protolith of one of the gneisses being no older than 2400 Ma (U-Pb on zircon; van Breemen et al., 1987a). The age of metamorphic zircon from a granulite-grade amphibolite sill within gneisses on the Hanbury River east of Sifton Lake support an age of less that 2000 Ma for the metamorphism (van Breemen et al., 1987a).

Units dominated by compositionally and texturally heterogeneous granitoid gneisses to migmatitic gneisses occur in several zones across the domain. The Sifton gneiss along the west margin of the domain is one such unit. It is a foliated pink and grey layered gneiss with locally developed coarse feldspar and finer black hornblende porphyroclasts; where best preserved it can be seen to be derived from a granitoid migmatite not unlike the Archean Smart granitoid migmatite of the central zone. However, a fairly concordant U-Pb zircon age of about 1975 Ma has been interpreted as a minimal age of metamorphism and migmatization of the Sifton gneiss (van Breemen and Henderson, 1988) with no hint of an Archean component. Particularly common within the granitoid gneisses in a zone between Morain Lake and the Hanbury River but occuring elsewhere as well are a series of narrow bodies ranging in composition from black gabbro through anorthositic gabbro to rarely white anorthosite. These bodies have been illustrated on the map as lenses but may well be more extensive along strike than indicated.

Two granitic units differ from the previously discussed major units of the domain in that they form discrete bodies that are significantly younger and do not appear to have undergone granulite grade metamorphism. The Crichchil granite forms a prominent pluton in the northeast part of the area. Foliated pink and black megacrystic granite that than normal for the Archean low-pressure, high-temperature facies-series typical of the Slave Province, its growth outlasted presumed Proterozoic shear structures (James, 1985) and it is close to the very high grade Proterozoic metamorphic rocks of the eastern domain. Within the major shear zone in the eastern belt of metasediment, migmatite layering becomes completely disrupted to form a relatively homogeneous porphyroclastic rock.

A major zone of amphibolite up to 3 km wide occurs within the eastern belt of Yellowknife rocks north of Sifton Lake. The rocks consist of compositionally and texturally layered amphibolites to migmatitic amphibolites with variably-developed, white, biotite-bearing leucosome. Thin metasedimentary units are locally present but rare. Although no primary textures or structures are preserved, this unit, because of its thickness, mafic composition and association with the other metasedimentary migmatites, is thought to be derived from a volcanic unit. West of Sifton Lake the metasedimentary migmatites contain abundant thin, silli-like amphibolite bodies that may represent more or less syn-volcanic mafic intrusives into the sediments as is seen elsewhere in the Slave Province. Other small mafic units along with rare carbonate lenses occur to the south within this supracrustal belt.

Granitoid intrusive rocks make up a major proportion of the domain. All contain a variably developed foliation. The oldest unit is the Sifton granitoid complex in the eastern part of the domain. It is a grey, medium-grained to coarse, megacrystic, biotite granodiorite to tonalite with varied proportions of lenssets veins of pink to white, mafic-poor, generally coarser-grained leucosome that is commonly gradational with the grey phase. The grey tonalitic phase has a U-Pb age of 2661 +/-2 Ma (van Breemen and Henderson, 1988). To the north the Muscliw granite forms a large, homogeneous, foliated, pink and black megacrystic granite that has a U-Pb zircon age of 2605 +3/-2 Ma (van Breemen and Henderson, 1988). Both the Smart and Muscliw granites have close but typically more deformed lithological analogues in the eastern domain although U-Pb geochronological data indicate that they are distinctly younger. More strongly foliated and locally migmatitic phases of the Smart granite and two-mica Ptarmigan granite of the western domain extend across the boundary faults into the central domain. The remaining granitoid rocks have been left in an undifferentiated unit that consists of an heterogeneous assemblage of foliated granite to tonalite and includes dioreitic derived from the Yellowknife metasediment. Sound, where the rocks of the domain are strongly deformed, it is commonly difficult to differentiate between porphyroclastic gneisses derived from Yellowknife migmatites and biotite-rich meta-tonalites.

The same sets of Proterozoic diabase dykes present in the western domain also occur in the central zone. The northeasterly to east-northeasterly set tends to have a more northeasterly trend and dominates in this domain. The trend of the easterly set is somewhat variable but is generally east-northeast to more northeasterly trend of the other set is said due to rotation by internal deformation of the domain but may be more a response to the pre-existing foliation in the domain. Both dyke sets become metamorphosed and foliated to some degree toward the Thelon Tectonic Zone, as was seen to the north in the Healey Lake area (Henderson and Thompson, 1982). Dykes in the southern part of the central zone are less metamorphosed than those to the north suggesting that there is also a north-south Proterozoic metamorphic gradient as well.

Eastern Domain

The eastern domain, representing part of the Thelon Tectonic Zone, is bounded by faults with the western and central domains of the Slave province. Despite this, the boundary with the central domain is the more poorly defined of any of the domain boundaries, being a shear zone between the somewhat more homogeneous, deformed granitoids and gneisses on the Slave side and the somewhat more heterogeneous, deformed granitoid gneisses of the Sifton gneiss of the Thelon Tectonic Zone. It is best defined near the southern border of the map area where James (1985), on the basis of detailed mapping, placed the boundary about 1.5 km east of the zone of Yellowknife supracrustal rocks.

The eastern domain consists of a heterogeneous assemblage of severely deformed granitoid and supracrustal gneisses, most of which are, or have been retrograded from, granulite metamorphic grade. Mylonitic shear zones occur throughout the domain, resulting in a generally much more linear map pattern relative to the domains to the west. The domain has been divided into several subdivisions, that are commonly bounded by pronounced, NNE striking, shear zones. The subdivision may have differing lithological proportions, locally contrasting structural trends and style and, more rarely, contrasting psphysical expression. These shear zones are similar in trend to those that bound the central domain and form the major shear zone on the east side of the central domain (the 'straight zone' (Henderson and Thompson, 1982), and are parallel as well to the Sifton Lake shear zones that occur within the western domain.

The eastern domain is divided into five major units: two gneiss units of presumed supracrustal origin, a heterogeneous layered gneiss and two granitoid gneiss units. In many cases the rock packages are totally gradational such that the unit designation indicates the probable dominant rock type present in the region with all or any of the other rock types also possibly present. Both the supracrustal and granitoid gneisses are commonly migmatitic with highly varied proportions of in-situ leucosome together with introduced granitoid phases. The common occurrence of orthopyroxene indicates that much of the domain is at granulite metamorphic grade. This is particularly true in the area immediately north of Mary Frances Lake where the occurrence of orthopyroxene in rocks of appropriate composition is rather sporadic. Reconnaissance geothermobarometric measurements in the domain suggest a pressure of 7.4 kbar at 780°C for a relatively undeformed sample while several pyroxene/mylonite samples gave significantly lower values between 4 and 5 kbar at 700°C (Haggart, 1985).

Migmatite consisting of biotite-rich paleosome and white quartz-plagioclase leucosome of presumed metasedimentary parentage dominates in the southern part of the domain. An introduced pink granitic phase is commonly present at all scales. Rarely do the migmatites contain sillimanite but where it is present, the migmatites resemble the high-grade migmatites in the western part of the central domain. More commonly however, hornblende and clinopyroxene are present in the biotite-rich rocks, suggesting a more calcic parent. As a result, previous correlations of this unit with the Yellowknife (Fraser, 1972; Henderson and Macfie, 1985) is in some doubt. A possible correlative occurs at lower grade south of the McDonald fault, 150 km WSW of the most westerly occurrence of the biotite migmatite in the Artillery Lake area. There the 'external paragneisses' of Hanmer and Needham (1988) consist of 'micaceous meta-arenite, rhythmically interlayered meta-arenite and metapelite, calc-silicate and amphibolite' of which the calcsilicate forms a major component. Associated with it are 'internal paragneisses' that consist of meta-arenite and extensive units of felsic gneisses, amphibole-rich rocks ranging from amphibolites to amphibolitic gneisses and amphibolitic migmatites that may represent mafic to intermediate volcanics in a supracrustal succession. Like the biotite migmatites, an introduced, concordant granitic component is common. No geochronological data on the supracrustal gneisses are presently available.

The largest unit in the domain is a heterogeneous assemblage of undifferentiated layered gneisses. A large component consists of quartzofeldspathic amphibole-bearing gneisses, possibly of supracrustal origin. Sheets and veins of granitoid gneiss occur throughout the unit. Some into the layered gneiss, south and east of Morraine Lake is a series of discontinuous layers and lenses of granitoid gneiss commonly associated with these gneisses. Immediately north of the area at Morraine Lake, gneisses dated at two localities indicate an age of metamorphism of between 1950 and about 1990 Ma (U-Pb on monazite and zircon), with evidence of the protolith of one of the gneisses being no older than 2400 Ma (U-Pb on zircon; van Breemen et al., 1987a). The age of metamorphic zircon from a granulite-grade amphibolite sill within gneisses on the Hanbury River east of Sifton Lake support an age of less that 2000 Ma for the metamorphism (van Breemen et al., 1987a).

Units dominated by compositionally and texturally heterogeneous granitoid gneisses to migmatitic gneisses occur in several zones across the domain. The Sifton gneiss along the west margin of the domain is one such unit. It is a foliated pink and grey layered gneiss with locally developed coarse feldspar and finer black hornblende porphyroclasts; where best preserved it can be seen to be derived from a granitoid migmatite not unlike the Archean Smart granitoid migmatite of the central zone. However, a

Economic Geology

Lead mineralization is associated with the contact between the east's Proterozoic Artillery Lake dolomite and the underlying granitoid basement at Artillery Lake. The structurally controlled, epigenetic veins are localized along fractures and faults formed during folding and fracturing of the Artillery Lake formation (Gandhi, 1984, 1985; Rosoce et al., 1987). Rare occurrences of low grade nickel-copper mineralization have been reported in the western Thelon Tectonic Zone in the vicinity of Smart Lake – Campbell Lake (Thorpe, 1972; Padgham et al, 1976). This discovery resulted in the staking of several thousand claims in the region together with considerable airborne and ground EM and magnetometer surveys, geological mapping and some drilling. Although no significant mineralization was found, numerous EM conductors were identified, for the most part, can be related to graphitic bearing metasediments (Fraser, 1972; McGrath and Henderson, 1985). Silicate-sulphide facies iron formation was recognized at one locality in the southern domain at the eastern end of Mary Frances Lake (63°16.9' 106°04.9'). It occurs at the contact between migmatite grade biotite rich metasediments and amphibole rich metavolcanics in association with thin marble and calc silicate units. Lithium mineral bearing pegmatites occur at two localities east of Clinton-Colden Lake, one at 63°51.2' 107°31.1' and the other just north of the border at 64°01.2' 107°48.7'. These pegmatites may be related to the two-mica Ptarmigan granites which occur in the immediate area and may be similar to the lithium mineral bearing pegmatites associated with the two-mica granites in the Healey Lake area to the Yellowknife. A small occurrence of graphite occurs in the eastern part of the domain at Clinton-Colden Lake (Sexton, 1983). Associated with the same volcanic unit but in the Healey Lake area to the north is an occurrence of sphalerite boulders whose source is as yet unidentified (Gibbins et al., 1977).

References

Fraser, J.A.
1972: Artillery Lake map area, District of Mackenzie; Geological Survey of Canada, Paper 71-28, 17 p.

Frith, R.A.
1982: Geology of the Beechey Lake-Duggan Lake Map Area, 76G, E/2 & 76H, W/2; Geological Survey of Canada, Open File 831.

Gandhi, S.S.
1984: Cartra-sphalerite-chalcopyrite veins in Aphebian dolomite and Archean basement at Artillery Lake, Northwest Territories; in Current Research, Part B, Geological Survey of Canada, Paper 84-1B, p. 33-44.

Gandhi, S.S.
1985: Geology of the Artillery Pb-Zn-Cu district, Northwest Territories; in Current Research, Part C, Geological Survey of Canada, Paper 85-1A, p. 359-363.

Geological Survey of Canada
1968: Artillery Lake, District of Mackenzie, Northwest Territories, Geological Survey of Canada, Aeromagnetic Series Map 21502.

Gibbins, W.A., Sexton, J.B., Lapeyre, F.J.,
Murphy, J.D., Hurdle, E.J., Padgham, W.A.
1977: Mineral Industry Report 1974, Northwest Territories; Department of Indian and Northern Affairs, Canada, 267 p.

Haggart, M. J.
1987: Geothermobarometry of the Slave/Churchill Structural Provinces boundary, Artillery Lake map area, District of Mackenzie; unpublished B.Sc. thesis, Department of Geological Sciences, Queen's University, Kingston.

Henderson, J.B.
1975: Sedimentological studies of the Proterozoic Healey Formation in the Slave Structural Province; Geological Research, Part A, Geological Survey of Canada, Paper 75-1A, p. 325-330.

Henderson, J.B.
1985: Geology of the Yellowknife-Hearne Lake area, District of Mackenzie: a segment across the Archean basin; Geological Survey of Canada, Memoir 414.

Henderson, J.B. and Macfie, R.I.
1985: The northern Artillery Lake area: a transect across the Thelon Front; in Current Research, Part A, Geological Survey of Canada, Paper 85-1A, p. 441-448.

Henderson, J.B. and Macfie, R.I.
1986: Artillery Lake map area, District of Mackenzie: a transect across the Thelon Front; in Current Research, Part A, Geological Survey of Canada, Paper 86-1A, p. 107-116.

Henderson, J.B., McGrath, P.H., James, D.T. and Macfie, R.I.
1987: An integrated geological, gravity and magnetic study of the Artillery Lake area and the Thelon Tectonic Zone, District of Mackenzie; in Current Research, Part A, Geological Survey of Canada, Paper 87-1A, p. 203-814.

Henderson, J.B. and Thompson, P.H.
1982: Geology, Healey Lake, District of Mackenzie, Northwest Territories; Geological Survey of Canada, Open File 859.

James, D.T.
1985: Geology of the Moraine Lake area and the Thelon Front, District of Mackenzie; in Current Research, Part A, Geological Survey of Canada, Paper 85-1A, p. 645-652.

Macfie, R.I.
1987: The Clinton-Colden hornblende gabbro-anorthosite intrusion, Artillery Lake map area, District of Mackenzie; in Current Research, Part A, Geological Survey of Canada, Paper 87-1A, p. 691-697.

McGrath, P.H. and Henderson, J.B.
1985: Reconnaissance ground magnetic and VLF profile data in the vicinity of the Thelon Front, Artillery Lake map area, District of Mackenzie; in Current Research, Part A, Geological Survey of Canada, Paper 85-1A, p. 649-652.

Miller, S.M.
1987: Kinematic analysis and metamorphic character of a shear zone in the Thelon Front, Artillery Lake area, District of Mackenzie, N.W.T.; unpublished B.Sc. thesis, Department of Geology, McMaster University, Hamilton.

Observatories Division
1969: Artillery Lake, Northwest Territories; Observatories Division, Gravity Map Series No. 39.

Padgham T., Caine, T.W., Hughes, D.R., Jefferson, C.W., Kennedy, M.W., Murphy, J.D.
1978: Mineral Industry Report, 1969 and 1970, Northwest Territories, West of 104°W Longitude, Department of Indian and Northern Affairs, Canada, v. 3, 199 p.

Rocoe, S.M., Gandhi, S.S., Charbonneau, B.W., Maurice, Y.T., and Gibb, R.A.
1987: Mineral resource assessment of the area in the East Arm (Great Slave Lake) and Artillery Lake region, N.W.T., proposed as a national park (NTS 75J,K,L,N,O); Geological Survey of Canada, Open File 1434, 93p.

Seaton, J.B.
1983: Slave Structural Province; in Mineral Industry Report 1972, Northwest Territories, ed. J.A. Brophy, Indian and Northern Affairs, Canada, chapter 7, p. 217-244.

Thomas, M.D., Gibb, R.A. and Quince, J.A.
1976: New evidence from offset aeromagnetic anomalies for transcurrent faulting associated with the Bathurst and McDonald faults, Northwest Territories; Canadian Journal of Earth Sciences, v. 13, p. 1246-1260.

Thompson, P.H.
1986: Geology, Tinney Hills-Overby Lake (West Half), District of Mackenzie, Northwest Territories; Geological Survey of Canada, Open File 1316.

Thorpe R.I.
1972: Mineral exploration and mining activities, mainland Northwest Territories, 1966 to 1968, (excluding the Coppermine River area); Geological Survey of Canada, Paper 70-39, 204 p.

van Breemen, O. and Henderson, J.B.
1988: U-Pb zircon and monazite ages from the eastern Slave Province and Thelon Tectonic Zone, Artillery lake area, N.W.T.; in Radiogenic Age and Isotopic Studies: Report 2, Geological Survey of Canada, Paper 88-2, (in press).

van Breemen, O., Henderson, J.B. Loveridge, W.D. and Thompson, P.H.
1987a: U-Pb zircon and monazite geochronology and evolution of granitoid and gneissic rocks in the Thelon Tectonic Zone, Healey Lake and Artillery Lake map areas, N.W.T.; in Current Research, Geological Survey of Canada, Paper 87-2, p. 103-110.

van Breemen,O., Henderson,J.B., Sullivan,R.W. and Thompson,P.H.
1987b: U-Pb zircon and monazite ages from the eastern Slave Province, Healey Lake area, N.W.T.; in Radiogenic Age and Isotopic Studies: Report 1, Geological Survey of Canada, Paper 87-2, p. 107-110.

Wodicka, N.
1987: Geology and geochemistry of part of the Clinton-Colden greenstone belt, N.W.T.; unpublished B.Sc. thesis, Department of Geology, University of Ottawa, Ottawa, 85 p.



ARTILLERY LAKE

DISTRICT OF MACKENZIE

NORTHWEST TERRITORIES

LEGEND

WESTERN and CENTRAL DOMAINS

Middle Proterozoic

- Mackenzie Dykes: diabase, gabbro, in most cases position of dykes defined by aeromagnetic expression, also occur in both eastern and southern domains but not in part obscured by extreme variation of magnetic pattern in those domains

Early Proterozoic

- Artillery Lake formation: dolomite, pale yellow-brown to orange weathering, locally stromatolitic and intraformational conglomeratic, locally in lens of minor, fine, well rounded quartz pebble conglomerate

Archean

- Mafic dykes: diabase, gabbro to amphibolite, occurs in an easterly, subordinately to a west-northwesterly and northerly trending sets, dykes in the western domain are massive and only weakly altered while those in the central domain, in particular the northern half, tend to be metamorphosed and foliated to varied degrees

The following five units represent 'probability' units as opposed to the more clearly defined units described above in that the lithology indicated by the unit descriptor is the rock type generally present at any given locality. However, smaller quantities of the other four units may also be present.

- Ptarmigan granite: buff to white to pale yellow, leucocratic, medium to coarse grained, locally megacrystic, equigranular, contains moderate amounts of biotite and muscovite in varied proportions; muscovite bearing pegmatite locally present
- Smart granitoid migmatite: heterogeneous migmatitic granitoid, medium to coarse grained, grey, locally pink; biotite and/or hornblende granodiorite to tonalite with lesser amounts of locally defined leucocratic to pink, coarser grained, leucocratic, granitic leucosome; moderately foliated with local protomylonite zones
- Muscliw granite: pink to red and dark grey to black, homogeneous, weakly to moderately foliated, inequigranular granite with coarse microcline megacrysts in a biotite, biotite-rich matrix
- Charlotin granodiorite: granodiorite to tonalite, pinkish grey to white, massive, medium grained, equigranular, biotite, locally hornblende bearing
- Pyroxenite: black, massive, medium to coarse grained; occurs as small plugs or lenses
- Tarantula tonalite: dominantly quartz diorite but ranging from tonalite to gabbro, massive, massive, medium to coarse grained, equigranular, hornblende and biotite bearing

Undifferentiated granitoid rocks: heterogeneous assemblage of granite to diorite to granitoid gneiss, moderately to strongly foliated, includes in part dioreitic derived from Yellowknife metasediment

Anorthositic gabbro: hornblende gabbro to leucogabbro to anorthosite, dark green to grey to white, massive, heterogenous; extreme grain size variation

Diorite: diorite to quartz diorite to gabbro, dark grey, massive, medium to fine grained

Yellowknife Supergroup

- Greywacke, siltstone, mudstone; greenschist grade greywacke-siltstone turbidites interbedded with graphitic mudstone, sedimentary textures and structures locally preserved
- Schist, anamictic to pelitic: amphibolite grade equivalent of above, contains various assemblages of quartz-plagioclase-biotite-muscovite-garnet-staurolite-andalusite-sillimanite (kyanite locally in northeastern metasediments)
- Migmatite: migmatitic equivalent of above
- Felsic volc